

$a = 11.1641 (4) \text{ \AA}$
 $b = 12.2744 (5) \text{ \AA}$
 $c = 13.2265 (5) \text{ \AA}$
 $\alpha = 106.924 (2)^\circ$
 $\beta = 99.622 (3)^\circ$
 $\gamma = 103.452 (2)^\circ$
 $V = 1631.79 (11) \text{ \AA}^3$
 $Z = 1$
 $\text{Mo } K\alpha \text{ radiation}$
 $\mu = 1.05 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
 $0.26 \times 0.22 \times 0.20 \text{ mm}$

Hexakis(μ_2 -4-amino-3,5-dimethyl-4H-1,2,4-triazole)hexaaquatricobalt(II) naphthalene-1,5-disulfonate tetrachloride

Zhi-Guo Gu,* Bao-Xiang Wang, Chun-Yan Pang and Fei-Fei Bao

School of Chemical and Material Engineering, Jiangnan University, Wuxi 214122, People's Republic of China
Correspondence e-mail: zhiguogu@jiangnan.edu.cn


Received 29 March 2012; accepted 7 May 2012

Key indicators: single-crystal X-ray study; $T = 100 \text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005 \text{ \AA}$; disorder in solvent or counterion; R factor = 0.048; wR factor = 0.108; data-to-parameter ratio = 16.6.

In the centrosymmetric trinuclear cation of the title compound, $[\text{Co}_3(\text{C}_4\text{H}_8\text{N}_4)_6(\text{H}_2\text{O})_6](\text{C}_{10}\text{H}_6\text{O}_6\text{S}_2)\text{Cl}_4$, the three Co^{II} atoms are bridged by six triazole molecules via the N atoms in the 1,2-positions. The central Co^{II} atom, lying on an inversion center, is coordinated by six triazole N atoms while the terminal Co^{II} atoms are coordinated by three triazole N atoms and three water O atoms in a distorted octahedral geometry. The naphthalenedisulfonate anion is also centrosymmetric. The four chloride counter anions are half-occupied; the H atoms of the amino groups show an occupancy of 2/3. O—H···Cl, O—H···O and N—H···O hydrogen bonds link the complex cations and the chloride and naphthalene-1,5-disulfonate anions.

Related literature

For the structure of the title cation as hydrated nitrate salt, see: Tong *et al.* (2011).



Experimental

Crystal data

$[\text{Co}_3(\text{C}_4\text{H}_8\text{N}_4)_6(\text{H}_2\text{O})_6](\text{C}_{10}\text{H}_6\text{O}_6\text{S}_2)\text{Cl}_4$

$M_r = 1385.82$
Triclinic, $P\bar{1}$

Data collection

Bruker SMART APEX CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2001)
 $T_{\min} = 0.763$, $T_{\max} = 0.815$

23481 measured reflections
6410 independent reflections
5057 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.108$
 $S = 1.06$
6410 reflections

386 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.67 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.75 \text{ e } \text{\AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1W—H1X···Cl3 ⁱ	0.98	2.06	2.765 (3)	127
O1W—H1Y···Cl3 ⁱⁱ	0.98	1.89	2.701 (3)	138
O2W—H2X···Cl1	0.98	2.28	2.929 (3)	122
O2W—H2Y···Cl3 ⁱⁱ	0.98	2.82	3.749 (3)	158
O3W—H3X···O2 ⁱⁱⁱ	0.98	1.89	2.773 (3)	148
O3W—H3Y···Cl4 ⁱⁱⁱ	0.98	2.18	3.114 (3)	159
N8—H8A···O3 ⁱ	0.91	2.21	3.009 (3)	146

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x, y, z - 1$; (iii) $-x + 2, -y + 1, -z + 1$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *SHELXTL*.

This work was supported by the National Natural Science Foundation of China (grant No. 21101078), the Program for New Century Excellent Talents in Universities of China (NCET-11-0657), the Natural Science Foundation of Jiangsu Province (BK2011143), the Fundamental Research Funds for the Central Universities (JUSRP21111), the State Key Laboratory of Coordination Chemistry of Nanjing University and the Scientific Research Foundation of Jiangnan University (20101951).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2535).

References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Tong, Y.-Z., Wang, Q.-L., Si, M., Qi, J., Yan, S.-P., Yang, G.-M., Cheng, P. & Liao, D.-Z. (2011). *Polyhedron*, **30**, 3151–3157.

supporting information

Acta Cryst. (2012). E68, m862 [https://doi.org/10.1107/S1600536812020417]

Hexakis(μ_2 -4-amino-3,5-dimethyl-4*H*-1,2,4-triazole)hexaaquaticobalt(II) naphthalene-1,5-disulfonate tetrachloride

Zhi-Guo Gu, Bao-Xiang Wang, Chun-Yan Pang and Fei-Fei Bao

S1. Comment


A recent study has reported a centrosymmetric trinuclear cobalt nitrate complex of 4-amino-3,5-dimethyl-1,2,4-triazole ($C_4H_8N_4$), which crystallizes in the monoclinic system, $P2_1/n$ space group (Tong *et al.*, 2011). The charge of the hexavalent cation, $[Co_3(C_4H_8N_4)_6(H_2O)_6]^{6+}$, is balanced by the nitrate anions. In the trinuclear unit of the title compound, the central Co^{II} atom and two terminal Co^{II} atoms are bridged by six triazole ligands (Fig. 1). The central Co^{II} atom, lying on an inversion center, is coordinated by six N atoms from the triazole ligands whereas each terminal Co^{II} atom completes its octahedral geometry by three water molecules. The title compound displays a similar structure to the previously reported cobalt analogs (Tong *et al.*, 2011), but it crystallizes in the triclinic system, $P\bar{1}$ space group. The charge of the trinuclear cation is balanced by one naphthalene-1,5-disulfonate dianion and four chloride anions. O—H···Cl, O—H···O and N—H···O hydrogen bonds link the complex cations, chloride and naphthalene-1,5-disulfonate anions (Table 1).

S2. Experimental

A 20 ml ethanol solution of 4-amino-3,5-dimethyl-1,2,4-triazole (0.6 mmol) was mixed with a 10 ml aqueous solution of $CoCl_2 \cdot 6H_2O$ (0.3 mmol), naphthalene-1,5-disulfonate acid (0.1 mmol) in 5 ml of ethanol was added to this reaction mixture with continuous stirring. The resulting solution was filtered and left to stand at room temperature. Pink bolck-shaped crystals of the title compound were obtained by slow evaporation of the solvent within two weeks (yield: 48%).

S3. Refinement

H atoms on water molecules and on N8 atom were located from a difference Fourier map and refined as riding atoms, with $U_{iso}(H) = 1.2U_{eq}(O)$ or $1.5U_{eq}(N)$. The other H atoms were positioned geometrically and refined as riding atoms, with C—H = 0.95 (CH) and 0.98 (CH₃) and N—H = 0.88 Å and with $U_{iso}(H) = 1.2(1.5$ for methyl) $U_{eq}(C, N)$. Cl atoms were refined with a fixed occupancy factor of 0.50.

**Figure 1**

The molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level.
[Symmetry codes: (i) 1-x, 1-y, 1-z; (ii) 2-x, -y, 2-z.]

Hexakis(μ_2 -4-amino-3,5-dimethyl-4H-1,2,4-triazole)hexaaquaticobalt(II) naphthalene-1,5-disulfonate tetrachloride

Crystal data



$M_r = 1385.82$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 11.1641 (4)$ Å

$b = 12.2744 (5)$ Å

$c = 13.2265 (5)$ Å

$\alpha = 106.924 (2)^\circ$

$\beta = 99.622 (3)^\circ$

$\gamma = 103.452 (2)^\circ$

$V = 1631.79 (11)$ Å³

$Z = 1$

$F(000) = 715$

$D_x = 1.410 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3738 reflections

$\theta = 2.2\text{--}25.7^\circ$

$\mu = 1.05 \text{ mm}^{-1}$

$T = 100$ K

Block, pink

$0.26 \times 0.22 \times 0.20$ mm

Data collection

Bruker SMART APEX CCD
diffractometer

Radiation source: sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2001)

$T_{\min} = 0.763$, $T_{\max} = 0.815$

23481 measured reflections

6410 independent reflections

5057 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.043$

$\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 1.7^\circ$

$h = -13 \rightarrow 13$

$k = -15 \rightarrow 15$

$l = -16 \rightarrow 16$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.048$$

$$wR(F^2) = 0.108$$

$$S = 1.06$$

6410 reflections

386 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0427P)^2 + 2.0543P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.67 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.75 \text{ e \AA}^{-3}$$

Extinction correction: *SHELXTL* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0020 (5)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Co1	0.63106 (4)	0.64428 (4)	0.31065 (4)	0.02470 (13)	
Co2	0.5000	0.5000	0.5000	0.02126 (15)	
O1W	0.5966 (2)	0.56654 (19)	0.14192 (17)	0.0265 (5)	
H1X	0.5055	0.5253	0.1106	0.040*	
H1Y	0.6234	0.6285	0.1105	0.040*	
O2W	0.6118 (2)	0.79286 (19)	0.27619 (18)	0.0261 (5)	
H2X	0.6232	0.8578	0.3444	0.039*	
H2Y	0.5942	0.7753	0.1971	0.039*	
O3W	0.8229 (2)	0.70517 (19)	0.31226 (18)	0.0268 (5)	
H3X	0.8292	0.7098	0.2406	0.040*	
H3Y	0.8764	0.7196	0.3845	0.040*	
N1	0.6573 (2)	0.4793 (2)	0.3235 (2)	0.0223 (6)	
N2	0.6086 (2)	0.4269 (2)	0.3951 (2)	0.0215 (5)	
N3	0.7090 (2)	0.3163 (2)	0.3066 (2)	0.0213 (5)	
N4	0.7638 (3)	0.2234 (2)	0.2677 (2)	0.0229 (6)	
H4A	0.7329	0.1885	0.1940	0.034*	0.667
H4B	0.7436	0.1680	0.3001	0.034*	0.667
H4C	0.8499	0.2545	0.2842	0.034*	0.667
N5	0.4307 (2)	0.5840 (2)	0.2976 (2)	0.0223 (6)	
N6	0.3797 (2)	0.5345 (2)	0.3702 (2)	0.0210 (5)	
N7	0.2246 (2)	0.5399 (2)	0.2502 (2)	0.0214 (5)	
N8	0.0988 (2)	0.5242 (2)	0.1957 (2)	0.0233 (6)	
H8A	0.0997	0.5532	0.1398	0.035*	0.667

H8B	0.0604	0.5640	0.2434	0.035*	0.667
H8C	0.0549	0.4451	0.1687	0.035*	0.667
N9	0.6756 (2)	0.7319 (2)	0.4850 (2)	0.0234 (6)	
N10	0.6331 (2)	0.6759 (2)	0.5555 (2)	0.0214 (5)	
N11	0.7398 (2)	0.8597 (2)	0.6501 (2)	0.0228 (6)	
N12	0.7993 (3)	0.9696 (2)	0.7368 (2)	0.0302 (7)	
H12A	0.8319	1.0268	0.7095	0.045*	0.667
H12B	0.8632	0.9615	0.7839	0.045*	0.667
H12C	0.7411	0.9911	0.7727	0.045*	0.667
C1	0.7852 (3)	0.4237 (3)	0.1853 (3)	0.0292 (7)	
H1D	0.8529	0.5001	0.2143	0.044*	
H1E	0.7246	0.4222	0.1217	0.044*	
H1F	0.8225	0.3584	0.1635	0.044*	
C2	0.7179 (3)	0.4095 (3)	0.2710 (2)	0.0213 (6)	
C3	0.6427 (3)	0.3295 (3)	0.3836 (2)	0.0204 (6)	
C4	0.6189 (3)	0.2409 (3)	0.4412 (2)	0.0206 (6)	
H4D	0.5658	0.2627	0.4911	0.031*	
H4E	0.7003	0.2410	0.4829	0.031*	
H4F	0.5751	0.1611	0.3874	0.031*	
C5	0.3332 (3)	0.6312 (3)	0.1326 (3)	0.0209 (6)	
H5D	0.4186	0.6815	0.1396	0.031*	
H5E	0.2731	0.6780	0.1321	0.031*	
H5F	0.3068	0.5630	0.0642	0.031*	
C6	0.3351 (3)	0.5871 (3)	0.2273 (2)	0.0219 (6)	
C7	0.2558 (3)	0.5085 (3)	0.3398 (2)	0.0219 (6)	
C8	0.1563 (3)	0.4586 (3)	0.3906 (3)	0.0287 (7)	
H8D	0.1968	0.4413	0.4533	0.043*	
H8E	0.0959	0.3850	0.3367	0.043*	
H8F	0.1110	0.5169	0.4150	0.043*	
C9	0.8092 (3)	0.9393 (3)	0.5089 (3)	0.0281 (7)	
H9A	0.7569	0.9922	0.5030	0.042*	
H9B	0.8251	0.9038	0.4377	0.042*	
H9C	0.8905	0.9853	0.5627	0.042*	
C10	0.7414 (3)	0.8433 (3)	0.5442 (2)	0.0211 (6)	
C11	0.6736 (3)	0.7529 (3)	0.6545 (2)	0.0231 (7)	
C12	0.6622 (3)	0.7396 (3)	0.7613 (3)	0.0245 (7)	
H12D	0.6123	0.6577	0.7490	0.037*	
H12E	0.6196	0.7953	0.7974	0.037*	
H12F	0.7473	0.7569	0.8079	0.037*	
S1	1.06020 (7)	0.26397 (7)	0.94747 (6)	0.02274 (18)	
O1	1.1780 (2)	0.32512 (18)	1.03387 (17)	0.0229 (5)	
O2	1.0829 (2)	0.21065 (18)	0.84071 (17)	0.0222 (5)	
O3	0.9768 (2)	0.33792 (18)	0.94574 (17)	0.0238 (5)	
C13	0.7946 (3)	0.0438 (3)	1.0283 (2)	0.0211 (6)	
H13	0.7140	0.0431	1.0427	0.025*	
C14	0.8632 (3)	0.1383 (3)	1.0024 (3)	0.0235 (7)	
H14	0.8283	0.2011	0.9992	0.028*	
C15	0.9796 (3)	0.1417 (3)	0.9818 (3)	0.0229 (7)	

C16	1.0358 (3)	0.0482 (3)	0.9874 (3)	0.0226 (6)	
C17	1.1565 (3)	0.0484 (3)	0.9670 (3)	0.0245 (7)	
H17	1.2041	0.1126	0.9501	0.029*	
Cl1	0.45668 (16)	0.91733 (15)	0.39605 (14)	0.0331 (4)	0.50
Cl2	0.47349 (13)	0.01196 (12)	0.18426 (11)	0.0180 (3)	0.50
Cl3	0.57922 (14)	0.65370 (13)	0.97649 (13)	0.0249 (3)	0.50
Cl4	0.95318 (12)	0.22467 (12)	0.48318 (11)	0.0172 (3)	0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0198 (2)	0.0236 (2)	0.0234 (2)	0.00047 (17)	0.00243 (17)	0.00400 (18)
Co2	0.0170 (3)	0.0209 (3)	0.0225 (3)	0.0040 (2)	0.0033 (2)	0.0048 (2)
O1W	0.0259 (12)	0.0256 (11)	0.0212 (11)	0.0007 (9)	0.0044 (9)	0.0047 (9)
O2W	0.0255 (12)	0.0253 (11)	0.0179 (11)	-0.0031 (9)	0.0005 (9)	0.0047 (9)
O3W	0.0229 (12)	0.0313 (12)	0.0218 (11)	0.0013 (9)	0.0037 (9)	0.0094 (10)
N1	0.0169 (13)	0.0225 (13)	0.0235 (14)	0.0027 (10)	0.0053 (11)	0.0043 (11)
N2	0.0226 (13)	0.0186 (12)	0.0208 (13)	0.0064 (10)	0.0067 (11)	0.0024 (10)
N3	0.0245 (13)	0.0222 (13)	0.0171 (13)	0.0120 (11)	0.0064 (11)	0.0022 (10)
N4	0.0327 (15)	0.0256 (14)	0.0203 (13)	0.0210 (12)	0.0139 (11)	0.0085 (11)
N5	0.0223 (14)	0.0188 (13)	0.0214 (14)	0.0036 (10)	0.0026 (11)	0.0039 (10)
N6	0.0180 (13)	0.0211 (13)	0.0213 (13)	0.0045 (10)	0.0047 (10)	0.0047 (11)
N7	0.0188 (13)	0.0228 (13)	0.0201 (13)	0.0043 (10)	0.0037 (10)	0.0059 (11)
N8	0.0166 (13)	0.0243 (13)	0.0256 (14)	0.0034 (10)	-0.0009 (11)	0.0092 (11)
N9	0.0232 (14)	0.0245 (13)	0.0205 (13)	0.0048 (11)	0.0048 (11)	0.0070 (11)
N10	0.0196 (13)	0.0203 (13)	0.0206 (13)	0.0054 (10)	0.0033 (10)	0.0029 (11)
N11	0.0217 (13)	0.0240 (13)	0.0172 (13)	0.0037 (11)	0.0059 (10)	0.0012 (10)
N12	0.0273 (15)	0.0239 (14)	0.0193 (14)	-0.0075 (11)	0.0020 (11)	-0.0079 (11)
C1	0.0257 (17)	0.0386 (19)	0.0260 (17)	0.0113 (15)	0.0110 (14)	0.0111 (15)
C2	0.0212 (15)	0.0205 (15)	0.0180 (15)	0.0050 (12)	0.0011 (12)	0.0036 (12)
C3	0.0204 (15)	0.0189 (14)	0.0169 (15)	0.0054 (12)	0.0011 (12)	0.0011 (12)
C4	0.0187 (15)	0.0227 (15)	0.0230 (16)	0.0110 (12)	0.0063 (12)	0.0074 (13)
C5	0.0148 (14)	0.0225 (15)	0.0265 (16)	0.0034 (12)	0.0041 (12)	0.0125 (13)
C6	0.0221 (16)	0.0212 (15)	0.0148 (15)	0.0014 (12)	0.0024 (12)	0.0003 (12)
C7	0.0194 (16)	0.0245 (15)	0.0153 (15)	0.0025 (12)	0.0023 (12)	0.0018 (12)
C8	0.0263 (17)	0.0353 (18)	0.0307 (18)	0.0078 (14)	0.0070 (14)	0.0217 (15)
C9	0.0279 (17)	0.0238 (16)	0.0290 (18)	0.0020 (13)	0.0129 (14)	0.0052 (14)
C10	0.0240 (16)	0.0233 (15)	0.0147 (14)	0.0075 (12)	0.0069 (12)	0.0034 (12)
C11	0.0246 (16)	0.0223 (15)	0.0190 (15)	0.0064 (13)	0.0062 (13)	0.0023 (12)
C12	0.0239 (16)	0.0196 (15)	0.0242 (16)	0.0021 (12)	0.0050 (13)	0.0032 (13)
S1	0.0218 (4)	0.0219 (4)	0.0222 (4)	0.0040 (3)	0.0031 (3)	0.0076 (3)
O1	0.0187 (11)	0.0242 (11)	0.0227 (11)	0.0010 (9)	0.0035 (9)	0.0088 (9)
O2	0.0221 (11)	0.0241 (11)	0.0225 (11)	0.0046 (9)	0.0091 (9)	0.0108 (9)
O3	0.0239 (11)	0.0239 (11)	0.0273 (12)	0.0080 (9)	0.0003 (9)	0.0169 (9)
C13	0.0220 (15)	0.0191 (15)	0.0184 (15)	0.0072 (12)	0.0050 (12)	0.0002 (12)
C14	0.0200 (16)	0.0253 (16)	0.0215 (16)	0.0072 (13)	0.0035 (12)	0.0032 (13)
C15	0.0238 (16)	0.0169 (14)	0.0246 (16)	0.0019 (12)	0.0058 (13)	0.0055 (12)
C16	0.0195 (15)	0.0212 (15)	0.0207 (15)	0.0032 (12)	0.0002 (12)	0.0028 (12)

C17	0.0237 (16)	0.0220 (15)	0.0225 (16)	0.0017 (13)	0.0042 (13)	0.0051 (13)
Cl1	0.0314 (9)	0.0308 (8)	0.0284 (9)	-0.0015 (7)	0.0048 (7)	0.0074 (7)
Cl2	0.0182 (7)	0.0195 (7)	0.0128 (6)	0.0016 (5)	0.0015 (5)	0.0050 (5)
Cl3	0.0226 (7)	0.0253 (8)	0.0287 (8)	0.0041 (6)	0.0114 (6)	0.0115 (6)
Cl4	0.0133 (6)	0.0211 (7)	0.0223 (7)	0.0067 (5)	0.0008 (5)	0.0155 (6)

Geometric parameters (Å, °)

Co1—O2W	2.053 (2)	N12—H12A	0.9100
Co1—O1W	2.084 (2)	N12—H12B	0.9100
Co1—O3W	2.092 (2)	N12—H12C	0.9100
Co1—N5	2.146 (3)	C1—C2	1.490 (4)
Co1—N9	2.156 (3)	C1—H1D	0.9800
Co1—N1	2.162 (3)	C1—H1E	0.9800
Co2—N2 ⁱ	2.132 (3)	C1—H1F	0.9800
Co2—N2	2.132 (3)	C3—C4	1.501 (4)
Co2—N10	2.151 (2)	C4—H4D	0.9800
Co2—N10 ⁱ	2.151 (2)	C4—H4E	0.9800
Co2—N6 ⁱ	2.201 (3)	C4—H4F	0.9800
Co2—N6	2.201 (3)	C5—C6	1.502 (4)
O1W—H1X	0.9796	C5—H5D	0.9800
O1W—H1Y	0.9802	C5—H5E	0.9800
O2W—H2X	0.9801	C5—H5F	0.9800
O2W—H2Y	0.9797	C7—C8	1.494 (4)
O3W—H3X	0.9793	C8—H8D	0.9800
O3W—H3Y	0.9807	C8—H8E	0.9800
N1—C2	1.325 (4)	C8—H8F	0.9800
N1—N2	1.404 (4)	C9—C10	1.485 (4)
N2—C3	1.312 (4)	C9—H9A	0.9800
N3—C2	1.346 (4)	C9—H9B	0.9800
N3—C3	1.350 (4)	C9—H9C	0.9800
N3—N4	1.423 (3)	C11—C12	1.492 (4)
N4—H4A	0.9100	C12—H12D	0.9800
N4—H4B	0.9100	C12—H12E	0.9800
N4—H4C	0.9100	C12—H12F	0.9800
N5—C6	1.308 (4)	S1—O3	1.445 (2)
N5—N6	1.407 (3)	S1—O1	1.462 (2)
N6—C7	1.308 (4)	S1—O2	1.464 (2)
N7—C6	1.358 (4)	S1—C15	1.781 (3)
N7—C7	1.366 (4)	C13—C17 ⁱⁱ	1.377 (4)
N7—N8	1.409 (3)	C13—C14	1.398 (4)
N8—H8A	0.9100	C13—H13	0.9500
N8—H8B	0.9100	C14—C15	1.365 (4)
N8—H8C	0.9100	C14—H14	0.9500
N9—C10	1.319 (4)	C15—C16	1.444 (4)
N9—N10	1.391 (3)	C16—C17	1.417 (5)
N10—C11	1.302 (4)	C16—C16 ⁱⁱ	1.425 (6)
N11—C10	1.359 (4)	C17—C13 ⁱⁱ	1.377 (4)

N11—C11	1.369 (4)	C17—H17	0.9500
N11—N12	1.416 (3)		
O2W—Co1—O1W	84.31 (9)	H12A—N12—H12B	109.5
O2W—Co1—O3W	86.35 (9)	N11—N12—H12C	109.5
O1W—Co1—O3W	86.73 (9)	H12A—N12—H12C	109.5
O2W—Co1—N5	90.77 (9)	H12B—N12—H12C	109.5
O1W—Co1—N5	89.79 (9)	C2—C1—H1D	109.5
O3W—Co1—N5	175.68 (9)	C2—C1—H1E	109.5
O2W—Co1—N9	94.08 (9)	H1D—C1—H1E	109.5
O1W—Co1—N9	176.90 (9)	C2—C1—H1F	109.5
O3W—Co1—N9	90.53 (9)	H1D—C1—H1F	109.5
N5—Co1—N9	92.89 (10)	H1E—C1—H1F	109.5
O2W—Co1—N1	171.49 (10)	N1—C2—N3	108.6 (3)
O1W—Co1—N1	87.76 (9)	N1—C2—C1	128.7 (3)
O3W—Co1—N1	90.13 (9)	N3—C2—C1	122.7 (3)
N5—Co1—N1	92.27 (9)	N2—C3—N3	108.8 (3)
N9—Co1—N1	93.70 (10)	N2—C3—C4	130.0 (3)
N2 ⁱ —Co2—N2	180.00 (11)	N3—C3—C4	121.1 (3)
N2 ⁱ —Co2—N10	87.58 (9)	C3—C4—H4D	109.5
N2—Co2—N10	92.42 (10)	C3—C4—H4E	109.5
N2 ⁱ —Co2—N10 ⁱ	92.42 (10)	H4D—C4—H4E	109.5
N2—Co2—N10 ⁱ	87.58 (9)	C3—C4—H4F	109.5
N10—Co2—N10 ⁱ	180.0	H4D—C4—H4F	109.5
N2 ⁱ —Co2—N6 ⁱ	93.28 (9)	H4E—C4—H4F	109.5
N2—Co2—N6 ⁱ	86.72 (9)	C6—C5—H5D	109.5
N10—Co2—N6 ⁱ	87.17 (9)	C6—C5—H5E	109.5
N10 ⁱ —Co2—N6 ⁱ	92.83 (9)	H5D—C5—H5E	109.5
N2 ⁱ —Co2—N6	86.72 (9)	C6—C5—H5F	109.5
N2—Co2—N6	93.28 (9)	H5D—C5—H5F	109.5
N10—Co2—N6	92.83 (9)	H5E—C5—H5F	109.5
N10 ⁱ —Co2—N6	87.17 (9)	N5—C6—N7	109.2 (3)
N6 ⁱ —Co2—N6	180.000 (1)	N5—C6—C5	130.4 (3)
Co1—O1W—H1X	109.4	N7—C6—C5	120.4 (3)
Co1—O1W—H1Y	109.6	N6—C7—N7	108.7 (3)
H1X—O1W—H1Y	109.5	N6—C7—C8	129.9 (3)
Co1—O2W—H2X	109.4	N7—C7—C8	121.4 (3)
Co1—O2W—H2Y	109.5	C7—C8—H8D	109.5
H2X—O2W—H2Y	141.1	C7—C8—H8E	109.5
Co1—O3W—H3X	109.3	H8D—C8—H8E	109.5
Co1—O3W—H3Y	109.6	C7—C8—H8F	109.5
H3X—O3W—H3Y	141.1	H8D—C8—H8F	109.5
C2—N1—N2	106.9 (2)	H8E—C8—H8F	109.5
C2—N1—Co1	130.1 (2)	C10—C9—H9A	109.5
N2—N1—Co1	123.00 (18)	C10—C9—H9B	109.5
C3—N2—N1	107.4 (2)	H9A—C9—H9B	109.5
C3—N2—Co2	129.0 (2)	C10—C9—H9C	109.5
N1—N2—Co2	123.60 (18)	H9A—C9—H9C	109.5

C2—N3—C3	108.1 (2)	H9B—C9—H9C	109.5
C2—N3—N4	123.8 (2)	N9—C10—N11	108.1 (3)
C3—N3—N4	128.1 (2)	N9—C10—C9	129.2 (3)
N3—N4—H4A	109.5	N11—C10—C9	122.8 (3)
N3—N4—H4B	109.5	N10—C11—N11	108.2 (3)
H4A—N4—H4B	109.5	N10—C11—C12	131.1 (3)
N3—N4—H4C	109.5	N11—C11—C12	120.7 (3)
H4A—N4—H4C	109.5	C11—C12—H12D	109.5
H4B—N4—H4C	109.5	C11—C12—H12E	109.5
C6—N5—N6	107.3 (2)	H12D—C12—H12E	109.5
C6—N5—Co1	129.4 (2)	C11—C12—H12F	109.5
N6—N5—Co1	123.37 (18)	H12D—C12—H12F	109.5
C7—N6—N5	107.7 (2)	H12E—C12—H12F	109.5
C7—N6—Co2	130.1 (2)	O3—S1—O1	112.19 (13)
N5—N6—Co2	122.12 (18)	O3—S1—O2	113.46 (13)
C6—N7—C7	107.2 (2)	O1—S1—O2	112.75 (13)
C6—N7—N8	128.9 (3)	O3—S1—C15	106.52 (14)
C7—N7—N8	123.9 (2)	O1—S1—C15	105.66 (13)
N7—N8—H8A	109.5	O2—S1—C15	105.50 (14)
N7—N8—H8B	109.5	C17 ⁱⁱ —C13—C14	120.3 (3)
H8A—N8—H8B	109.5	C17 ⁱⁱ —C13—H13	119.9
N7—N8—H8C	109.5	C14—C13—H13	119.9
H8A—N8—H8C	109.5	C15—C14—C13	121.1 (3)
H8B—N8—H8C	109.5	C15—C14—H14	119.5
C10—N9—N10	107.7 (2)	C13—C14—H14	119.5
C10—N9—Co1	129.0 (2)	C14—C15—C16	120.7 (3)
N10—N9—Co1	123.31 (18)	C14—C15—S1	118.5 (2)
C11—N10—N9	108.3 (2)	C16—C15—S1	120.8 (2)
C11—N10—Co2	127.8 (2)	C17—C16—C16 ⁱⁱ	119.7 (4)
N9—N10—Co2	123.28 (18)	C17—C16—C15	122.7 (3)
C10—N11—C11	107.6 (2)	C16 ⁱⁱ —C16—C15	117.6 (4)
C10—N11—N12	123.8 (3)	C13 ⁱⁱ —C17—C16	120.6 (3)
C11—N11—N12	128.5 (3)	C13 ⁱⁱ —C17—H17	119.7
N11—N12—H12A	109.5	C16—C17—H17	119.7
N11—N12—H12B	109.5		
O1W—Co1—N1—C2	42.1 (3)	Co1—N1—C2—N3	179.84 (19)
O3W—Co1—N1—C2	−44.6 (3)	N2—N1—C2—C1	179.3 (3)
N5—Co1—N1—C2	131.8 (3)	Co1—N1—C2—C1	−1.0 (5)
N9—Co1—N1—C2	−135.2 (3)	C3—N3—C2—N1	−0.7 (3)
O1W—Co1—N1—N2	−138.2 (2)	N4—N3—C2—N1	−179.8 (3)
O3W—Co1—N1—N2	135.1 (2)	C3—N3—C2—C1	−180.0 (3)
N5—Co1—N1—N2	−48.5 (2)	N4—N3—C2—C1	0.9 (4)
N9—Co1—N1—N2	44.5 (2)	N1—N2—C3—N3	−1.0 (3)
C2—N1—N2—C3	0.6 (3)	Co2—N2—C3—N3	178.92 (19)
Co1—N1—N2—C3	−179.2 (2)	N1—N2—C3—C4	178.6 (3)
C2—N1—N2—Co2	−179.37 (19)	Co2—N2—C3—C4	−1.4 (5)
Co1—N1—N2—Co2	0.9 (3)	C2—N3—C3—N2	1.1 (3)

N10—Co2—N2—C3	132.7 (3)	N4—N3—C3—N2	-179.8 (3)
N10 ⁱ —Co2—N2—C3	-47.3 (3)	C2—N3—C3—C4	-178.6 (3)
N6 ⁱ —Co2—N2—C3	45.7 (3)	N4—N3—C3—C4	0.5 (5)
N6—Co2—N2—C3	-134.3 (3)	N6—N5—C6—N7	-1.0 (3)
N10—Co2—N2—N1	-47.3 (2)	Co1—N5—C6—N7	-179.58 (19)
N10 ⁱ —Co2—N2—N1	132.7 (2)	N6—N5—C6—C5	179.7 (3)
N6 ⁱ —Co2—N2—N1	-134.3 (2)	Co1—N5—C6—C5	1.1 (5)
N6—Co2—N2—N1	45.7 (2)	C7—N7—C6—N5	0.9 (3)
O2W—Co1—N5—C6	40.3 (3)	N8—N7—C6—N5	-179.4 (3)
O1W—Co1—N5—C6	-44.0 (3)	C7—N7—C6—C5	-179.7 (3)
N9—Co1—N5—C6	134.5 (3)	N5—N6—C7—N7	-0.2 (3)
N1—Co1—N5—C6	-131.7 (3)	Co2—N6—C7—N7	-175.57 (19)
O2W—Co1—N5—N6	-138.0 (2)	N5—N6—C7—C8	-177.4 (3)
O1W—Co1—N5—N6	137.7 (2)	Co2—N6—C7—C8	7.3 (5)
N9—Co1—N5—N6	-43.9 (2)	C6—N7—C7—N6	-0.4 (3)
N1—Co1—N5—N6	49.9 (2)	N8—N7—C7—N6	179.9 (3)
C6—N5—N6—C7	0.8 (3)	C6—N7—C7—C8	177.1 (3)
Co1—N5—N6—C7	179.4 (2)	N8—N7—C7—C8	-2.7 (5)
C6—N5—N6—Co2	176.57 (19)	N10—N9—C10—N11	-1.1 (3)
Co1—N5—N6—Co2	-4.8 (3)	Co1—N9—C10—N11	177.3 (2)
N2 ⁱ —Co2—N6—C7	-48.8 (3)	N10—N9—C10—C9	178.4 (3)
N2—Co2—N6—C7	131.2 (3)	Co1—N9—C10—C9	-3.2 (5)
N10—Co2—N6—C7	-136.2 (3)	C11—N11—C10—N9	1.9 (4)
N10 ⁱ —Co2—N6—C7	43.8 (3)	N12—N11—C10—N9	-178.9 (3)
N2 ⁱ —Co2—N6—N5	136.4 (2)	C11—N11—C10—C9	-177.7 (3)
N2—Co2—N6—N5	-43.6 (2)	N12—N11—C10—C9	1.5 (5)
N10—Co2—N6—N5	49.0 (2)	N9—N10—C11—N11	1.3 (3)
N10 ⁱ —Co2—N6—N5	-131.0 (2)	Co2—N10—C11—N11	-169.8 (2)
O2W—Co1—N9—C10	-35.7 (3)	N9—N10—C11—C12	-176.3 (3)
O3W—Co1—N9—C10	50.7 (3)	Co2—N10—C11—C12	12.5 (5)
N5—Co1—N9—C10	-126.7 (3)	C10—N11—C11—N10	-2.0 (4)
N1—Co1—N9—C10	140.9 (3)	N12—N11—C11—N10	178.9 (3)
O2W—Co1—N9—N10	142.5 (2)	C10—N11—C11—C12	175.9 (3)
O3W—Co1—N9—N10	-131.1 (2)	N12—N11—C11—C12	-3.2 (5)
N5—Co1—N9—N10	51.5 (2)	C17 ⁱⁱ —C13—C14—C15	-0.3 (5)
N1—Co1—N9—N10	-41.0 (2)	C13—C14—C15—C16	-0.7 (5)
C10—N9—N10—C11	-0.2 (3)	C13—C14—C15—S1	179.1 (2)
Co1—N9—N10—C11	-178.7 (2)	O3—S1—C15—C14	-1.7 (3)
C10—N9—N10—Co2	171.5 (2)	O1—S1—C15—C14	117.8 (3)
Co1—N9—N10—Co2	-7.0 (3)	O2—S1—C15—C14	-122.6 (3)
N2 ⁱ —Co2—N10—C11	41.0 (3)	O3—S1—C15—C16	178.2 (2)
N2—Co2—N10—C11	-139.0 (3)	O1—S1—C15—C16	-62.3 (3)
N6 ⁱ —Co2—N10—C11	-52.4 (3)	O2—S1—C15—C16	57.3 (3)
N6—Co2—N10—C11	127.6 (3)	C14—C15—C16—C17	-179.7 (3)
N2 ⁱ —Co2—N10—N9	-128.9 (2)	S1—C15—C16—C17	0.4 (4)
N2—Co2—N10—N9	51.1 (2)	C14—C15—C16—C16 ⁱⁱ	1.4 (5)
N6 ⁱ —Co2—N10—N9	137.7 (2)	S1—C15—C16—C16 ⁱⁱ	-178.5 (3)

N6—Co2—N10—N9	−42.3 (2)	C16 ⁱⁱ —C16—C17—C13 ⁱⁱ	−0.1 (5)
N2—N1—C2—N3	0.1 (3)	C15—C16—C17—C13 ⁱⁱ	−179.0 (3)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+2, -y, -z+2$.

Hydrogen-bond geometry (\AA , °)

D—H···A	D—H	H···A	D···A	D—H···A
O1W—H1X···Cl3 ⁱ	0.98	2.06	2.765 (3)	127
O1W—H1Y···Cl3 ⁱⁱⁱ	0.98	1.89	2.701 (3)	138
O2W—H2X···Cl1	0.98	2.28	2.929 (3)	122
O2W—H2Y···Cl3 ⁱⁱⁱ	0.98	2.82	3.749 (3)	158
O3W—H3X···O2 ^{iv}	0.98	1.89	2.773 (3)	148
O3W—H3Y···Cl4 ^{iv}	0.98	2.18	3.114 (3)	159
N8—H8A···O3 ⁱ	0.91	2.21	3.009 (3)	146

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (iii) $x, y, z-1$; (iv) $-x+2, -y+1, -z+1$.