

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,9-Dimethyl-1,10-phenanthrolin-1-ium tetrachloridoferrate(III) methanol monosolvate

Ehsan Bahojb Noruzi, Nasser Safari,* Vahid Amani and Behrouz Notash

Department of Chemistry, Shahid Beheshti University, G. C., Evin, Tehran 1983963113, Iran

Correspondence e-mail: n-safari@sbu.ac.ir


Received 29 May 2012; accepted 1 June 2012

 Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.045; wR factor = 0.099; data-to-parameter ratio = 22.0.

In the title compound, $(\text{C}_{14}\text{H}_{13}\text{N}_2)[\text{FeCl}_4]\cdot\text{CH}_3\text{OH}$, the 2,9-dimethyl-1,10-phenanthrolin-1-ium cation, FeCl_4^- anion and methanol solvent molecule lie on a twofold rotation axis. Due to symmetry, the H atom on the N atom of the cation is half-occupied. In the anion, the Fe^{III} atom has a tetrahedral geometry. H atoms of the methanol molecule are disordered over two sets of sites around the twofold axis. In the crystal, $\pi-\pi$ contacts between the pyridine rings and between the pyridine and benzene rings [centroid-centroid distances = 3.6535 (16) and 3.5522 (17) Å] and intermolecular $\text{O}-\text{H}\cdots\text{N}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds stabilize the structure.

Related literature

For related structures, see: Abboud *et al.* (2005); Amani *et al.* (2007, 2009); Khavasi *et al.* (2008); Moreno *et al.* (2006); Morsali (2005); Veidis *et al.* (1981); Yousefi *et al.* (2007); Yu *et al.* (2006).



Experimental

Crystal data

$(\text{C}_{14}\text{H}_{13}\text{N}_2)[\text{FeCl}_4]\cdot\text{CH}_3\text{O}$
 $M_r = 438.96$
 Monoclinic, $C2/c$

$a = 13.067$ (3) Å
 $b = 20.377$ (4) Å
 $c = 7.2810$ (15) Å

$\beta = 109.03$ (3)°
 $V = 1832.7$ (7) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 1.41$ mm⁻¹
 $T = 120$ K
 $0.20 \times 0.15 \times 0.10$ mm

Data collection

Stoe IPDS-2T diffractometer
 Absorption correction: numerical
 (*X-SHAPE* and *X-RED*; Stoe & Cie, 2002)
 $T_{\text{min}} = 0.766$, $T_{\text{max}} = 0.872$

9967 measured reflections
 2464 independent reflections
 2075 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.099$
 $S = 1.10$
 2464 reflections
 112 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.84$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.84$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots N1	0.84	2.33	2.751 (4)	112
N1—H1D \cdots O1	0.65 (12)	2.10 (10)	2.751 (4)	175 (14)

Data collection: *X-Area* (Stoe & Cie, 2002); cell refinement: *X-Area*; data reduction: *X-RED* (Stoe & Cie, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We are grateful to the Shahid Beheshti University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2554).

References

- Abboud, K., Harrowfield, J. M., James, B. D., Skelton, B. W. & White, A. H. (2005). *Inorg. Chim. Acta*, **358**, 1293–1297.
- Amani, V., Safari, N. & Khavasi, H. R. (2007). *Polyhedron*, **26**, 4257–4262.
- Amani, V., Safari, N., Notash, B. & Khavasi, H. R. (2009). *J. Coord. Chem.* **62**, 1939–1950.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Khavasi, H. R., Amani, V. & Safari, N. (2008). *Z. Kristallogr. New Cryst. Struct.* **223**, 41–42.
- Moreno, M. A., Haukka, M., Kallinen, M. & Pakkanen, T. A. (2006). *Appl. Organomet. Chem.* **20**, 51–69.
- Morsali, A. (2005). *Anal. Sci.* **21**, x21–x22.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoe & Cie (2002). *X-Area*, *X-RED* and *X-SHAPE*. Stoe & Cie, Darmstadt, Germany.
- Veidis, M. V., Witten, E. H., Reiff, W. M., Brennan, T. F. & Garafalo, A. R. (1981). *Inorg. Chim. Acta*, **54**, L133–L135.
- Yousefi, M., Ahmadi, R., Amani, V. & Khavasi, H. R. (2007). *Acta Cryst.* **E63**, m3114–m3115.
- Yu, Y.-Q., Ding, C.-F., Zhang, M.-L., Li, X.-M. & Zhang, S.-S. (2006). *Acta Cryst.* **E62**, o2187–o2189.

supporting information

Acta Cryst. (2012). E68, m905 [https://doi.org/10.1107/S1600536812025111]

2,9-Dimethyl-1,10-phenanthroline-1-ium tetrachloridoferrate(III) methanol monosolvate

Ehsan Bahojb Noruzi, Nasser Safari, Vahid Amani and Behrouz Notash

S1. Comment

In the recent years, we reported the synthesis and crystal structures of iron(III) proton transfer complexes, such as [Fe(bipy)Cl₄][bipy.H] (Amani *et al.*, 2007), [Fe(phen)Cl₄][phen.H] (Khavasi *et al.*, 2008), [Fe(4,4'-dmbpy)Cl₄][4,4'-dmbpy.H] (Amani *et al.*, 2009) (bipy = 2,2'-bipyridine, phen = 1,10-phenanthroline, 4,4'-dmbpy = 4,4'-dimethyl-2,2'-bipyridine). Several proton transfer systems using 2,9-dimethyl-1,10-phenanthroline as proton donor molecules, such as [Me₂phen.H](ClO₄) (Morsali, 2005), [Me₂phen.H](NO₃) (Yu *et al.*, 2006), [Me₂phen.H][Ru(CO)₃Cl₃] (Moreno *et al.*, 2006) and [Me₂phen.H]₂[PtCl₆] (Yousefi *et al.*, 2007) (Me₂phen.H = 2,9-dimethyl-1,10-phenanthroline) have been synthesized and characterized by single-crystal X-ray diffraction methods. Also, the structure of [Me₂phen.H][FeCl₄] was reported (Veidis *et al.*, 1981). We report herein the synthesis and crystal structure of the title compound.


The molecular structure of the title compound is shown in Fig. 1. The asymmetric unit of the title compound contains half of a protonated 2,9-dimethyl-1,10-phenanthroline cation, half of a FeCl₄⁻ anion and half of a methanol solvent molecule. In the anion, the Fe^{III} atom has a tetrahedral coordination. The Fe—Cl bond lengths and angles are within normal range (Abboud *et al.*, 2005; Amani *et al.*, 2007). In the crystal, intermolecular O—H⋯N and N—H⋯O hydrogen bonds (Table 1) and π—π interactions between the pyridine and benzene rings, Cg1⋯Cg1ⁱ = 3.6535 (16) and Cg1⋯Cg2ⁱⁱ = 3.5522 (17) Å [Cg1 and Cg2 are the centroids of the N1/C2–C6 ring and the C5, C6, C7, C5ⁱⁱⁱ, C6ⁱⁱⁱ, C7ⁱⁱⁱ ring; symmetry codes: (i) *x*, 1-*y*, -1/2+*z*; (ii) 2-*x*, 1-*y*, -*z*; (iii) 2-*x*, *y*, 1/2-*z*], stabilize the structure (Fig. 2).

S2. Experimental

For the preparation of the title compound, a solution of 2,9-dimethyl-1,10-phenanthroline (0.25 g, 1.20 mmol) in acetonitrile (10 ml) was added to a solution of FeCl₃·6H₂O (0.11 g, 0.40 mmol) in methanol (10 ml) and the resulting yellow solution was stirred at 313 K for 2 h. This solution was left to evaporate slowly at room temperature. After 5 days, yellow needle crystals of the title compound were isolated (yield: 0.13 g, 74.0%; m. p. 426 K).


S3. Refinement

H atoms bonded to N atom was found in difference Fourier map and refined isotropically. H atoms bonded to C and O atoms were positioned geometrically and refined as riding atoms, with O—H = 0.84, C—H = 0.95 (aromatic) and 0.98 (methyl) Å and with $U_{\text{iso}}(\text{H}) = 1.2(1.5 \text{ for methyl and hydroxyl})U_{\text{eq}}(\text{C}, \text{O})$. H atoms of the methanol solvent molecule are disordered over two sets of sites around a twofold axis.

**Figure 1**

The molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level.

[Symmetry code: (i) $2-x, y, 1/2-z$.]

**Figure 2**

Crystal packing diagram for the title compound. Hydrogen bonds are shown as dashed lines.

2,9-Dimethyl-1,10-phenanthroline-1-ium tetrachloridoferrate(III) methanol monosolvate

Crystal data

(C₁₄H₁₃N₂)[FeCl₄]·CH₄O $M_r = 438.96$ Monoclinic, *C2/c*

Hall symbol: -C 2yc

 $a = 13.067$ (3) Å $b = 20.377$ (4) Å $c = 7.2810$ (15) Å $\beta = 109.03$ (3)° $V = 1832.7$ (7) Å³ $Z = 4$ $F(000) = 892$ $D_x = 1.591$ Mg m⁻³Mo *K* α radiation, $\lambda = 0.71073$ Å

Cell parameters from 2464 reflections

 $\theta = 3.1$ – 29.2 ° $\mu = 1.41$ mm⁻¹ $T = 120$ K

Needle, yellow

 $0.20 \times 0.15 \times 0.10$ mm

Data collection

Stoe IPDS-2T

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω scans

Absorption correction: numerical

(X-SHAPE and X-RED; Stoe & Cie, 2002) $T_{\min} = 0.766$, $T_{\max} = 0.872$

9967 measured reflections

2464 independent reflections

2075 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.049$ $\theta_{\text{max}} = 29.2$ °, $\theta_{\text{min}} = 3.1$ ° $h = -17 \rightarrow 17$ $k = -27 \rightarrow 27$ $l = -9 \rightarrow 9$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.045$ $wR(F^2) = 0.099$ $S = 1.10$

2464 reflections

112 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0412P)^2 + 3.5467P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} = 0.001$ $\Delta\rho_{\text{max}} = 0.84$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.84$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Fe1	1.0000	0.15100 (2)	0.2500	0.01928 (13)	
Cl1	0.85420 (5)	0.21251 (3)	0.17847 (11)	0.03440 (18)	
Cl2	0.99637 (5)	0.08774 (3)	0.00604 (10)	0.03183 (17)	

O1	1.0000	0.68189 (19)	0.2500	0.139 (3)	
H1	0.9766	0.6682	0.1350	0.208*	0.50
N1	0.89657 (16)	0.56531 (10)	0.1150 (3)	0.0181 (4)	
C1	0.7488 (2)	0.63346 (14)	-0.0768 (4)	0.0304 (6)	
H1A	0.7225	0.6522	0.0235	0.046*	
H1B	0.6883	0.6291	-0.1985	0.046*	
H1C	0.8039	0.6624	-0.0974	0.046*	
C2	0.79690 (19)	0.56745 (13)	-0.0131 (3)	0.0232 (5)	
C3	0.7407 (2)	0.50876 (14)	-0.0831 (4)	0.0290 (6)	
H3	0.6692	0.5103	-0.1729	0.035*	
C4	0.7888 (2)	0.44989 (14)	-0.0223 (4)	0.0305 (6)	
H4	0.7508	0.4105	-0.0709	0.037*	
C5	0.8949 (2)	0.44681 (12)	0.1129 (4)	0.0242 (5)	
C6	0.94610 (18)	0.50710 (11)	0.1799 (3)	0.0178 (4)	
C7	0.9500 (2)	0.38679 (12)	0.1836 (4)	0.0323 (6)	
H7	0.9157	0.3462	0.1363	0.039*	
C8	1.0000	0.7446 (2)	0.2500	0.073 (2)	
H8A	0.9305	0.7606	0.2564	0.088*	0.50
H8B	1.0107	0.7606	0.1307	0.088*	0.50
H8C	1.0587	0.7606	0.3629	0.088*	0.50
H1D	0.924 (9)	0.592 (5)	0.147 (16)	0.088*	0.50

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.0158 (2)	0.0150 (2)	0.0223 (2)	0.000	-0.00025 (17)	0.000
Cl1	0.0220 (3)	0.0221 (3)	0.0523 (4)	0.0066 (2)	0.0028 (3)	0.0083 (3)
Cl2	0.0328 (3)	0.0324 (3)	0.0289 (3)	-0.0084 (3)	0.0083 (3)	-0.0095 (2)
O1	0.154 (5)	0.0212 (19)	0.141 (5)	0.000	-0.090 (5)	0.000
N1	0.0187 (9)	0.0185 (9)	0.0181 (9)	-0.0025 (7)	0.0074 (7)	-0.0004 (7)
C1	0.0194 (11)	0.0393 (15)	0.0290 (13)	0.0023 (10)	0.0030 (9)	0.0067 (11)
C2	0.0200 (11)	0.0314 (13)	0.0195 (11)	-0.0052 (9)	0.0085 (9)	-0.0008 (9)
C3	0.0216 (11)	0.0415 (15)	0.0254 (12)	-0.0124 (10)	0.0097 (10)	-0.0077 (11)
C4	0.0309 (13)	0.0343 (14)	0.0327 (14)	-0.0189 (11)	0.0192 (11)	-0.0154 (11)
C5	0.0307 (12)	0.0204 (11)	0.0297 (13)	-0.0092 (10)	0.0211 (10)	-0.0062 (9)
C6	0.0192 (10)	0.0181 (10)	0.0193 (10)	-0.0012 (8)	0.0107 (9)	-0.0012 (8)
C7	0.0450 (15)	0.0153 (11)	0.0507 (17)	-0.0049 (10)	0.0349 (14)	-0.0062 (10)
C8	0.099 (5)	0.026 (2)	0.070 (4)	0.000	-0.005 (4)	0.000

Geometric parameters (Å, °)

Fe1—Cl2	2.1826 (8)	C2—C3	1.409 (4)
Fe1—Cl2 ⁱ	2.1826 (8)	C3—C4	1.359 (4)
Fe1—Cl1 ⁱ	2.1966 (8)	C3—H3	0.9500
Fe1—Cl1	2.1966 (8)	C4—C5	1.416 (4)
O1—C8	1.277 (6)	C4—H4	0.9500
O1—H1	0.8400	C5—C6	1.408 (3)
N1—C2	1.333 (3)	C5—C7	1.427 (4)

N1—C6	1.360 (3)	C6—C6 ⁱ	1.445 (4)
N1—H1D	0.64 (11)	C7—C7 ⁱ	1.349 (6)
C1—C2	1.493 (4)	C7—H7	0.9500
C1—H1A	0.9800	C8—H8A	0.9800
C1—H1B	0.9800	C8—H8B	0.9800
C1—H1C	0.9800	C8—H8C	0.9800
Cl2—Fe1—Cl2 ⁱ	107.61 (5)	C4—C3—H3	120.0
Cl2—Fe1—Cl1 ⁱ	108.48 (3)	C2—C3—H3	120.0
Cl2 ⁱ —Fe1—Cl1 ⁱ	110.92 (4)	C3—C4—C5	120.6 (2)
Cl2—Fe1—Cl1	110.92 (4)	C3—C4—H4	119.7
Cl2 ⁱ —Fe1—Cl1	108.48 (3)	C5—C4—H4	119.7
Cl1 ⁱ —Fe1—Cl1	110.41 (4)	C6—C5—C4	116.7 (2)
C8—O1—H1	109.5	C6—C5—C7	119.7 (2)
C2—N1—C6	121.2 (2)	C4—C5—C7	123.6 (2)
C2—N1—H1D	122 (10)	N1—C6—C5	121.5 (2)
C6—N1—H1D	117 (10)	N1—C6—C6 ⁱ	119.29 (12)
C2—C1—H1A	109.5	C5—C6—C6 ⁱ	119.25 (14)
C2—C1—H1B	109.5	C7 ⁱ —C7—C5	121.03 (16)
H1A—C1—H1B	109.5	C7 ⁱ —C7—H7	119.5
C2—C1—H1C	109.5	C5—C7—H7	119.5
H1A—C1—H1C	109.5	O1—C8—H8A	109.5
H1B—C1—H1C	109.5	O1—C8—H8B	109.5
N1—C2—C3	120.0 (2)	H8A—C8—H8B	109.5
N1—C2—C1	117.6 (2)	O1—C8—H8C	109.5
C3—C2—C1	122.3 (2)	H8A—C8—H8C	109.5
C4—C3—C2	120.0 (2)	H8B—C8—H8C	109.5
C6—N1—C2—C3	0.3 (3)	C2—N1—C6—C6 ⁱ	-179.5 (3)
C6—N1—C2—C1	-179.8 (2)	C4—C5—C6—N1	-1.1 (3)
N1—C2—C3—C4	-1.0 (4)	C7—C5—C6—N1	179.5 (2)
C1—C2—C3—C4	179.1 (3)	C4—C5—C6—C6 ⁱ	179.2 (3)
C2—C3—C4—C5	0.6 (4)	C7—C5—C6—C6 ⁱ	-0.2 (4)
C3—C4—C5—C6	0.4 (4)	C6—C5—C7—C7 ⁱ	1.1 (5)
C3—C4—C5—C7	179.7 (3)	C4—C5—C7—C7 ⁱ	-178.2 (3)
C2—N1—C6—C5	0.8 (3)		

Symmetry code: (i) $-x+2, y, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 \cdots N1	0.84	2.33	2.751 (4)	112
N1—H1D \cdots O1	0.65 (12)	2.10 (10)	2.751 (4)	175 (14)