

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-(4-Bromophenylsulfonyl)-5-chloro-2-methyl-1-benzofuran

 Hong Dae Choi,^a Pil Ja Seo^a and Uk Lee^{b*}

^aDepartment of Chemistry, Donggeui University, San 24 Kaya-dong, Busanjin-gu, Busan 614-714, Republic of Korea, and ^bDepartment of Chemistry, Pukyong National University, 599-1 Daeyeon 3-dong, Nam-gu, Busan 608-737, Republic of Korea

Correspondence e-mail: uklee@pknu.ac.kr


Received 24 May 2012; accepted 30 May 2012

 Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.036; wR factor = 0.097; data-to-parameter ratio = 18.7.

In the title compound, $\text{C}_{15}\text{H}_{10}\text{BrClO}_3\text{S}$, the 4-bromo-substituted benzene ring forms a dihedral angle of $72.55(6)^\circ$ with the mean plane [mean deviation = $0.008(2)$ Å] of the benzofuran ring system. In the crystal, molecules are linked by weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds into chains along [001]. There are also $\pi-\pi$ interactions between the furan and benzene rings of symmetry-related benzofuran systems [centroid-centroid distances = $3.549(3)$ and $3.632(3)$ Å].

Related literature

For background information and the crystal structures of related compounds, see: Choi *et al.* (2008, 2010).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{10}\text{BrClO}_3\text{S}$
 $M_r = 385.65$
 Triclinic, $P\bar{1}$
 $a = 7.1089(2)$ Å
 $b = 10.4169(3)$ Å
 $c = 11.0217(3)$ Å
 $\alpha = 69.659(2)^\circ$
 $\beta = 89.526(2)^\circ$
 $\gamma = 71.632(1)^\circ$
 $V = 721.58(4)$ Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 3.18$ mm⁻¹
 $T = 173$ K
 $0.38 \times 0.31 \times 0.27$ mm

Data collection

Bruker SMART APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.448$, $T_{\max} = 0.746$
 13182 measured reflections
 3570 independent reflections
 3007 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.097$
 $S = 1.08$
 3570 reflections
 191 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.45$ e Å⁻³
 $\Delta\rho_{\min} = -0.43$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C11}-\text{H11}\cdots\text{O2}^i$	0.95	2.53	3.203 (3)	128
$\text{C15}-\text{H15}\cdots\text{O3}^{ii}$	0.95	2.57	3.195 (3)	124

 Symmetry codes: (i) $-x + 1, -y, -z + 2$; (ii) $-x + 1, -y, -z + 1$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and DIAMOND (Brandenburg, 1998); software used to prepare material for publication: SHELXL97.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5482).

References

- Brandenburg, K. (1998). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Bruker (2009). *APEX2*, *SADABS* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2008). *Acta Cryst.* **E64**, o1190.
 Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2010). *Acta Cryst.* **E66**, o2350.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2012). E68, o2027 [https://doi.org/10.1107/S1600536812024592]

3-(4-Bromophenylsulfonyl)-5-chloro-2-methyl-1-benzofuran**Hong Dae Choi, Pil Ja Seo and Uk Lee****S1. Comment**

As a part of our ongoing study of 5-chloro-2-methyl-1-benzofuran derivatives containing 3-phenylsulfonyl (Choi *et al.*, 2008) and 3-(4-fluorophenylsulfonyl) (Choi *et al.*, 2010) substituents, we report herein the crystal structure of the title compound.

In the title molecule (Fig. 1), the benzofuran unit is essentially planar, with a mean deviation of 0.008 (2) Å from the least-squares plane defined by the nine constituent atoms. The dihedral angle between the 4-bromo substituted benzene ring and the mean plane of the benzofuran fragment is 72.55 (6)°. In the crystal structure (Fig. 2), molecules are connected by weak C—H···O hydrogen bonds (Table 1). The crystal packing (Fig. 3) also exhibits offset π – π interactions between the furan and benzene rings of neighbouring molecules, with Cg1···Cg2ⁱⁱⁱ & Cg1···Cg2^{iv} distances of 3.549 (3) Å & 3.632 (3) Å [Symmetry codes: (iii) - x + 1, - y + 1, - z + 1; (iv) - x, - y + 1, - z + 1] and interplanar distances of 3.510 (3) Å & 3.343 (3) Å resulting in slippages of 0.511 (3) Å & 1.420 (3) Å (Cg1 and Cg2 are the centroids of the C1/C2/C7/O1/C8 furan ring and the C2–C7 benzene ring, respectively).

S2. Experimental

3-Chloroperoxybenzoic acid (77%, 381 mg, 1.7 mmol) was added in small portions to a stirred solution of 3-(4-bromophenylsulfonyl)-5-chloro-2-methyl-1-benzofuran (318 mg, 0.8 mmol) in dichloromethane (40 mL) at 273 K. After being stirred at room temperature for 8h, the mixture was washed with saturated sodium bicarbonate solution and the organic layer was separated, dried over magnesium sulfate, filtered and concentrated at reduced pressure. The residue was purified by column chromatography (benzene) to afford the title compound as a colorless solid [yield 76%, m.p. 454–456 K; R_f = 0.71 (benzene)]. Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of the title compound in ethyl acetate at room temperature.

S3. Refinement

All H atoms were positioned geometrically and refined using a riding model, with C—H = 0.95 Å for aryl and 0.98 Å for methyl H atoms. $U_{iso}(H) = 1.2U_{eq}(C)$ for aryl and $1.5U_{eq}(C)$ for methyl H atoms. The positions of methyl hydrogens were optimized rotationally.




Figure 1

The molecular structure of the title compound with displacement ellipsoids drawn at the 50% probability level. H atoms are presented as small spheres of arbitrary radius.




Figure 2

A view of the C—H···O interactions (dotted lines) in the packing of the title compound. H atoms non-participating in hydrogen-bonding are omitted for clarity. [Symmetry codes: (i) $-x + 1, -y, -z + 2$; (ii) $-x + 1, -y, -z + 1$].




Figure 3

A view of the π - π interactions (dotted lines) in the crystal packing of the title compound. All H atoms are omitted for clarity. [Symmetry codes: (iii) $-x + 1, -y + 1, -z + 1$; (iv) $-x, -y + 1, -z + 1$].

3-(4-Bromophenylsulfonyl)-5-chloro-2-methyl-1-benzofuran

Crystal data

$C_{15}H_{10}BrClO_3S$
 $M_r = 385.65$
 Triclinic, $P\bar{1}$
 Hall symbol: -P 1
 $a = 7.1089$ (2) Å
 $b = 10.4169$ (3) Å
 $c = 11.0217$ (3) Å
 $\alpha = 69.659$ (2)°
 $\beta = 89.526$ (2)°
 $\gamma = 71.632$ (1)°
 $V = 721.58$ (4) Å³

$Z = 2$
 $F(000) = 384$
 $D_x = 1.775$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 7922 reflections
 $\theta = 2.5$ – 28.3 °
 $\mu = 3.18$ mm⁻¹
 $T = 173$ K
 Block, colourless
 $0.38 \times 0.31 \times 0.27$ mm

Data collection

Bruker SMART APEXII CCD
 diffractometer
 Radiation source: rotating anode
 Graphite multilayer monochromator
 Detector resolution: 10.0 pixels mm⁻¹
 φ and ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2009)
 $T_{\min} = 0.448$, $T_{\max} = 0.746$

13182 measured reflections
 3570 independent reflections
 3007 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$
 $\theta_{\max} = 28.3$ °, $\theta_{\min} = 2.0$ °
 $h = -9 \rightarrow 8$
 $k = -13 \rightarrow 13$
 $l = -14 \rightarrow 14$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.097$
 $S = 1.08$
 3570 reflections
 191 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: difference Fourier map
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0512P)^2 + 0.0238P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.45$ e Å⁻³
 $\Delta\rho_{\min} = -0.43$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	-0.05797 (4)	-0.23812 (2)	0.82146 (3)	0.03829 (11)
Cl1	0.31879 (13)	0.47313 (9)	0.12675 (7)	0.0514 (2)
S1	0.53111 (8)	0.13394 (6)	0.70691 (5)	0.02350 (14)

O1	0.1919 (2)	0.53895 (17)	0.62583 (18)	0.0320 (4)
O2	0.6309 (3)	0.10707 (19)	0.83068 (17)	0.0320 (4)
O3	0.6452 (2)	0.10389 (17)	0.60557 (16)	0.0282 (4)
C1	0.3820 (3)	0.3152 (2)	0.6417 (2)	0.0237 (4)
C2	0.3255 (3)	0.4018 (2)	0.5047 (2)	0.0242 (5)
C3	0.3615 (4)	0.3785 (3)	0.3887 (2)	0.0288 (5)
H3	0.4391	0.2863	0.3874	0.035*
C4	0.2782 (4)	0.4967 (3)	0.2754 (3)	0.0345 (6)
C5	0.1640 (4)	0.6333 (3)	0.2737 (3)	0.0406 (7)
H5	0.1098	0.7108	0.1929	0.049*
C6	0.1294 (4)	0.6564 (3)	0.3885 (3)	0.0378 (7)
H6	0.0539	0.7491	0.3898	0.045*
C7	0.2097 (4)	0.5387 (3)	0.5014 (3)	0.0293 (5)
C8	0.2974 (4)	0.4014 (3)	0.7101 (3)	0.0297 (5)
C9	0.2938 (4)	0.3784 (3)	0.8495 (3)	0.0400 (6)
H9A	0.1613	0.3780	0.8744	0.060*
H9B	0.3238	0.4568	0.8661	0.060*
H9C	0.3940	0.2850	0.9008	0.060*
C10	0.3644 (3)	0.0345 (2)	0.7367 (2)	0.0246 (5)
C11	0.2993 (4)	-0.0064 (3)	0.8589 (2)	0.0333 (6)
H11	0.3425	0.0202	0.9252	0.040*
C12	0.1709 (4)	-0.0863 (3)	0.8835 (3)	0.0364 (6)
H12	0.1249	-0.1146	0.9667	0.044*
C13	0.1106 (4)	-0.1243 (2)	0.7867 (3)	0.0291 (5)
C14	0.1743 (4)	-0.0834 (3)	0.6636 (2)	0.0305 (5)
H14	0.1313	-0.1105	0.5976	0.037*
C15	0.3008 (4)	-0.0028 (3)	0.6388 (2)	0.0286 (5)
H15	0.3444	0.0272	0.5550	0.034*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.03630 (17)	0.02806 (14)	0.0493 (2)	-0.01477 (11)	0.00645 (13)	-0.00910 (12)
Cl1	0.0655 (5)	0.0720 (5)	0.0242 (3)	-0.0428 (4)	0.0035 (3)	-0.0084 (3)
S1	0.0238 (3)	0.0256 (3)	0.0201 (3)	-0.0069 (2)	0.0006 (2)	-0.0082 (2)
O1	0.0265 (9)	0.0271 (8)	0.0463 (11)	-0.0081 (7)	0.0063 (8)	-0.0187 (8)
O2	0.0309 (9)	0.0384 (9)	0.0253 (9)	-0.0116 (7)	-0.0032 (7)	-0.0097 (7)
O3	0.0263 (9)	0.0290 (8)	0.0267 (9)	-0.0060 (7)	0.0046 (7)	-0.0101 (7)
C1	0.0205 (11)	0.0259 (10)	0.0255 (12)	-0.0089 (9)	0.0020 (9)	-0.0092 (9)
C2	0.0192 (11)	0.0253 (10)	0.0294 (12)	-0.0104 (9)	0.0007 (9)	-0.0088 (9)
C3	0.0286 (12)	0.0300 (11)	0.0274 (13)	-0.0130 (10)	0.0027 (10)	-0.0071 (10)
C4	0.0331 (14)	0.0455 (14)	0.0267 (13)	-0.0246 (11)	0.0010 (11)	-0.0051 (11)
C5	0.0333 (14)	0.0357 (13)	0.0408 (17)	-0.0189 (11)	-0.0064 (12)	0.0071 (12)
C6	0.0244 (13)	0.0228 (11)	0.0604 (19)	-0.0103 (9)	-0.0014 (12)	-0.0059 (12)
C7	0.0219 (11)	0.0290 (11)	0.0397 (14)	-0.0129 (9)	0.0030 (10)	-0.0116 (10)
C8	0.0235 (12)	0.0351 (12)	0.0354 (14)	-0.0120 (10)	0.0042 (10)	-0.0165 (11)
C9	0.0392 (15)	0.0550 (16)	0.0365 (15)	-0.0181 (13)	0.0100 (12)	-0.0274 (13)
C10	0.0260 (12)	0.0228 (10)	0.0225 (11)	-0.0069 (9)	0.0015 (9)	-0.0061 (9)

C11	0.0420 (15)	0.0394 (13)	0.0227 (12)	-0.0190 (11)	0.0048 (11)	-0.0114 (11)
C12	0.0431 (15)	0.0384 (13)	0.0251 (13)	-0.0182 (12)	0.0050 (11)	-0.0042 (11)
C13	0.0275 (12)	0.0215 (10)	0.0334 (13)	-0.0065 (9)	0.0051 (10)	-0.0059 (9)
C14	0.0316 (13)	0.0337 (12)	0.0297 (13)	-0.0111 (10)	0.0020 (11)	-0.0153 (10)
C15	0.0312 (13)	0.0307 (11)	0.0242 (12)	-0.0097 (10)	0.0057 (10)	-0.0108 (10)

Geometric parameters (Å, °)

Br1—C13	1.887 (2)	C6—C7	1.374 (4)
Cl1—C4	1.748 (3)	C6—H6	0.9500
S1—O2	1.4380 (18)	C8—C9	1.472 (4)
S1—O3	1.4392 (17)	C9—H9A	0.9800
S1—C1	1.735 (2)	C9—H9B	0.9800
S1—C10	1.765 (2)	C9—H9C	0.9800
O1—C8	1.374 (3)	C10—C11	1.387 (3)
O1—C7	1.378 (3)	C10—C15	1.389 (3)
C1—C8	1.364 (3)	C11—C12	1.385 (4)
C1—C2	1.446 (3)	C11—H11	0.9500
C2—C3	1.389 (3)	C12—C13	1.373 (4)
C2—C7	1.392 (3)	C12—H12	0.9500
C3—C4	1.383 (4)	C13—C14	1.393 (4)
C3—H3	0.9500	C14—C15	1.378 (4)
C4—C5	1.392 (4)	C14—H14	0.9500
C5—C6	1.375 (4)	C15—H15	0.9500
C5—H5	0.9500		
O2—S1—O3	120.14 (11)	C1—C8—O1	110.0 (2)
O2—S1—C1	108.61 (11)	C1—C8—C9	134.5 (2)
O3—S1—C1	106.53 (10)	O1—C8—C9	115.6 (2)
O2—S1—C10	107.53 (11)	C8—C9—H9A	109.5
O3—S1—C10	107.70 (10)	C8—C9—H9B	109.5
C1—S1—C10	105.43 (11)	H9A—C9—H9B	109.5
C8—O1—C7	107.14 (17)	C8—C9—H9C	109.5
C8—C1—C2	107.8 (2)	H9A—C9—H9C	109.5
C8—C1—S1	126.30 (19)	H9B—C9—H9C	109.5
C2—C1—S1	125.94 (17)	C11—C10—C15	120.7 (2)
C3—C2—C7	119.6 (2)	C11—C10—S1	119.19 (19)
C3—C2—C1	135.8 (2)	C15—C10—S1	120.08 (18)
C7—C2—C1	104.6 (2)	C12—C11—C10	119.5 (2)
C4—C3—C2	116.4 (2)	C12—C11—H11	120.3
C4—C3—H3	121.8	C10—C11—H11	120.3
C2—C3—H3	121.8	C13—C12—C11	119.5 (2)
C3—C4—C5	123.4 (3)	C13—C12—H12	120.2
C3—C4—Cl1	118.2 (2)	C11—C12—H12	120.2
C5—C4—Cl1	118.4 (2)	C12—C13—C14	121.5 (2)
C6—C5—C4	120.2 (2)	C12—C13—Br1	119.62 (19)
C6—C5—H5	119.9	C14—C13—Br1	118.8 (2)
C4—C5—H5	119.9	C15—C14—C13	118.9 (2)

C7—C6—C5	116.6 (2)	C15—C14—H14	120.5
C7—C6—H6	121.7	C13—C14—H14	120.5
C5—C6—H6	121.7	C14—C15—C10	119.8 (2)
C6—C7—O1	125.6 (2)	C14—C15—H15	120.1
C6—C7—C2	123.9 (2)	C10—C15—H15	120.1
O1—C7—C2	110.5 (2)		
O2—S1—C1—C8	-27.8 (2)	C1—C2—C7—O1	-0.6 (2)
O3—S1—C1—C8	-158.5 (2)	C2—C1—C8—O1	-0.8 (3)
C10—S1—C1—C8	87.2 (2)	S1—C1—C8—O1	179.12 (16)
O2—S1—C1—C2	152.06 (19)	C2—C1—C8—C9	178.4 (3)
O3—S1—C1—C2	21.3 (2)	S1—C1—C8—C9	-1.7 (4)
C10—S1—C1—C2	-92.9 (2)	C7—O1—C8—C1	0.4 (2)
C8—C1—C2—C3	-179.8 (3)	C7—O1—C8—C9	-179.0 (2)
S1—C1—C2—C3	0.3 (4)	O2—S1—C10—C11	22.3 (2)
C8—C1—C2—C7	0.8 (2)	O3—S1—C10—C11	153.12 (19)
S1—C1—C2—C7	-179.05 (17)	C1—S1—C10—C11	-93.4 (2)
C7—C2—C3—C4	0.4 (3)	O2—S1—C10—C15	-157.24 (19)
C1—C2—C3—C4	-178.8 (2)	O3—S1—C10—C15	-26.4 (2)
C2—C3—C4—C5	0.1 (4)	C1—S1—C10—C15	87.0 (2)
C2—C3—C4—C11	-179.96 (17)	C15—C10—C11—C12	0.5 (4)
C3—C4—C5—C6	0.3 (4)	S1—C10—C11—C12	-179.0 (2)
C11—C4—C5—C6	-179.7 (2)	C10—C11—C12—C13	0.3 (4)
C4—C5—C6—C7	-1.1 (4)	C11—C12—C13—C14	-0.5 (4)
C5—C6—C7—O1	-179.8 (2)	C11—C12—C13—Br1	178.11 (19)
C5—C6—C7—C2	1.7 (4)	C12—C13—C14—C15	0.0 (4)
C8—O1—C7—C6	-178.5 (2)	Br1—C13—C14—C15	-178.66 (18)
C8—O1—C7—C2	0.2 (2)	C13—C14—C15—C10	0.8 (4)
C3—C2—C7—C6	-1.4 (4)	C11—C10—C15—C14	-1.1 (4)
C1—C2—C7—C6	178.1 (2)	S1—C10—C15—C14	178.47 (18)
C3—C2—C7—O1	179.9 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C11—H11 \cdots O2 ⁱ	0.95	2.53	3.203 (3)	128
C15—H15 \cdots O3 ⁱⁱ	0.95	2.57	3.195 (3)	124

Symmetry codes: (i) $-x+1, -y, -z+2$; (ii) $-x+1, -y, -z+1$.