

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 1-(2-Chloroethyl)-1*H*-pyrazolo[3,4-*d*]-pyrimidin-4(5*H*)-one

 Mohammed Iqbal A. Khazi,<sup>a</sup> Nikhath Fathima,<sup>b</sup>  
 Ningaraddi S. Belavagi,<sup>a</sup> Noor Shahina Begum<sup>b</sup> and  
 I. M. Khazi<sup>a\*</sup>
<sup>a</sup>Department of Chemistry, Karnatak University, Dharwad 580 003, India, and  
<sup>b</sup>Department of Studies in Chemistry, Bangalore University, Bangalore 560 001,  
 Karnataka, India  
 Correspondence e-mail: dr\_imk@yahoo.com


Received 24 May 2012; accepted 2 June 2012

 Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  
 $R$  factor = 0.035;  $wR$  factor = 0.100; data-to-parameter ratio = 13.1.

In the title compound,  $\text{C}_7\text{H}_7\text{ClN}_4\text{O}$ , the pyrazolopyrimidine ring is essentially planar, the r.m.s. deviation of the fitted atoms being 0.0071 Å. The crystal structure features strong  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds and further consolidated by weak  $\text{C}-\text{H}\cdots\text{O}$ ,  $\text{C}-\text{H}\cdots\text{N}$  and  $\text{C}-\text{H}\cdots\text{Cl}$  interactions.

## Related literature

For the biological activity of pyrazolopyrimidines, see: Carraro *et al.* (2006). For a related structure, see: Dolzhenko *et al.* (2009). For the graph-set analysis of hydrogen bonding, see: Bernstein *et al.* (1995).



## Experimental

### Crystal data

 $\text{C}_7\text{H}_7\text{ClN}_4\text{O}$   
 $M_r = 198.61$   
 Monoclinic,  $P2_1/n$   
 $a = 4.6448$  (1) Å

 $b = 8.0792$  (1) Å  
 $c = 22.7335$  (4) Å  
 $\beta = 93.554$  (1)°  
 $V = 851.46$  (3) Å<sup>3</sup>
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.41$  mm<sup>-1</sup>
 $T = 296$  K  
 $0.18 \times 0.16 \times 0.16$  mm

### Data collection

 Bruker SMART APEX CCD  
 detector diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 1998)  
 $T_{\min} = 0.930$ ,  $T_{\max} = 0.937$ 

 7660 measured reflections  
 1548 independent reflections  
 1353 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.023$ 

### Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.035$   
 $wR(F^2) = 0.100$   
 $S = 0.87$   
 1548 reflections

 118 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.19$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.37$  e Å<sup>-3</sup>

**Table 1**  
 Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}^{\text{i}}$	0.86	1.96	2.810 (2)	170
$\text{C5}-\text{H5A}\cdots\text{N4}^{\text{ii}}$	0.93	2.79	3.676 (2)	160
$\text{C2}-\text{H2A}\cdots\text{Cl1}^{\text{iii}}$	0.97	2.84	3.779 (2)	164
$\text{C2}-\text{H2B}\cdots\text{N2}^{\text{iv}}$	0.97	2.59	3.463 (2)	150
$\text{C3}-\text{H3}\cdots\text{O1}^{\text{v}}$	0.93	2.35	3.254 (2)	163

Symmetry codes: (i)  $-x + 1, -y + 2, -z + 2$ ; (ii)  $x, y + 1, z$ ; (iii)  $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$ ;  
 (iv)  $-x + \frac{5}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (v)  $-x + 1, -y + 1, -z + 2$ .

Data collection: SMART (Bruker, 1998); cell refinement: SAINT-Plus (Bruker, 1998); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and CAMERON (Watkin *et al.*, 1996); software used to prepare material for publication: WinGX (Farrugia, 1999).

IMK is thankful to the University Grants Commission (UGC), India, for financial assistance.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2552).

## References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N. L. (1995). *Angew. Chem.* **34**, 1555–1573.
- Bruker. (1998). SMART, SAINT-Plus and SADABS. Bruker Axs Inc., Madison, Wisconsin, USA.
- Carraro, F., Naldini, A., Pucci, A., Locatelli, G. A., Maga, G., Schenone, S., Bruno, O., Ranise, A., Bondavalli, F., Brullo, C., Fossa, P., Menozzi, G., Mosti, L., Modugno, M., Tintori, C., Manetti, F. & Botta, M. (2006). *J. Med. Chem.* **49**, 1549–1561.
- Dolzhenko, A. V., Pastorin, G., Dolzhenko, A. V., Tan, G. K. & Koh, L. L. (2009). *Acta Cryst.* **E65**, o1720–o1721.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Watkin, D. J., Prout, C. K. & Pearce, L. J. (1996). CAMERON. Chemical Crystallography Laboratory, University of Oxford, England.

## supporting information

*Acta Cryst.* (2012). E68, o2083 [https://doi.org/10.1107/S1600536812025184]

**1-(2-Chloroethyl)-1*H*-pyrazolo[3,4-*d*]pyrimidin-4(5*H*)-one**

**Mohammed Iqbal A. Khazi, Nikhath Fathima, Ningaraddi S. Belavagi, Noor Shahina Begum and I. M. Khazi**

**S1. Comment**

Pyrazolo[3,4-*d*]pyrimidines are purine analogues which exhibit a number of pharmacological properties such as antitryproliferative (Carraro *et al.*, 2006).

In the title compound (Fig. 1), the fused pyrazolopyrimidine ring is substituted with 2-chloro-ethyl group on one side and the oxo group on the other side. The pyrazolopyrimidine ring is planar with the maximum deviation from the mean statistical plane being 0.0115 (14) Å for C3. The *cis* orientation of 2-chloro-ethyl group with respect to the C2—N2 bond is described by the torsion angle N2—C2—N3—C3, -2.204 (4)°.


The crystal structure is stabilized by some interesting features that comprise of intermolecular N—H···O, C—H···O, C—H···N and C—H···Cl interactions (Fig. 2 and Tab. 1). The C—H···O and the N—H···O interactions result in centrosymmetric head-to-head dimers corresponding to the graph set  $R^2_2(10)$  and  $R^2_2(8)$  motif (Bernstein *et al.*, 1995). There are two types of C—H···N interactions, one of which forms a helix, the other forms sheets along the crystallographic *b*-axis. The C—H···Cl intermolecular interaction result in one dimensional molecular chain along *b*-axis. The bond lengths and bond angles in the title molecule agree very well with the corresponding bond distances and bond angles reported in a closely related compound (Dolzhenko *et al.*, 2009).

**S2. Experimental**

A mixture of 5-amino-1-(2-chloro-ethyl)-1*H*-pyrazole-4-carbonitrile (1 g, 5.8 mmol) and formic acid (15 ml) was heated under reflux for 10 h. The excess of formic acid was removed under reduced pressure and the solid separated was washed with water and recrystallized from ethanol. (Yield = 0.86 g, 75% and m.p. = 470–472 K).

**S3. Refinement**

The H atoms were placed at calculated positions in the riding model approximation with N—H = 0.86 Å and C—H = 0.93, and 0.97 Å for aryl and methylene H-atoms respectively, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N/C})$ .



**Figure 1**

The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as small spheres of arbitrary radius.




Figure 2

A view of the intermolecular hydrogen bonds (dotted lines) in the crystal structure of the title compound. H atoms non participating in H-bonding were omitted for clarity.

### 1-(2-Chloroethyl)-1*H*-pyrazolo[3,4-*d*]pyrimidin-4(5*H*)-one

#### Crystal data

$C_7H_7ClN_4O$

$M_r = 198.61$

Monoclinic,  $P2_1/n$

Hall symbol:  $-P\ 2_1n$

$a = 4.6448$  (1) Å

$b = 8.0792$  (1) Å

$c = 22.7335$  (4) Å

$\beta = 93.554$  (1)°

$V = 851.46$  (3) Å<sup>3</sup>

$Z = 4$

$F(000) = 416$

$D_x = 1.565$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1548 reflections

$\theta = 1.8$ – $25.2$ °

$\mu = 0.41$  mm<sup>-1</sup>

$T = 296$  K

Block, yellow

$0.18 \times 0.16 \times 0.16$  mm

#### Data collection

Bruker SMART APEX CCD detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 1998)

$T_{\min} = 0.930$ ,  $T_{\max} = 0.937$

7660 measured reflections

1548 independent reflections

1353 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.023$

$\theta_{\max} = 25.2$ °,  $\theta_{\min} = 1.8$ °

$h = -5 \rightarrow 5$

$k = -9 \rightarrow 9$

$l = -27 \rightarrow 25$

Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.035$	H-atom parameters constrained
$wR(F^2) = 0.100$	$w = 1/[\sigma^2(F_o^2) + (0.0642P)^2 + 0.5291P]$
$S = 0.87$	where $P = (F_o^2 + 2F_c^2)/3$
1548 reflections	$(\Delta/\sigma)_{\max} = 0.001$
118 parameters	$\Delta\rho_{\max} = 0.19 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.37 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.89765 (12)	0.64259 (8)	0.72864 (2)	0.0600 (2)
O1	0.5103 (3)	0.78930 (17)	0.99906 (6)	0.0516 (4)
N1	0.7733 (3)	0.95763 (18)	0.94256 (6)	0.0381 (4)
H1	0.7021	1.0429	0.9590	0.046*
N2	1.0887 (3)	0.86963 (19)	0.87124 (7)	0.0396 (4)
N3	1.0967 (3)	0.57267 (18)	0.86397 (6)	0.0376 (4)
N4	0.9763 (3)	0.43809 (19)	0.88974 (7)	0.0434 (4)
C5	0.9640 (4)	0.9842 (2)	0.90068 (8)	0.0401 (4)
H5A	1.0092	1.0935	0.8924	0.048*
C6	1.0057 (4)	0.7151 (2)	0.88711 (7)	0.0331 (4)
C1	1.2904 (4)	0.5504 (3)	0.81692 (8)	0.0418 (4)
H1A	1.4385	0.4711	0.8295	0.050*
H1B	1.3848	0.6548	0.8096	0.050*
C2	1.1398 (4)	0.4907 (2)	0.76054 (8)	0.0442 (5)
H2A	1.0330	0.3907	0.7683	0.053*
H2B	1.2826	0.4635	0.7327	0.053*
C4	0.6848 (4)	0.8026 (2)	0.96076 (7)	0.0371 (4)
C7	0.8174 (4)	0.6729 (2)	0.92974 (7)	0.0349 (4)
C3	0.8092 (4)	0.4993 (2)	0.92932 (8)	0.0424 (4)
H3	0.7002	0.4356	0.9538	0.051*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0545 (3)	0.0640 (4)	0.0607 (4)	0.0019 (2)	-0.0027 (3)	0.0011 (3)

O1	0.0692 (9)	0.0403 (8)	0.0488 (8)	-0.0011 (6)	0.0335 (7)	-0.0006 (6)
N1	0.0486 (8)	0.0314 (8)	0.0355 (8)	0.0000 (6)	0.0122 (6)	-0.0036 (6)
N2	0.0475 (9)	0.0357 (8)	0.0370 (8)	-0.0044 (7)	0.0137 (7)	-0.0006 (6)
N3	0.0437 (8)	0.0351 (8)	0.0354 (8)	0.0002 (6)	0.0121 (6)	-0.0014 (6)
N4	0.0550 (9)	0.0326 (8)	0.0436 (9)	0.0000 (7)	0.0117 (7)	0.0026 (7)
C5	0.0489 (10)	0.0352 (10)	0.0372 (9)	-0.0061 (8)	0.0098 (8)	0.0021 (8)
C6	0.0360 (9)	0.0343 (9)	0.0295 (8)	-0.0008 (7)	0.0060 (7)	-0.0005 (7)
C1	0.0397 (9)	0.0451 (11)	0.0418 (10)	0.0050 (8)	0.0132 (8)	-0.0037 (8)
C2	0.0505 (11)	0.0395 (11)	0.0442 (10)	0.0004 (8)	0.0159 (8)	-0.0060 (8)
C4	0.0452 (10)	0.0365 (10)	0.0304 (9)	-0.0030 (8)	0.0088 (7)	-0.0002 (7)
C7	0.0413 (9)	0.0347 (9)	0.0293 (8)	-0.0019 (7)	0.0076 (7)	0.0006 (7)
C3	0.0528 (11)	0.0364 (10)	0.0393 (10)	-0.0025 (8)	0.0141 (8)	0.0033 (8)

*Geometric parameters (Å, °)*

C11—C2	1.788 (2)	C5—H5A	0.9300
O1—C4	1.231 (2)	C6—C7	1.388 (2)
N1—C5	1.357 (2)	C1—C2	1.501 (3)
N1—C4	1.389 (2)	C1—H1A	0.9700
N1—H1	0.8600	C1—H1B	0.9700
N2—C5	1.299 (2)	C2—H2A	0.9700
N2—C6	1.362 (2)	C2—H2B	0.9700
N3—C6	1.344 (2)	C4—C7	1.424 (2)
N3—N4	1.371 (2)	C7—C3	1.403 (3)
N3—C1	1.451 (2)	C3—H3	0.9300
N4—C3	1.320 (2)		
C5—N1—C4	124.71 (15)	C2—C1—H1B	109.0
C5—N1—H1	117.6	H1A—C1—H1B	107.8
C4—N1—H1	117.6	C1—C2—C11	112.00 (14)
C5—N2—C6	111.97 (15)	C1—C2—H2A	109.2
C6—N3—N4	111.42 (13)	C11—C2—H2A	109.2
C6—N3—C1	128.26 (15)	C1—C2—H2B	109.2
N4—N3—C1	120.30 (15)	C11—C2—H2B	109.2
C3—N4—N3	105.48 (15)	H2A—C2—H2B	107.9
N2—C5—N1	125.45 (17)	O1—C4—N1	120.63 (16)
N2—C5—H5A	117.3	O1—C4—C7	127.58 (17)
N1—C5—H5A	117.3	N1—C4—C7	111.79 (14)
N3—C6—N2	125.45 (15)	C6—C7—C3	105.00 (16)
N3—C6—C7	106.86 (15)	C6—C7—C4	118.38 (16)
N2—C6—C7	127.68 (16)	C3—C7—C4	136.61 (17)
N3—C1—C2	113.08 (15)	N4—C3—C7	111.23 (16)
N3—C1—H1A	109.0	N4—C3—H3	124.4
C2—C1—H1A	109.0	C7—C3—H3	124.4
N3—C1—H1B	109.0		
C6—N3—N4—C3	-0.1 (2)	C5—N1—C4—O1	-179.89 (18)
C1—N3—N4—C3	-178.72 (16)	C5—N1—C4—C7	-0.4 (2)

C6—N2—C5—N1	0.1 (3)	N3—C6—C7—C3	0.2 (2)
C4—N1—C5—N2	0.7 (3)	N2—C6—C7—C3	-179.13 (18)
N4—N3—C6—N2	179.28 (17)	N3—C6—C7—C4	-179.02 (15)
C1—N3—C6—N2	-2.2 (3)	N2—C6—C7—C4	1.6 (3)
N4—N3—C6—C7	-0.1 (2)	O1—C4—C7—C6	178.81 (19)
C1—N3—C6—C7	178.40 (17)	N1—C4—C7—C6	-0.6 (2)
C5—N2—C6—N3	179.47 (17)	O1—C4—C7—C3	-0.1 (4)
C5—N2—C6—C7	-1.3 (3)	N1—C4—C7—C3	-179.6 (2)
C6—N3—C1—C2	-108.1 (2)	N3—N4—C3—C7	0.2 (2)
N4—N3—C1—C2	70.3 (2)	C6—C7—C3—N4	-0.3 (2)
N3—C1—C2—C11	66.78 (19)	C4—C7—C3—N4	178.7 (2)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O1 <sup>i</sup>	0.86	1.96	2.810 (2)	170
C5—H5A...N4 <sup>ii</sup>	0.93	2.79	3.676 (2)	160
C2—H2A...C11 <sup>iii</sup>	0.97	2.84	3.779 (2)	164
C2—H2B...N2 <sup>iv</sup>	0.97	2.59	3.463 (2)	150
C3—H3...O1 <sup>v</sup>	0.93	2.35	3.254 (2)	163

Symmetry codes: (i)  $-x+1, -y+2, -z+2$ ; (ii)  $x, y+1, z$ ; (iii)  $-x+3/2, y-1/2, -z+3/2$ ; (iv)  $-x+5/2, y-1/2, -z+3/2$ ; (v)  $-x+1, -y+1, -z+2$ .