

Di- μ -adipato- κ^4 O¹:O⁶-bis{aqua[5,6-di-phenyl-3-(pyridin-2-yl)-1,2,4-triazine- κ^2 N²,N³]copper(II)}

Wei Xu* and Jin-Li Qi

Center of Applied Solid State Chemistry Research, Ningbo University, Ningbo 315211, People's Republic of China
 Correspondence e-mail: xuwei@nbu.edu.cn


Received 16 May 2012; accepted 5 June 2012

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å;
 R factor = 0.057; wR factor = 0.135; data-to-parameter ratio = 15.9.

In the centrosymmetric binuclear title complex, $[\text{Cu}_2(\text{C}_6\text{H}_8\text{O}_4)_2(\text{C}_{20}\text{H}_{14}\text{N}_4)_2(\text{H}_2\text{O})_2]$ or $[\text{Cu}_2(\text{PDPT})_2(\text{C}_6\text{H}_8\text{O}_4)_2(\text{H}_2\text{O})_2]$ ($\text{PDPT} = 3$ -(2-pyridyl)-5,6-diphenyl-1,2,4-triazine), the Cu atom displays a distorted square-pyramidal coordination environment with the basal plane occupied by two PDPT N atoms and two O atoms from different adipate dianions while a water molecule is located at the apical position. Of the two water H atoms, one participates in an intramolecular hydrogen bond whereas the second participates in an intermolecular hydrogen bond, which leads to the formation of a chain along [010].

Related literature

For the biological activity and applications of triazines, see: Garcia *et al.* (1995); Mashaly *et al.* (1999); Croot & Hunter (2000); Soudi *et al.* (2005); Kawamichi *et al.* (2009).



Experimental

Crystal data

$[\text{Cu}_2(\text{C}_6\text{H}_8\text{O}_4)_2(\text{C}_{20}\text{H}_{14}\text{N}_4)_2(\text{H}_2\text{O})_2]$ $M_r = 1072.08$

Triclinic, $P\bar{1}$	$V = 1186.4$ (5) Å ³
$a = 9.4825$ (19) Å	$Z = 1$
$b = 10.616$ (2) Å	Mo $K\alpha$ radiation
$c = 13.080$ (3) Å	$\mu = 0.97$ mm ⁻¹
$\alpha = 78.96$ (3)°	$T = 295$ K
$\beta = 68.76$ (3)°	$0.30 \times 0.19 \times 0.11$ mm
$\gamma = 76.85$ (3)°	

Data collection

Rigaku R-AXIS RAPID CCD diffractometer	11646 measured reflections
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	5295 independent reflections
$T_{\min} = 0.756$, $T_{\max} = 0.863$	3207 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.061$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.135$	$\Delta\rho_{\text{max}} = 0.61$ e Å ⁻³
$S = 1.02$	$\Delta\rho_{\text{min}} = -0.51$ e Å ⁻³
5295 reflections	
333 parameters	
3 restraints	

Table 1
 Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
O5—H5A···O3 ⁱ	0.83	1.92	2.721 (3)	161
O5—H5B···O1 ⁱⁱ	0.83	2.06	2.878 (4)	167

Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $-x, -y + 1, -z + 1$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSC, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

This project was sponsored by the K. C. Wong Magna Fund in Ningbo University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2363).

References

- Croot, P. L. & Hunter, K. A. (2000). *Anal. Chim. Acta*, **406**, 289–302.
- Garcia, G., Solano, I., Sanchez, G. & Lopez, G. (1995). *Polyhedron*, **14**, 1855–1863.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Kawamichi, T., Haneda, T., Kawano, M. & Fujita, M. (2009). *Nature (London)*, **461**, 633–635.
- Mashaly, M., Bayoumi, H. A. & Taha, A. (1999). *J. Serb. Chem. Soc.* **64**, 621–635.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2004). *CrystalStructure*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Soudi, A. A., Marandi, F., Morsali, A., Kempe, R. & Hertle, I. (2005). *J. Coord. Chem.* **58**, 1631–1637.

supporting information

Acta Cryst. (2012). E68, m906 [https://doi.org/10.1107/S1600536812025676]

Di- μ -adipato- $\kappa^4O^1:O^6$ -bis{aqua[5,6-diphenyl-3-(pyridin-2-yl)-1,2,4-triazine- κ^2N^2,N^3]copper(II)}

Wei Xu and Jin-Li Qi

S1. Comment

Research on coordination chemistry of triazine-derived ligands has progressed very rapidly during the past two decades (Kawamichi *et al.*, 2009). The 1,2,4-triazine compounds are well-known in natural materials and show interesting biological, pharmacological and medicinal properties (Garcia *et al.*, 1995). The 3-(2-pyridyl)-5,6-diphenyl-1,2,4-triazine (*PDPT*) represents a principal class of *N*-donor heterocyclic ligands that exhibit interesting pharmacological properties such as blood platelet aggregation inhibition, significant activity towards leukemia and ovarian cancer, and anti-HIV activity (Mashaly *et al.*, 1999; Soudi *et al.*, 2005). It also has been widely used as a sensitive reagent for the determination of Fe(II) by spectrophotometric methods, in natural and waste water (Croot & Hunter, 2000). The title complex, was recently prepared and its crystal structure is reported here.

The title compound crystal structure is composed of centrosymmetric binuclear $[\text{Cu}_2(\text{H}_2\text{O})_2(\text{PDPT})_2(\text{C}_6\text{H}_8\text{O}_4)_2]$ complex molecule (Fig. 1). The dinuclear complex molecules are centered at the crystallographic $2e$ positions. Each Cu atom is coordinated by two N atoms of the chelating *PDPT* ligand and three O atoms of one H_2O molecule and two bis-monodentate adipato ligands to form a slightly distorted square-pyramidal coordination with H_2O molecule located at the apical position ($d(\text{Cu}-\text{N}) = 2.028$ (3) \AA , 2.029 (3) \AA , the basal $d(\text{Cu}-\text{O}) = 1.921$ (3) \AA , 1.961 (3) \AA , the axial $d(\text{Cu}-\text{O}) = 2.375$ (3) \AA). Through the adipato ligands, the square-pyramidal coordinated Cu atoms are linked to form centrosymmetric dinuclear. As expected, the Cu atom is slightly shifted toward the apical water O atom by 0.026 (2) \AA from the least-squares plane defined by the four equatorial coordinating atoms. The triazine ring adopts a slight twist conformation. The dihedral angle between the two phenyl rings is 61.9 (2) $^\circ$.


As shown in the Fig. 2 and Table 1, within the crystal structure, the water molecule O5 forms a strong intramolecular hydrogen bond to the uncoordinated carboxyl O3ⁱ with $O5 \cdots O3^i = 2.721$ (3) \AA and angle O5—H5A…O3ⁱ = 161° . Moreover, it forms an intermolecular hydrogen bond to the coordinated carboxyl O1ⁱⁱ atoms with $O5 \cdots O1^{ii} = 2.878$ (4) \AA and angle O5—H5B…O1ⁱⁱ = 167° to connect the dinuclear complexes along the [0 1 0] direction. Symmetry codes: (i) - x , - y , - $z+1$; (ii) - x , - $y+1$, - $z+1$.

S2. Experimental


Addition of 2.0 mL (1.0 *M*) NaOH to a stirred aqueous of 0.172 g (1.0 mmol) $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ in 5.0 mL H_2O yield a blue precipitate, which was then separated by centrifugation, followed by washing with double-distilled water until no detectable Cl^- anions in supernatant. The precipitate was added to a stirred ethanolic aqueous solution of 0.146 g (1.0 mmol) adipic acid in 20 mL $\text{EtOH}/\text{H}_2\text{O}$ (v:v = 1: 1). To the resulting suspension was added 0.310 g (1.0 mmol) 3-(2-pyridyl)-5,6-diphenyl-1,2,4-triazine (*PDPT*). The mixture was further stirred for approximately 15 min and the insoluble solid was filtered off. The filtrate ($\text{pH} = 6.5$) was allowed to stand at room temperature. Slow evaporation for two weeks afforded a small amount of brown crystals (yield 58% based on the initial $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ input).

S3. Refinement

All H atoms bound to C were position geometrically and refined as riding, with C—H = 0.93 Å and 0.97 Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The H atoms attached to O were located in difference Fourier maps and refined freely with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.


**Figure 1**

The binuclear structure of title compound with the atom numbering scheme. Displacement ellipsoids are drawn at 40% probability level. H atoms are presented as a small spheres of arbitrary radius. Symmetry code: (i) -x, -y, -z+1. Di- μ -adipato- $\kappa^4O^1:O^5$ -bis{aqua[5,6-diphenyl-3-(pyridin-2-yl)-1,2,4-triazine- κ^2N^2,N^3]copper(II)}

**Figure 2**

One dimensional chain through hydrogen bond along [0 1 0]. The C—H bonds omitted for clarity.

Di- μ -adipato- $\kappa^4O^1:O^6$ -bis{aqua[5,6-diphenyl-3-(pyridin-2-yl)-1,2,4-triazine- κ^2N^2,N^3]copper(II)}

Crystal data

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.4825 (19)$ Å

$b = 10.616 (2)$ Å

$c = 13.080 (3)$ Å

$$\alpha = 78.96 (3)^\circ$$

$$\beta = 68.76 (3)^\circ$$

$$\gamma = 76.85 (3)^\circ$$

$$V = 1186.4 (5) \text{ Å}^3$$

$$Z = 1$$

$$F(000) = 554$$

$$D_x = 1.501 \text{ Mg m}^{-3}$$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 8025 reflections
 $\theta = 3.3\text{--}27.5^\circ$
 $\mu = 0.97 \text{ mm}^{-1}$

$T = 295 \text{ K}$
 Plate, brown
 $0.30 \times 0.19 \times 0.11 \text{ mm}$

Data collection

Rigaku R-AXIS RAPID CCD
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω -scans
 Absorption correction: multi-scan
 (ABSCOR; Higashi, 1995)
 $T_{\min} = 0.756$, $T_{\max} = 0.863$

11646 measured reflections
 5295 independent reflections
 3207 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.061$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.3^\circ$
 $h = -12 \rightarrow 11$
 $k = -13 \rightarrow 13$
 $l = -16 \rightarrow 16$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.135$
 $S = 1.02$
 5295 reflections
 333 parameters
 3 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0516P)^2 + 0.8435P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.61 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.51 \text{ e \AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.24237 (6)	0.28612 (5)	0.46540 (4)	0.03879 (17)
N1	0.4452 (4)	0.2512 (3)	0.4938 (3)	0.0374 (8)
N2	0.3453 (3)	0.4301 (3)	0.3587 (3)	0.0333 (7)
N3	0.5595 (3)	0.5336 (3)	0.3071 (3)	0.0343 (7)
N4	0.3041 (3)	0.4987 (3)	0.2743 (3)	0.0355 (7)
O1	0.0833 (3)	0.3122 (3)	0.3970 (2)	0.0390 (6)
O2	0.2814 (3)	0.1907 (3)	0.2883 (3)	0.0488 (7)
O3	0.0418 (3)	-0.2164 (3)	0.3245 (3)	0.0570 (8)
O4	-0.1845 (3)	-0.1321 (3)	0.4379 (3)	0.0488 (8)
O5	0.0969 (3)	0.4299 (3)	0.5999 (3)	0.0495 (8)
H5A	0.042 (4)	0.375 (3)	0.634 (4)	0.074*
H5B	0.040 (4)	0.501 (2)	0.593 (4)	0.074*

C1	0.4946 (5)	0.1514 (4)	0.5589 (3)	0.0436 (10)
H1A	0.4325	0.0892	0.5955	0.052*
C2	0.6347 (5)	0.1373 (4)	0.5739 (3)	0.0460 (11)
H2A	0.6684	0.0647	0.6170	0.055*
C3	0.7228 (5)	0.2320 (4)	0.5244 (4)	0.0471 (11)
H3A	0.8152	0.2262	0.5362	0.057*
C4	0.6740 (4)	0.3366 (4)	0.4566 (3)	0.0404 (9)
H4A	0.7322	0.4020	0.4221	0.048*
C5	0.5364 (4)	0.3401 (4)	0.4422 (3)	0.0333 (8)
C6	0.7323 (4)	0.7259 (4)	0.1838 (4)	0.0414 (10)
H6A	0.7876	0.6483	0.2068	0.050*
C7	0.7968 (5)	0.8365 (4)	0.1501 (4)	0.0495 (11)
H7A	0.8964	0.8330	0.1486	0.059*
C8	0.7152 (5)	0.9530 (4)	0.1183 (4)	0.0520 (12)
H8A	0.7591	1.0279	0.0959	0.062*
C9	0.5676 (5)	0.9572 (4)	0.1200 (4)	0.0497 (11)
H9A	0.5117	1.0358	0.0996	0.060*
C10	0.5033 (5)	0.8481 (4)	0.1509 (4)	0.0419 (10)
H10A	0.4046	0.8525	0.1502	0.050*
C11	0.5839 (4)	0.7290 (3)	0.1840 (3)	0.0331 (8)
C12	0.2046 (4)	0.6241 (4)	0.1017 (4)	0.0423 (10)
H12A	0.1287	0.6023	0.1676	0.051*
C13	0.1724 (5)	0.6510 (4)	0.0040 (4)	0.0492 (11)
H13A	0.0761	0.6445	0.0043	0.059*
C14	0.2816 (5)	0.6872 (4)	-0.0934 (4)	0.0510 (11)
H14A	0.2587	0.7071	-0.1586	0.061*
C15	0.4258 (5)	0.6942 (4)	-0.0942 (3)	0.0444 (10)
H15A	0.4999	0.7191	-0.1601	0.053*
C16	0.4600 (5)	0.6642 (4)	0.0026 (3)	0.0379 (9)
H16A	0.5579	0.6675	0.0011	0.045*
C17	0.3498 (4)	0.6293 (3)	0.1022 (3)	0.0323 (8)
C18	0.4794 (4)	0.4412 (3)	0.3645 (3)	0.0308 (8)
C19	0.5105 (4)	0.6140 (3)	0.2307 (3)	0.0320 (8)
C20	0.3880 (4)	0.5838 (3)	0.2046 (3)	0.0322 (8)
C21	0.1506 (4)	0.2554 (4)	0.3093 (3)	0.0363 (9)
C22	0.0629 (5)	0.2733 (4)	0.2294 (4)	0.0424 (10)
H22A	-0.0456	0.2783	0.2717	0.051*
H22B	0.0762	0.3559	0.1834	0.051*
C23	0.1097 (5)	0.1676 (4)	0.1553 (3)	0.0441 (10)
H23A	0.2148	0.1694	0.1071	0.053*
H23B	0.0456	0.1878	0.1090	0.053*
C24	0.0991 (5)	0.0304 (4)	0.2136 (4)	0.0525 (11)
H24A	0.1160	-0.0266	0.1590	0.063*
H24B	0.1818	0.0018	0.2443	0.063*
C25	-0.0463 (5)	0.0133 (4)	0.3027 (4)	0.0595 (13)
H25A	-0.1292	0.0445	0.2724	0.071*
H25B	-0.0616	0.0685	0.3583	0.071*
C26	-0.0602 (5)	-0.1238 (4)	0.3594 (3)	0.0393 (9)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0376 (3)	0.0390 (3)	0.0392 (3)	-0.0141 (2)	-0.0134 (2)	0.0069 (2)
N1	0.0387 (18)	0.0370 (18)	0.034 (2)	-0.0073 (14)	-0.0125 (15)	0.0035 (15)
N2	0.0339 (16)	0.0338 (17)	0.0325 (19)	-0.0103 (13)	-0.0124 (14)	0.0033 (14)
N3	0.0330 (17)	0.0318 (17)	0.036 (2)	-0.0066 (13)	-0.0103 (14)	-0.0015 (14)
N4	0.0364 (17)	0.0380 (18)	0.0308 (19)	-0.0067 (14)	-0.0130 (14)	0.0026 (14)
O1	0.0372 (14)	0.0379 (15)	0.0414 (18)	-0.0135 (12)	-0.0097 (13)	-0.0023 (13)
O2	0.0369 (16)	0.0528 (18)	0.058 (2)	-0.0044 (14)	-0.0162 (14)	-0.0125 (15)
O3	0.0568 (19)	0.0408 (17)	0.061 (2)	-0.0117 (15)	-0.0073 (16)	0.0011 (16)
O4	0.0459 (17)	0.0403 (16)	0.050 (2)	-0.0119 (13)	-0.0076 (15)	0.0081 (14)
O5	0.0481 (18)	0.0387 (16)	0.056 (2)	-0.0060 (13)	-0.0129 (16)	-0.0035 (15)
C1	0.045 (2)	0.041 (2)	0.037 (3)	-0.0053 (18)	-0.0106 (19)	0.0029 (19)
C2	0.045 (2)	0.050 (3)	0.035 (3)	0.007 (2)	-0.016 (2)	-0.001 (2)
C3	0.034 (2)	0.062 (3)	0.044 (3)	0.000 (2)	-0.015 (2)	-0.010 (2)
C4	0.038 (2)	0.041 (2)	0.043 (3)	-0.0060 (18)	-0.0153 (19)	-0.0050 (19)
C5	0.035 (2)	0.034 (2)	0.029 (2)	-0.0043 (16)	-0.0095 (17)	-0.0043 (16)
C6	0.037 (2)	0.036 (2)	0.048 (3)	-0.0079 (17)	-0.0110 (19)	-0.0018 (19)
C7	0.045 (2)	0.046 (2)	0.060 (3)	-0.020 (2)	-0.012 (2)	-0.007 (2)
C8	0.069 (3)	0.037 (2)	0.055 (3)	-0.027 (2)	-0.018 (2)	0.001 (2)
C9	0.068 (3)	0.030 (2)	0.054 (3)	-0.009 (2)	-0.026 (2)	0.001 (2)
C10	0.046 (2)	0.033 (2)	0.050 (3)	-0.0080 (17)	-0.019 (2)	-0.0043 (19)
C11	0.037 (2)	0.0273 (18)	0.035 (2)	-0.0111 (15)	-0.0100 (17)	-0.0014 (16)
C12	0.035 (2)	0.050 (2)	0.037 (2)	-0.0076 (18)	-0.0116 (18)	0.0057 (19)
C13	0.042 (2)	0.058 (3)	0.047 (3)	-0.004 (2)	-0.021 (2)	0.001 (2)
C14	0.064 (3)	0.046 (3)	0.039 (3)	0.002 (2)	-0.023 (2)	0.000 (2)
C15	0.057 (3)	0.036 (2)	0.029 (2)	-0.0072 (19)	-0.002 (2)	-0.0014 (18)
C16	0.040 (2)	0.033 (2)	0.037 (2)	-0.0084 (16)	-0.0085 (18)	-0.0025 (17)
C17	0.041 (2)	0.0256 (18)	0.031 (2)	-0.0087 (15)	-0.0126 (17)	-0.0012 (16)
C18	0.0319 (19)	0.0302 (19)	0.029 (2)	-0.0081 (15)	-0.0077 (16)	-0.0034 (16)
C19	0.0287 (18)	0.0308 (19)	0.032 (2)	-0.0051 (15)	-0.0064 (16)	-0.0005 (16)
C20	0.0288 (18)	0.0264 (18)	0.037 (2)	-0.0041 (15)	-0.0071 (16)	-0.0027 (16)
C21	0.036 (2)	0.031 (2)	0.042 (3)	-0.0163 (17)	-0.0109 (18)	0.0036 (18)
C22	0.046 (2)	0.038 (2)	0.043 (3)	-0.0154 (18)	-0.017 (2)	0.0094 (19)
C23	0.049 (2)	0.052 (3)	0.034 (2)	-0.021 (2)	-0.016 (2)	0.008 (2)
C24	0.064 (3)	0.048 (3)	0.041 (3)	-0.020 (2)	-0.006 (2)	-0.007 (2)
C25	0.059 (3)	0.046 (3)	0.062 (4)	-0.013 (2)	-0.007 (3)	-0.002 (2)
C26	0.046 (2)	0.034 (2)	0.038 (3)	-0.0121 (18)	-0.014 (2)	-0.0003 (18)

Geometric parameters (\AA , $^\circ$)

Cu1—O1	1.961 (3)	C8—C9	1.382 (6)
Cu1—O4 ⁱ	1.921 (3)	C8—H8A	0.9300
Cu1—O5	2.375 (3)	C9—C10	1.356 (6)
Cu1—N1	2.029 (3)	C9—H9A	0.9300
Cu1—N2	2.028 (3)	C10—C11	1.401 (5)
N1—C1	1.336 (5)	C10—H10A	0.9300

N1—C5	1.341 (5)	C11—C19	1.463 (5)
N2—C18	1.333 (4)	C12—C13	1.382 (6)
N2—N4	1.334 (4)	C12—C17	1.394 (5)
N3—C18	1.321 (5)	C12—H12A	0.9300
N3—C19	1.332 (4)	C13—C14	1.371 (6)
N4—C20	1.329 (5)	C13—H13A	0.9300
O1—C21	1.283 (5)	C14—C15	1.382 (6)
O2—C21	1.235 (4)	C14—H14A	0.9300
O3—C26	1.236 (5)	C15—C16	1.380 (6)
O4—C26	1.260 (5)	C15—H15A	0.9300
O4—Cu1 ⁱ	1.921 (3)	C16—C17	1.390 (5)
O5—H5A	0.829 (18)	C16—H16A	0.9300
O5—H5B	0.830 (18)	C17—C20	1.477 (5)
C1—C2	1.383 (5)	C19—C20	1.438 (5)
C1—H1A	0.9300	C21—C22	1.516 (5)
C2—C3	1.367 (6)	C22—C23	1.508 (6)
C2—H2A	0.9300	C22—H22A	0.9700
C3—C4	1.385 (6)	C22—H22B	0.9700
C3—H3A	0.9300	C23—C24	1.516 (6)
C4—C5	1.376 (5)	C23—H23A	0.9700
C4—H4A	0.9300	C23—H23B	0.9700
C5—C18	1.481 (5)	C24—C25	1.470 (6)
C6—C7	1.372 (6)	C24—H24A	0.9700
C6—C11	1.399 (5)	C24—H24B	0.9700
C6—H6A	0.9300	C25—C26	1.513 (6)
C7—C8	1.378 (6)	C25—H25A	0.9700
C7—H7A	0.9300	C25—H25B	0.9700
O1—Cu1—O5	94.47 (11)	C13—C12—H12A	119.7
O1—Cu1—N1	164.40 (13)	C17—C12—H12A	119.7
O1—Cu1—N2	92.22 (11)	C14—C13—C12	120.4 (4)
O4 ⁱ —Cu1—O1	96.02 (12)	C14—C13—H13A	119.8
O4 ⁱ —Cu1—O5	94.45 (12)	C12—C13—H13A	119.8
O4 ⁱ —Cu1—N1	90.71 (12)	C13—C14—C15	119.8 (4)
O4 ⁱ —Cu1—N2	168.98 (12)	C13—C14—H14A	120.1
N1—Cu1—O5	99.04 (12)	C15—C14—H14A	120.1
N2—Cu1—O5	92.19 (12)	C16—C15—C14	120.1 (4)
N2—Cu1—N1	79.55 (12)	C16—C15—H15A	120.0
C1—N1—C5	117.8 (3)	C14—C15—H15A	120.0
C1—N1—Cu1	126.9 (3)	C15—C16—C17	120.8 (4)
C5—N1—Cu1	115.3 (2)	C15—C16—H16A	119.6
C18—N2—N4	117.7 (3)	C17—C16—H16A	119.6
C18—N2—Cu1	115.4 (2)	C16—C17—C12	118.3 (3)
N4—N2—Cu1	126.0 (2)	C16—C17—C20	121.7 (3)
C18—N3—C19	117.8 (3)	C12—C17—C20	119.6 (3)
C20—N4—N2	120.5 (3)	N3—C18—N2	124.6 (3)
C21—O1—Cu1	104.0 (2)	N3—C18—C5	120.5 (3)
C26—O4—Cu1 ⁱ	126.9 (3)	N2—C18—C5	114.9 (3)

Cu1—O5—H5A	90 (4)	N3—C19—C20	118.0 (3)
Cu1—O5—H5B	129 (4)	N3—C19—C11	115.9 (3)
H5A—O5—H5B	108 (3)	C20—C19—C11	126.0 (3)
N1—C1—C2	122.4 (4)	N4—C20—C19	118.7 (3)
N1—C1—H1A	118.8	N4—C20—C17	114.1 (3)
C2—C1—H1A	118.8	C19—C20—C17	127.0 (3)
C3—C2—C1	118.8 (4)	O2—C21—O1	122.7 (4)
C3—C2—H2A	120.6	O2—C21—C22	120.5 (4)
C1—C2—H2A	120.6	O1—C21—C22	116.8 (3)
C2—C3—C4	119.8 (4)	C23—C22—C21	115.3 (3)
C2—C3—H3A	120.1	C23—C22—H22A	108.5
C4—C3—H3A	120.1	C21—C22—H22A	108.5
C5—C4—C3	117.6 (4)	C23—C22—H22B	108.5
C5—C4—H4A	121.2	C21—C22—H22B	108.5
C3—C4—H4A	121.2	H22A—C22—H22B	107.5
N1—C5—C4	123.4 (3)	C22—C23—C24	115.8 (4)
N1—C5—C18	114.3 (3)	C22—C23—H23A	108.3
C4—C5—C18	122.2 (4)	C24—C23—H23A	108.3
C7—C6—C11	120.4 (4)	C22—C23—H23B	108.3
C7—C6—H6A	119.8	C24—C23—H23B	108.3
C11—C6—H6A	119.8	H23A—C23—H23B	107.4
C6—C7—C8	120.6 (4)	C25—C24—C23	116.0 (4)
C6—C7—H7A	119.7	C25—C24—H24A	108.3
C8—C7—H7A	119.7	C23—C24—H24A	108.3
C7—C8—C9	119.3 (4)	C25—C24—H24B	108.3
C7—C8—H8A	120.3	C23—C24—H24B	108.3
C9—C8—H8A	120.3	H24A—C24—H24B	107.4
C10—C9—C8	120.9 (4)	C24—C25—C26	116.6 (4)
C10—C9—H9A	119.6	C24—C25—H25A	108.2
C8—C9—H9A	119.6	C26—C25—H25A	108.2
C9—C10—C11	120.7 (4)	C24—C25—H25B	108.2
C9—C10—H10A	119.6	C26—C25—H25B	108.2
C11—C10—H10A	119.6	H25A—C25—H25B	107.3
C6—C11—C10	118.1 (3)	O3—C26—O4	125.7 (4)
C6—C11—C19	119.8 (3)	O3—C26—C25	120.3 (4)
C10—C11—C19	121.6 (3)	O4—C26—C25	113.9 (4)
C13—C12—C17	120.5 (4)		

Symmetry code: (i) $-x, -y, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O5—H5A \cdots O3 ⁱ	0.83	1.92	2.721 (3)	161
O5—H5B \cdots O1 ⁱⁱ	0.83	2.06	2.878 (4)	167

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $-x, -y+1, -z+1$.