

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

8-Methyl-2-oxo-4-(thiophen-2-yl)-1,2,5,6,7,8-hexahydroquinoline-3-carbonitrile

Abdullah M. Asiri,^{a,b,‡} Hassan M. Faidallah,^a
Alaa Anwar Ahmad Saqer,^{a,b} Seik Weng Ng^c and
Edward R. T. Tiekink^{c,*}

^aChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203, Jeddah, Saudi Arabia, ^bCenter of Excellence for Advanced Materials Research (CEAMR), King Abdulaziz University, PO Box 80203, Jeddah 21589, Saudi Arabia, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: edward.tiekink@gmail.com


Received 20 June 2012; accepted 22 June 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; disorder in main residue; R factor = 0.088; wR factor = 0.230; data-to-parameter ratio = 16.4.

In the title compound, $\text{C}_{15}\text{H}_{14}\text{N}_2\text{OS}$, the pyridinone ring in the fused-ring system is nearly planar (r.m.s. deviation = 0.011 Å) and the cyclohexene ring has a twisted half-boat conformation with the methylene C atom adjacent to the methine C atom deviating by 0.592 (7) Å from the plane defined by the remaining five atoms (r.m.s. deviation = 0.108 Å). The thienyl ring is disordered over two almost coplanar positions of opposite orientation in a 0.649 (4):0.351 (4) ratio, and forms dihedral angles of 51.4 (3) (major component) and 54.2 (3)°, respectively, with the pyridinone ring. In the crystal, inversion-related molecules associate *via* an eight-membered $\{\cdots\text{HNCO}\}_2$ synthon and these are linked into a linear supramolecular chain along the a axis by weak $\pi-\pi$ interactions that occur between centrosymmetrically related pyridinone rings [centroid-centroid distance = 3.889 (2) Å].

Related literature

For background to the cardiogenic and anti-inflammatory properties of this class of compounds, see: Behit & Baraka (2005); Girgis *et al.* (2007). For a related structure, see: Asiri *et al.* (2011).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{14}\text{N}_2\text{OS}$
 $M_r = 270.34$
Triclinic, $P\bar{1}$
 $a = 7.6443$ (3) Å
 $b = 9.6909$ (5) Å
 $c = 9.9852$ (5) Å
 $\alpha = 67.003$ (5)°
 $\beta = 80.869$ (4)°
 $\gamma = 76.108$ (4)°
 $V = 659.26$ (5) Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.24$ mm⁻¹
 $T = 100$ K
0.30 × 0.20 × 0.05 mm

Data collection

Agilent SuperNova Dual
diffractometer with an Atlas
detector
Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2012)
 $T_{\min} = 0.798$, $T_{\max} = 1.000$
9707 measured reflections
3041 independent reflections
2356 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.088$
 $wR(F^2) = 0.230$
 $S = 1.07$
3041 reflections
185 parameters
33 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 1.02$ e Å⁻³
 $\Delta\rho_{\min} = -0.78$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}n\cdots\text{O1}^i$	0.88	1.94	2.801 (4)	168

Symmetry code: (i) $-x + 2, -y + 1, -z + 1$.

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors are grateful to King Abdulaziz University for providing the research facilities. The authors also thank the Ministry of Higher Education (Malaysia) for funding structural studies through the High-Impact Research scheme (UM.C/HIR/MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5572).

‡ Additional correspondence author, e-mail: aasiri2@kau.edu.sa.

References

- Agilent (2012). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.
- Asiri, A. M., Faidallah, H. M., Al-Youbi, A. O., Alamry, K. A. & Ng, S. W. (2011). *Acta Cryst.* **E67**, o2472.
- Behit, A. A. & Baraka, A. M. (2005). *Eur. J. Med. Chem.* **40**, 1405–1413.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Girgis, A. S., Mishriky, N., Ellithey, M., Hosni, H. M. & Farag, H. (2007). *Bioorg. Med. Chem.* **15**, 2403–2413.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2012). E68, o2291–o2292 [https://doi.org/10.1107/S160053681202836X]

8-Methyl-2-oxo-4-(thiophen-2-yl)-1,2,5,6,7,8-hexahydroquinoline-3-carbonitrile

Abdullah M. Asiri, Hassan M. Faidallah, Alaa Anwar Ahmad Saqer, Seik Weng Ng and Edward R. T. Tiekink

S1. Comment

The title compound (I) is a member of a series of cyano-pyridinones that have been evaluated for their cardiotoxic (Behit & Baraka, 2005) and anti-inflammatory (Girgis *et al.*, (2007) properties. Herein, the crystal and molecular structures of (I) are described.

In (I), Fig. 1, the pyridyl ring in the fused ring system is planar [r.m.s. deviation = 0.011 Å] and the cyclohexene ring has a twisted half-boat conformation with the methylene-C3 atom lying -0.592 (7) Å above the plane defined by the remaining five atoms [r.m.s. deviation = 0.108 Å]. There are two orientations of the thienyl ring [co-planar, with a dihedral angle of 4.4 (4)°, and of opposite orientations] both of which are inclined with respect to the pyridyl ring, forming dihedral angles of 51.4 (3)° [major component] and 54.2 (3)°, respectively. The molecular structure of (I) resembles that found in a literature structure with the exception of the C2—C3 conformation which is fused to a benzene ring (Asiri *et al.*, 2011).


The familiar eight-membered centrosymmetric amide {…HNCO}₂ synthon is observed in the crystal packing, Table 1. These are connected into a linear supramolecular chain along the *a* axis by π — π interactions that occur between centrosymmetrically related pyridyl rings [inter-centroid distance = 3.889 (2) Å for symmetry operation 1 - *x*, 1 - *y*, 1 - *z*], Fig. 2. Chains assemble into layers in the *ab* plane and stack along the *c* axis being separated by hydrophobic interactions, Fig. 3.

S2. Experimental


A mixture of the thiophene-2-carboxaldehyde (1.1 g, 0.01 *M*), 2-methylcyclohexanone (1.12 g, 0.01 *M*), ethyl cyanoacetate (1.1 g, 0.01 *M*) and ammonium acetate (6.2 g, 0.08 *M*) in absolute ethanol (50 ml) was refluxed for 6 h. The reaction mixture was allowed to cool. The formed precipitate was filtered, washed with water, dried and recrystallized from ethanol as yellow crystals, *M. pt.*: 525–527 K. Yield: 72%.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions [C—H = 0.95–0.99 Å, $U_{\text{iso}}(\text{H}) = 1.2$ – $1.5U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation. The N-bound H-atom was treated similarly with N—H = 0.88 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$. The thienyl ring is disordered over two positions [co-planar and opposite orientation] in a 0.649 (4):0.351 (4) ratio. Pairs of 1,2-related distances were restrained to within 0.01 Å of each other, and the rings were restrained to be within 0.01 Å of a plane. The anisotropic displacement parameters, restrained to be nearly isotropic, of the primed atoms were set to those of the unprimed ones. The maximum and minimum residual electron density peaks of 1.02 and -0.78 e Å⁻³, respectively, were located 0.14 Å and 0.39 Å from the C15' and C13 atoms, respectively.

**Figure 1**

The molecular structure of (I) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.

**Figure 2**

A view of the supramolecular chain along the *a* axis in (I) mediated by N—H...O and π — π interactions, shown as orange and purple dashed lines, respectively.




Figure 3

A view in projection down the b axis of the unit-cell contents of (I). The N—H \cdots O and π — π interactions are shown as orange and purple dashed lines, respectively.

8-Methyl-2-oxo-4-(thiophen-2-yl)-1,2,5,6,7,8-hexahydroquinoline-3-carbonitrile

Crystal data

$C_{15}H_{14}N_2OS$

$M_r = 270.34$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 7.6443\ (3)\ \text{\AA}$

$b = 9.6909\ (5)\ \text{\AA}$

$c = 9.9852\ (5)\ \text{\AA}$

$\alpha = 67.003\ (5)^\circ$

$\beta = 80.869\ (4)^\circ$

$\gamma = 76.108\ (4)^\circ$

$V = 659.26\ (5)\ \text{\AA}^3$

$Z = 2$

$F(000) = 284$

$D_x = 1.362\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 3729 reflections

$\theta = 2.3\text{--}27.5^\circ$

$\mu = 0.24\ \text{mm}^{-1}$

$T = 100\ \text{K}$

Plate, yellow

$0.30 \times 0.20 \times 0.05\ \text{mm}$

Data collection

Agilent SuperNova Dual
 diffractometer with an Atlas detector
 Radiation source: SuperNova (Mo) X-ray
 Source
 Mirror monochromator
 Detector resolution: 10.4041 pixels mm⁻¹
 ω scan
 Absorption correction: multi-scan
 (CrysAlis PRO; Agilent, 2012)

$T_{\min} = 0.798$, $T_{\max} = 1.000$
 9707 measured reflections
 3041 independent reflections
 2356 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$
 $\theta_{\max} = 27.6^\circ$, $\theta_{\min} = 2.3^\circ$
 $h = -9 \rightarrow 9$
 $k = -12 \rightarrow 12$
 $l = -12 \rightarrow 12$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.088$
 $wR(F^2) = 0.230$
 $S = 1.07$
 3041 reflections
 185 parameters
 33 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0899P)^2 + 1.5187P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 1.02 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.78 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.8432 (4)	0.6552 (5)	0.4105 (4)	0.0607 (13)	
N1	0.7871 (4)	0.4615 (4)	0.6240 (4)	0.0359 (9)	
H1n	0.8984	0.4118	0.6160	0.043*	
N2	0.4540 (5)	0.9325 (4)	0.3270 (4)	0.0359 (9)	
C1	0.9207 (6)	0.2563 (6)	0.9181 (5)	0.0405 (11)	
H1A	1.0051	0.3102	0.8423	0.061*	
H1B	0.9847	0.1546	0.9765	0.061*	
H1C	0.8702	0.3144	0.9813	0.061*	
C2	0.7691 (5)	0.2402 (5)	0.8473 (4)	0.0303 (9)	
H2	0.8246	0.1755	0.7884	0.036*	
C3	0.6307 (6)	0.1585 (5)	0.9600 (5)	0.0337 (10)	
H3A	0.5668	0.1120	0.9147	0.040*	
H3B	0.6941	0.0752	1.0422	0.040*	
C4	0.4947 (6)	0.2674 (5)	1.0173 (5)	0.0361 (10)	
H4A	0.5584	0.3154	1.0611	0.043*	
H4B	0.4121	0.2101	1.0945	0.043*	

C5	0.3854 (5)	0.3914 (5)	0.8953 (4)	0.0300 (9)	
H5A	0.3015	0.3458	0.8667	0.036*	
H5B	0.3120	0.4703	0.9318	0.036*	
C6	0.5042 (5)	0.4659 (4)	0.7629 (4)	0.0219 (7)	
C7	0.6793 (5)	0.3935 (4)	0.7432 (4)	0.0244 (8)	
C8	0.7363 (5)	0.6006 (5)	0.5159 (5)	0.0367 (11)	
C9	0.5537 (5)	0.6763 (4)	0.5364 (4)	0.0252 (8)	
C10	0.4413 (5)	0.6133 (4)	0.6569 (4)	0.0204 (7)	
C11	0.4972 (5)	0.8193 (5)	0.4214 (4)	0.0258 (8)	
C12	0.2592 (5)	0.7030 (4)	0.6738 (4)	0.0311 (9)	
S1	0.0646 (2)	0.6528 (2)	0.7033 (2)	0.0426 (7)	0.649 (4)
S1'	0.2014 (7)	0.8647 (5)	0.6683 (4)	0.043*	0.351 (4)
C13	0.260 (2)	0.8772 (14)	0.6570 (7)	0.069 (4)	0.649 (4)
H13	0.3559	0.9306	0.6399	0.082*	0.649 (4)
C13'	0.079 (2)	0.627 (2)	0.6905 (12)	0.069*	0.351 (4)
H13'	0.0714	0.5285	0.6971	0.082*	0.351 (4)
C14	0.0544 (12)	0.9255 (9)	0.6783 (8)	0.0461 (18)	0.649 (4)
H14	0.0031	1.0261	0.6752	0.055*	0.649 (4)
C14'	-0.072 (2)	0.7647 (18)	0.6921 (14)	0.046*	0.351 (4)
H14'	-0.1961	0.7600	0.6992	0.055*	0.351 (4)
C15	-0.0505 (11)	0.8293 (9)	0.7006 (7)	0.0432 (18)	0.649 (4)
H15	-0.1783	0.8545	0.7141	0.052*	0.649 (4)
C15'	-0.0217 (15)	0.8881 (17)	0.6833 (11)	0.043*	0.351 (4)
H15'	-0.1023	0.9788	0.6852	0.052*	0.351 (4)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0203 (15)	0.071 (3)	0.0354 (18)	0.0116 (15)	0.0094 (13)	0.0234 (17)
N1	0.0157 (15)	0.041 (2)	0.0255 (18)	0.0083 (14)	0.0040 (13)	0.0050 (15)
N2	0.0270 (18)	0.035 (2)	0.0275 (18)	0.0013 (15)	-0.0005 (14)	0.0029 (15)
C1	0.026 (2)	0.045 (3)	0.031 (2)	-0.0043 (18)	-0.0045 (17)	0.0055 (19)
C2	0.026 (2)	0.027 (2)	0.025 (2)	0.0028 (15)	-0.0006 (15)	-0.0016 (16)
C3	0.031 (2)	0.0230 (19)	0.032 (2)	-0.0038 (16)	-0.0018 (17)	0.0049 (16)
C4	0.030 (2)	0.034 (2)	0.025 (2)	-0.0049 (17)	0.0056 (16)	0.0047 (17)
C5	0.0189 (18)	0.030 (2)	0.029 (2)	-0.0054 (15)	0.0050 (15)	0.0000 (16)
C6	0.0180 (17)	0.0239 (18)	0.0205 (17)	-0.0050 (13)	-0.0003 (13)	-0.0046 (14)
C7	0.0197 (17)	0.0267 (19)	0.0202 (17)	-0.0022 (14)	0.0012 (13)	-0.0039 (15)
C8	0.0157 (18)	0.045 (2)	0.024 (2)	0.0044 (16)	0.0023 (14)	0.0071 (18)
C9	0.0174 (17)	0.0285 (19)	0.0209 (18)	-0.0013 (14)	-0.0021 (13)	-0.0013 (15)
C10	0.0151 (16)	0.0236 (17)	0.0213 (17)	-0.0047 (13)	-0.0007 (13)	-0.0067 (14)
C11	0.0164 (17)	0.031 (2)	0.0221 (18)	-0.0023 (14)	0.0006 (13)	-0.0034 (16)
C12	0.0238 (19)	0.035 (2)	0.0189 (18)	0.0049 (16)	0.0027 (14)	-0.0018 (16)
S1	0.0090 (7)	0.0483 (10)	0.0447 (10)	-0.0033 (6)	-0.0002 (5)	0.0084 (7)
C13	0.069 (4)	0.069 (4)	0.068 (4)	-0.0132 (15)	-0.0044 (13)	-0.0250 (18)
C14	0.046 (2)	0.045 (2)	0.043 (2)	-0.0018 (12)	0.0006 (12)	-0.0171 (13)
C15	0.040 (2)	0.042 (2)	0.041 (2)	-0.0030 (12)	0.0010 (12)	-0.0128 (13)

Geometric parameters (Å, °)

O1—C8	1.246 (5)	C6—C10	1.429 (5)
N1—C7	1.366 (5)	C8—C9	1.436 (5)
N1—C8	1.371 (5)	C9—C10	1.381 (5)
N1—H1n	0.8800	C9—C11	1.433 (5)
N2—C11	1.148 (5)	C10—C12	1.477 (5)
C1—C2	1.516 (6)	C12—S1'	1.503 (5)
C1—H1A	0.9800	C12—S1	1.622 (4)
C1—H1B	0.9800	C12—C13	1.632 (13)
C1—H1C	0.9800	C12—C13'	1.668 (17)
C2—C7	1.513 (5)	S1—C15	1.717 (8)
C2—C3	1.530 (5)	S1'—C15'	1.656 (11)
C2—H2	1.0000	C13—C14	1.537 (15)
C3—C4	1.510 (6)	C13—H13	0.9500
C3—H3A	0.9900	C13'—C14'	1.541 (18)
C3—H3B	0.9900	C13'—H13'	0.9500
C4—C5	1.524 (6)	C14—C15	1.304 (12)
C4—H4A	0.9900	C14—H14	0.9500
C4—H4B	0.9900	C14'—C15'	1.309 (15)
C5—C6	1.511 (5)	C14'—H14'	0.9500
C5—H5A	0.9900	C15—H15	0.9500
C5—H5B	0.9900	C15'—H15'	0.9500
C6—C7	1.377 (5)		
C7—N1—C8	125.4 (3)	O1—C8—N1	121.6 (4)
C7—N1—H1n	117.3	O1—C8—C9	124.0 (4)
C8—N1—H1n	117.3	N1—C8—C9	114.4 (3)
C2—C1—H1A	109.5	C10—C9—C11	123.1 (3)
C2—C1—H1B	109.5	C10—C9—C8	122.1 (3)
H1A—C1—H1B	109.5	C11—C9—C8	114.9 (3)
C2—C1—H1C	109.5	C9—C10—C6	119.8 (3)
H1A—C1—H1C	109.5	C9—C10—C12	118.5 (3)
H1B—C1—H1C	109.5	C6—C10—C12	121.7 (3)
C7—C2—C1	111.3 (4)	N2—C11—C9	178.5 (4)
C7—C2—C3	111.3 (3)	C10—C12—S1'	130.3 (4)
C1—C2—C3	112.2 (4)	C10—C12—S1	129.1 (3)
C7—C2—H2	107.3	S1'—C12—S1	100.5 (3)
C1—C2—H2	107.3	C10—C12—C13	113.3 (6)
C3—C2—H2	107.3	S1—C12—C13	117.5 (6)
C4—C3—C2	111.4 (3)	C10—C12—C13'	119.2 (8)
C4—C3—H3A	109.4	S1'—C12—C13'	110.3 (8)
C2—C3—H3A	109.4	C13—C12—C13'	127.1 (11)
C4—C3—H3B	109.4	C12—S1—C15	92.6 (4)
C2—C3—H3B	109.4	C12—S1'—C15'	102.2 (6)
H3A—C3—H3B	108.0	C14—C13—C12	95.8 (9)
C3—C4—C5	110.8 (4)	C14—C13—H13	132.1
C3—C4—H4A	109.5	C12—C13—H13	132.1

C5—C4—H4A	109.5	C14'—C13'—C12	99.6 (13)
C3—C4—H4B	109.5	C14'—C13'—H13'	130.2
C5—C4—H4B	109.5	C12—C13'—H13'	130.2
H4A—C4—H4B	108.1	C15—C14—C13	120.5 (8)
C6—C5—C4	112.3 (3)	C15—C14—H14	119.8
C6—C5—H5A	109.2	C13—C14—H14	119.8
C4—C5—H5A	109.2	C15'—C14'—C13'	117.0 (15)
C6—C5—H5B	109.2	C15'—C14'—H14'	121.5
C4—C5—H5B	109.2	C13'—C14'—H14'	121.5
H5A—C5—H5B	107.9	C14—C15—S1	113.6 (6)
C7—C6—C10	118.1 (3)	C14—C15—H15	123.2
C7—C6—C5	120.2 (3)	S1—C15—H15	123.2
C10—C6—C5	121.7 (3)	C14'—C15'—S1'	110.9 (12)
N1—C7—C6	120.2 (3)	C14'—C15'—H15'	124.6
N1—C7—C2	114.9 (3)	S1'—C15'—H15'	124.6
C6—C7—C2	124.9 (3)		
C7—C2—C3—C4	-42.7 (5)	C9—C10—C12—S1'	-51.3 (6)
C1—C2—C3—C4	82.7 (5)	C6—C10—C12—S1'	127.3 (4)
C2—C3—C4—C5	62.8 (5)	C9—C10—C12—S1	128.4 (4)
C3—C4—C5—C6	-49.6 (5)	C6—C10—C12—S1	-52.9 (5)
C4—C5—C6—C7	19.3 (6)	C9—C10—C12—C13	-50.5 (4)
C4—C5—C6—C10	-159.4 (4)	C6—C10—C12—C13	128.2 (4)
C8—N1—C7—C6	-0.6 (7)	C9—C10—C12—C13'	123.6 (5)
C8—N1—C7—C2	-179.4 (4)	C6—C10—C12—C13'	-57.7 (5)
C10—C6—C7—N1	-0.8 (6)	C10—C12—S1—C15	-178.7 (4)
C5—C6—C7—N1	-179.5 (4)	S1'—C12—S1—C15	1.1 (3)
C10—C6—C7—C2	177.8 (4)	C13—C12—S1—C15	0.15 (17)
C5—C6—C7—C2	-1.0 (6)	C13'—C12—S1—C15	-155 (3)
C1—C2—C7—N1	65.3 (5)	C10—C12—S1'—C15'	175.7 (5)
C3—C2—C7—N1	-168.7 (4)	S1—C12—S1'—C15'	-4.1 (4)
C1—C2—C7—C6	-113.3 (5)	C13—C12—S1'—C15'	173.1 (11)
C3—C2—C7—C6	12.6 (6)	C13'—C12—S1'—C15'	0.42 (19)
C7—N1—C8—O1	179.8 (5)	C10—C12—C13—C14	178.9 (4)
C7—N1—C8—C9	0.7 (7)	S1'—C12—C13—C14	-3.2 (10)
O1—C8—C9—C10	-178.3 (5)	S1—C12—C13—C14	-0.1 (2)
N1—C8—C9—C10	0.8 (7)	C13'—C12—C13—C14	5.4 (5)
O1—C8—C9—C11	3.1 (7)	C10—C12—C13'—C14'	-175.7 (4)
N1—C8—C9—C11	-177.9 (4)	S1'—C12—C13'—C14'	0.2 (2)
C11—C9—C10—C6	176.3 (4)	S1—C12—C13'—C14'	25 (3)
C8—C9—C10—C6	-2.2 (6)	C13—C12—C13'—C14'	-2.5 (5)
C11—C9—C10—C12	-5.0 (6)	C12—C13—C14—C15	0.0 (4)
C8—C9—C10—C12	176.4 (4)	C12—C13'—C14'—C15'	-1.0 (5)
C7—C6—C10—C9	2.2 (5)	C13—C14—C15—S1	0.1 (6)
C5—C6—C10—C9	-179.1 (4)	C12—S1—C15—C14	-0.2 (4)
C7—C6—C10—C12	-176.4 (4)	C13'—C14'—C15'—S1'	1.3 (6)
C5—C6—C10—C12	2.3 (6)	C12—S1'—C15'—C14'	-1.0 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1 <i>n</i> ···O1 ⁱ	0.88	1.94	2.801 (4)	168

Symmetry code: (i) $-x+2, -y+1, -z+1$.