

**1-(3-Fluorophenyl)-3-(4-nitrophenyl)urea****Mu-Sen Lin,<sup>a</sup> Yu Shi,<sup>b</sup> Shi-Yu Zhang<sup>a</sup> and Yi-Liang Li<sup>a\*</sup>**

<sup>a</sup>School of Chinese Materia Medica, Tianjin University of Traditional Chinese Medicine, Tianjin, 300193, People's Republic of China, and <sup>b</sup>Tianjin Key Laboratory of Molecular Design and Drug Discovery, Tianjin Institute of Pharmaceutical Research, Tianjin, 300193, People's Republic of China  
Correspondence e-mail: liyl@tjipr.com


Received 12 April 2012; accepted 1 June 2012

Key indicators: single-crystal X-ray study;  $T = 113\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.052;  $wR$  factor = 0.131; data-to-parameter ratio = 15.3.

In the title compound,  $\text{C}_{13}\text{H}_{10}\text{FN}_3\text{O}_3$ , the dihedral angle between the fluorophenyl and nitrophenyl ring planes is  $6.51(9)^\circ$ . The crystal structure features  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds.

**Related literature**

The title compound is an activated fragment of sorafenib derivatives. Sorafenib is a VEGFR-2 inhibitor (Ferrara *et al.*, 2003; Peruzzi *et al.*, 2006) that has good therapeutic effect for renal carcinoma and liver cancer (Wan *et al.*, 2004; Wilhelm *et al.*, 2004).

**Experimental***Crystal data*

$\text{C}_{13}\text{H}_{10}\text{FN}_3\text{O}_3$	$V = 1219.5(11)\text{ \AA}^3$
$M_r = 275.24$	$Z = 4$
Monoclinic, $P2_1/n$	$\text{Mo } K\alpha$ radiation
$a = 8.351(4)\text{ \AA}$	$\mu = 0.12\text{ mm}^{-1}$
$b = 12.461(6)\text{ \AA}$	$T = 113\text{ K}$
$c = 11.912(6)\text{ \AA}$	$0.24 \times 0.22 \times 0.20\text{ mm}$
$\beta = 100.315(9)^\circ$	

**Data collection**

Rigaku Saturn CCD area-detector diffractometer	12466 measured reflections
Absorption correction: multi-scan	2900 independent reflections
<i>CrystalClear</i>	2459 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.045$
	$T_{\min} = 0.972$ , $T_{\max} = 0.977$

**Refinement**

$R[F^2 > 2\sigma(F^2)] = 0.052$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.131$	$\Delta\rho_{\max} = 0.30\text{ e \AA}^{-3}$
$S = 1.12$	$\Delta\rho_{\min} = -0.26\text{ e \AA}^{-3}$
2900 reflections	2 restraints
189 parameters	

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2A $\cdots$ O1 <sup>i</sup>	0.91 (1)	1.99 (1)	2.890 (2)	170 (2)
N3—H3A $\cdots$ O2 <sup>i</sup>	0.90 (1)	2.28 (1)	3.157 (2)	168 (2)

Symmetry code: (i)  $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL/PC* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL/PC*.

This work was supported by the Tianjin Natural Science Foundation (09JCZDJC21700).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZJ2072).

**References**

- Ferrara, N., Gerber, H. P. & Le, C. J. (2003). *Nat. Med.*, **9**, 669–679.
- Peruzzi, B. & Bottaro, D. P. (2006). *Clin. Cancer Res.*, **12**, 3657–3660.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Wan, P. T. C., Garnett, M. J., Roe, S. M., Lee, S., Niculescu-Duvaz, D., Good, V. M., Jones, C. M., Marshall, C. J., Springer, C. J., Barford, D. & Marais, R. (2004). *Cell*, **116**, 855–867.
- Wilhelm, S. M., Carter, C., Tang, L. Y., Wilkie, D., McNabola, A., Rong, H., Chen, C., Zhang, X. M., Vincent, P., McHugh, M., Cao, Y. C., Shujath, J., Gawlak, S., Eveleigh, D., Rowley, B., Liu, L., Adnane, L., Lynch, M., Auclair, D., Taylor, I., Gedrich, R., Voznesensky, A., Riedl, B., Post, L. E., Bollag, G. & Trail, P. A. (2004). *Cancer Res.*, **64**, 7099–7109.

# supporting information

*Acta Cryst.* (2012). E68, o2030 [https://doi.org/10.1107/S160053681202507X]

## 1-(3-Fluorophenyl)-3-(4-nitrophenyl)urea

Mu-Sen Lin, Yu Shi, Shi-Yu Zhang and Yi-Liang Li

### S1. Comment

Sorafenib is a VEGFR-2 inhibitor (Ferrara *et al.*, 2003; Peruzzi *et al.*, 2006) that has good therapeutic effect for renal carcinoma and liver cancer (Wan *et al.*, 2004; Wilhelm *et al.*, 2004). 1-(3-fluorophenyl)-3-(4-nitrophenyl) urea is an important activated fragment of sorafenib derivatives. We present here the structure characterization of the title compound.


In the molecule of the title compound (Fig.1) bond lengths and angles have normal values. The interplanar angle between the fluorobenzyl and nitrobenzyl ring planes is 6.51 (9) $^{\circ}$ . The crystal structure is stabilized by the intermolecular N—H $\cdots$ O hydrogen bonds. The crystal structure (Fig.2) is stabilized by intermolecular N—H $\cdots$ O hydrogen bonds (table 1).

### S2. Experimental


A solution of 4-nitroaniline (1.38 g, 10 mmol) in DCM (100 ml) was added dropwise to a stirred solution of bis(trichloromethyl) carbonate (5.92 g, 20 mmol) in DCM (20 ml) at the atmosphere of ice-bath. The reaction mixture was stirred for 2 hrs at 0–5°C. Then the reaction mixture was added dropwise to a refluxed and stirred solution of 3-fluoroaniline (1.11 g, 10 mmol) in DCM (40 ml). The reaction was completed within 2 hrs at the reflux temperature. The solvent was removed under reduced pressure. Acetone (100 ml) and H<sub>2</sub>O (300 ml) was added to the mixture. The solid was collected and washed with H<sub>2</sub>O, then gave a yellow solid. The yield was 2.08 g (75.6%). Put about 0.3 g of the product in the ampoule bottle and add 10 ml absolute ethyl alcohol, yellow single crystals suitable for X-ray diffraction analysis were obtained by slow evaporation of the solvent at room temperature after 3 weeks.

### S3. Refinement

All H atoms were detected in a difference map, nevertheless, the H-atoms attached to the nitrogen atoms were refined freely, and the H-atoms attached to the carbon atoms were placed in calculated positions and refined using a riding motion approximation, with C—H=0.95 Å, with  $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C})$ .

**Figure 1**

The molecular structure of the title compound, showing the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

**Figure 2**

Packing diagram of the title compound viewed along the *c* axis. Hydrogen bonds are shown as dashed lines.

### 1-(3-Fluorophenyl)-3-(4-nitrophenyl)urea

#### *Crystal data*



$M_r = 275.24$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 8.351 (4)$  Å

$b = 12.461 (6)$  Å

$c = 11.912 (6)$  Å

$\beta = 100.315 (9)^\circ$

$V = 1219.5 (11)$  Å<sup>3</sup>

$Z = 4$

$F(000) = 568$

$D_x = 1.499 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 4180 reflections

$\theta = 1.6\text{--}27.8^\circ$

$\mu = 0.12 \text{ mm}^{-1}$

$T = 113$  K

Prism, yellow

$0.24 \times 0.22 \times 0.20$  mm

*Data collection*

Rigaku Saturn CCD area-detector  
diffractometer  
Radiation source: rotating anode  
Multilayer monochromator  
Detector resolution: 14.63 pixels mm<sup>-1</sup>  
 $\omega$  and  $\varphi$  scans  
Absorption correction: multi-scan  
*CrystalClear*  
 $T_{\min} = 0.972$ ,  $T_{\max} = 0.977$

12466 measured reflections  
2900 independent reflections  
2459 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.045$   
 $\theta_{\max} = 27.8^\circ$ ,  $\theta_{\min} = 2.4^\circ$   
 $h = -10 \rightarrow 10$   
 $k = -16 \rightarrow 16$   
 $l = -15 \rightarrow 15$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.052$   
 $wR(F^2) = 0.131$   
 $S = 1.12$   
2900 reflections  
189 parameters  
2 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H atoms treated by a mixture of independent  
and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.063P)^2 + 0.1263P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.003$   
 $\Delta\rho_{\max} = 0.30 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^* / U_{\text{eq}}$
F1	0.29184 (14)	0.96861 (8)	-0.02312 (9)	0.0449 (3)
O1	-0.04969 (14)	0.17411 (10)	0.34582 (10)	0.0322 (3)
O2	0.01132 (14)	0.04318 (10)	0.24132 (10)	0.0340 (3)
O3	0.19105 (13)	0.60131 (9)	0.05581 (9)	0.0277 (3)
N1	0.01105 (16)	0.13960 (12)	0.26533 (12)	0.0265 (3)
N2	0.26220 (16)	0.43681 (11)	-0.00884 (11)	0.0255 (3)
N3	0.32644 (17)	0.58967 (11)	-0.09569 (11)	0.0258 (3)
C1	0.20093 (18)	0.36633 (13)	0.06336 (13)	0.0236 (3)
C2	0.19815 (19)	0.25683 (13)	0.03495 (13)	0.0252 (4)
H2	0.2380	0.2346	-0.0312	0.030*
C3	0.13910 (19)	0.18127 (13)	0.10071 (14)	0.0264 (4)
H3	0.1371	0.1074	0.0808	0.032*
C4	0.08209 (18)	0.21605 (13)	0.19783 (13)	0.0232 (3)
C5	0.08776 (19)	0.32275 (13)	0.22924 (14)	0.0258 (4)
H5	0.0505	0.3439	0.2967	0.031*

C6	0.14717 (19)	0.39910 (13)	0.16335 (13)	0.0258 (4)
H6	0.1517	0.4725	0.1852	0.031*
C7	0.25437 (18)	0.54780 (13)	-0.01016 (13)	0.0234 (3)
C8	0.35025 (18)	0.69898 (13)	-0.11978 (13)	0.0246 (4)
C9	0.42473 (19)	0.72135 (14)	-0.21284 (14)	0.0268 (4)
H9	0.4554	0.6642	-0.2573	0.032*
C10	0.4543 (2)	0.82664 (14)	-0.24089 (14)	0.0297 (4)
H10	0.5056	0.8406	-0.3044	0.036*
C11	0.4106 (2)	0.91169 (15)	-0.17805 (15)	0.0327 (4)
H11	0.4305	0.9840	-0.1969	0.039*
C12	0.3367 (2)	0.88660 (13)	-0.08702 (15)	0.0307 (4)
C13	0.30351 (19)	0.78355 (14)	-0.05540 (14)	0.0274 (4)
H13	0.2509	0.7705	0.0077	0.033*
H2A	0.314 (2)	0.4069 (14)	-0.0618 (13)	0.037 (5)*
H3A	0.372 (2)	0.5429 (15)	-0.1377 (15)	0.056 (7)*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
F1	0.0653 (8)	0.0268 (6)	0.0451 (7)	-0.0026 (5)	0.0172 (6)	-0.0077 (5)
O1	0.0318 (6)	0.0391 (7)	0.0288 (7)	-0.0028 (5)	0.0136 (5)	0.0027 (5)
O2	0.0370 (7)	0.0269 (6)	0.0393 (7)	-0.0018 (5)	0.0096 (5)	0.0044 (5)
O3	0.0300 (6)	0.0275 (6)	0.0278 (6)	0.0027 (5)	0.0116 (5)	-0.0012 (5)
N1	0.0231 (7)	0.0283 (7)	0.0281 (7)	-0.0001 (6)	0.0044 (5)	0.0032 (6)
N2	0.0292 (7)	0.0241 (7)	0.0265 (7)	0.0010 (6)	0.0141 (6)	0.0009 (6)
N3	0.0303 (7)	0.0235 (7)	0.0265 (7)	0.0016 (6)	0.0130 (6)	0.0012 (6)
C1	0.0206 (7)	0.0276 (8)	0.0236 (8)	0.0013 (6)	0.0060 (6)	0.0015 (6)
C2	0.0259 (8)	0.0284 (8)	0.0228 (8)	0.0016 (6)	0.0084 (6)	-0.0001 (6)
C3	0.0257 (8)	0.0250 (8)	0.0287 (9)	0.0008 (6)	0.0056 (7)	-0.0005 (6)
C4	0.0213 (7)	0.0262 (8)	0.0228 (8)	-0.0009 (6)	0.0057 (6)	0.0044 (6)
C5	0.0259 (8)	0.0285 (9)	0.0250 (8)	0.0023 (6)	0.0097 (6)	-0.0002 (6)
C6	0.0281 (8)	0.0245 (8)	0.0267 (8)	0.0002 (6)	0.0099 (7)	-0.0007 (6)
C7	0.0205 (7)	0.0257 (8)	0.0239 (8)	0.0009 (6)	0.0042 (6)	0.0006 (6)
C8	0.0208 (7)	0.0254 (8)	0.0266 (8)	0.0015 (6)	0.0018 (6)	0.0022 (6)
C9	0.0250 (8)	0.0292 (9)	0.0263 (8)	0.0008 (6)	0.0051 (6)	0.0037 (7)
C10	0.0278 (8)	0.0320 (9)	0.0287 (9)	-0.0021 (7)	0.0032 (7)	0.0051 (7)
C11	0.0352 (9)	0.0275 (9)	0.0342 (9)	-0.0054 (7)	0.0027 (7)	0.0030 (7)
C12	0.0344 (9)	0.0250 (8)	0.0315 (9)	-0.0005 (7)	0.0029 (7)	-0.0035 (7)
C13	0.0286 (8)	0.0279 (8)	0.0257 (8)	-0.0018 (7)	0.0048 (7)	0.0011 (7)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

F1—C12	1.366 (2)	C3—H3	0.9500
O1—N1	1.2392 (18)	C4—C5	1.380 (2)
O2—N1	1.235 (2)	C5—C6	1.381 (2)
O3—C7	1.2209 (19)	C5—H5	0.9500
N1—C4	1.441 (2)	C6—H6	0.9500
N2—C7	1.384 (2)	C8—C9	1.393 (2)

N2—C1	1.389 (2)	C8—C13	1.399 (2)
N2—H2A	0.907 (9)	C9—C10	1.387 (2)
N3—C7	1.376 (2)	C9—H9	0.9500
N3—C8	1.413 (2)	C10—C11	1.384 (2)
N3—H3A	0.897 (9)	C10—H10	0.9500
C1—C2	1.405 (2)	C11—C12	1.376 (3)
C1—C6	1.407 (2)	C11—H11	0.9500
C2—C3	1.372 (2)	C12—C13	1.380 (2)
C2—H2	0.9500	C13—H13	0.9500
C3—C4	1.397 (2)		
O2—N1—O1	122.33 (14)	C5—C6—C1	118.87 (16)
O2—N1—C4	119.72 (14)	C5—C6—H6	120.6
O1—N1—C4	117.95 (14)	C1—C6—H6	120.6
C7—N2—C1	128.07 (14)	O3—C7—N3	124.54 (16)
C7—N2—H2A	115.5 (12)	O3—C7—N2	124.28 (15)
C1—N2—H2A	116.4 (12)	N3—C7—N2	111.18 (14)
C7—N3—C8	127.73 (14)	C9—C8—C13	119.58 (15)
C7—N3—H3A	116.9 (14)	C9—C8—N3	116.97 (15)
C8—N3—H3A	115.1 (14)	C13—C8—N3	123.46 (15)
N2—C1—C2	117.18 (14)	C10—C9—C8	120.35 (16)
N2—C1—C6	123.33 (15)	C10—C9—H9	119.8
C2—C1—C6	119.47 (15)	C8—C9—H9	119.8
C3—C2—C1	121.41 (15)	C11—C10—C9	121.22 (16)
C3—C2—H2	119.3	C11—C10—H10	119.4
C1—C2—H2	119.3	C9—C10—H10	119.4
C2—C3—C4	118.09 (15)	C12—C11—C10	116.83 (16)
C2—C3—H3	121.0	C12—C11—H11	121.6
C4—C3—H3	121.0	C10—C11—H11	121.6
C5—C4—C3	121.57 (15)	F1—C12—C11	118.40 (15)
C5—C4—N1	118.89 (14)	F1—C12—C13	117.07 (15)
C3—C4—N1	119.51 (15)	C11—C12—C13	124.52 (16)
C4—C5—C6	120.54 (15)	C12—C13—C8	117.49 (16)
C4—C5—H5	119.7	C12—C13—H13	121.3
C6—C5—H5	119.7	C8—C13—H13	121.3
C7—N2—C1—C2	168.79 (15)	C8—N3—C7—O3	4.1 (3)
C7—N2—C1—C6	-13.1 (2)	C8—N3—C7—N2	-176.20 (14)
N2—C1—C2—C3	-179.65 (14)	C1—N2—C7—O3	0.8 (3)
C6—C1—C2—C3	2.2 (2)	C1—N2—C7—N3	-178.89 (14)
C1—C2—C3—C4	-0.3 (2)	C7—N3—C8—C9	-179.05 (15)
C2—C3—C4—C5	-1.5 (2)	C7—N3—C8—C13	1.1 (3)
C2—C3—C4—N1	176.62 (13)	C13—C8—C9—C10	0.8 (2)
O2—N1—C4—C5	-176.89 (14)	N3—C8—C9—C10	-179.09 (14)
O1—N1—C4—C5	3.6 (2)	C8—C9—C10—C11	-0.2 (2)
O2—N1—C4—C3	4.9 (2)	C9—C10—C11—C12	0.0 (2)
O1—N1—C4—C3	-174.56 (13)	C10—C11—C12—F1	-179.98 (14)
C3—C4—C5—C6	1.5 (2)	C10—C11—C12—C13	-0.3 (3)

N1—C4—C5—C6	−176.63 (14)	F1—C12—C13—C8	−179.49 (14)
C4—C5—C6—C1	0.4 (2)	C11—C12—C13—C8	0.8 (3)
N2—C1—C6—C5	179.79 (14)	C9—C8—C13—C12	−1.0 (2)
C2—C1—C6—C5	−2.1 (2)	N3—C8—C13—C12	178.82 (15)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N2—H2A···O1 <sup>i</sup>	0.91 (1)	1.99 (1)	2.890 (2)	170 (2)
N3—H3A···O2 <sup>i</sup>	0.90 (1)	2.28 (1)	3.157 (2)	168 (2)

Symmetry code: (i)  $x+1/2, -y+1/2, z-1/2$ .