

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,2'-Sulfonyldipyrazine 4-oxide

Ai-Min Li, Ya Zhang, Zhi-Wei Wang and Chong-Qing Wan*

Department of Chemistry, Capital Normal University, Beijing 100048, People's Republic of China

Correspondence e-mail: wanchqing@yahoo.com.cn


Received 4 June 2012; accepted 23 June 2012

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.063; wR factor = 0.203; data-to-parameter ratio = 16.8.

In the title compound, $\text{C}_8\text{H}_6\text{N}_4\text{O}_3\text{S}$, the dihedral angle between the pyrazine rings is $85.04(1)^\circ$. In the crystal, molecules are arranged along the a axis and are linked by $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds and pyrazine–pyrazine $\pi-\pi$ interactions [centroid–centroid distance = $3.800(1)$ Å, forming an infinite chain array. The chains are connected by $\text{C}-\text{H}\cdots\text{O}(\text{oxide})$ hydrogen bonds into layers lying parallel to the ab plane. Along the c axis, the layers are stacked and linked through $\text{C}-\text{H}\cdots\text{O}(\text{sulfonyl})$ interactions, forming a three-dimensional network.

Related literature

For metal complexes with 2,2'-sulfonyldipyrazine, see: Wan & Mak (2011). For crystal structures of pyridyl-based N -oxide and their metal complexes, see: Jia *et al.* (2008).



Experimental

Crystal data

 $\text{C}_8\text{H}_6\text{N}_4\text{O}_3\text{S}$
 $M_r = 238.23$

 Monoclinic, $P2_1/c$
 $a = 7.6860(16)$ Å

 $b = 15.841(3)$ Å
 $c = 9.0624(14)$ Å
 $\beta = 117.813(13)^\circ$
 $V = 975.9(3)$ Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.33$ mm⁻¹
 $T = 296$ K
 $0.45 \times 0.30 \times 0.25$ mm

Data collection

 Bruker APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2007)
 $T_{\min} = 0.688$, $T_{\max} = 1.000$

 6606 measured reflections
 2429 independent reflections
 1586 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.058$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.063$
 $wR(F^2) = 0.203$
 $S = 1.07$
 2429 reflections

 145 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.85$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.46$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C2}-\text{H2A}\cdots\text{O1}^i$	0.93	2.32	3.130 (5)	146
$\text{C3}-\text{H3A}\cdots\text{O3}^{ii}$	0.93	2.56	3.419 (4)	153
$\text{C7}-\text{H7A}\cdots\text{N1}^{iii}$	0.93	2.57	3.449 (3)	157

 Symmetry codes: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x + 2, -y, -z + 1$; (iii) $-x + 2, -y + 1, -z + 1$.

Data collection: APEX2 (Bruker, 2007); cell refinement: APEX2 and SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

The authors are grateful for financial support from the Natural Science Foundation of Beijing Municipality (grant No. 2122011) and the Beijing Municipal Education Commission (KM201210028018) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZQ2171).

References

- Bruker (2007). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Jia, J. H., Blake, A. J., Champness, N. R., Hubberstey, P., Wilson, C. & Schröder, M. (2008). *Inorg. Chem.* **47**, 8652–8664.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Wan, C. Q. & Mak, T. C. W. (2011). *New J. Chem.* **35**, 319–327.

supporting information

Acta Cryst. (2012). E68, o2285 [https://doi.org/10.1107/S1600536812028607]

2,2'-Sulfonyldipyrazine 4-oxide

Ai-Min Li, Ya Zhang, Zhi-Wei Wang and Chong-Qing Wan

S1. Comment

Pyridyl based sulfonyl derivatives were widely used in supramolecular assemblies of transition metal complexes (Wan & Mak, 2011). Pyridyl based N-oxide derivatives have also been demonstrated as versatile building blocks to construct supramolecular architectures of various metal complexes (Jia *et al.*, 2008). In the present context, we report the structure of the title compound, a new N-oxide compound derived from 2,2'-sulfonyldipyrazine.

In the title compound, the value of the C1(sp²)—S1—C5(sp²) angle is 103.92 (1)° with two attached pyrazinyl rings exhibiting a dihedral angle of 85.04 (1)°, as shown in Fig. 1. The angular-shaped molecules are arranged along the *a* axis. As shown in Fig. 2, two adjacent molecules arranged with an inversion center are interconnected through C7—H7A···N1ⁱⁱⁱ and $\pi\cdots\pi$ interactions (Cg···Cgⁱⁱⁱ = 3.800 (1) Å, Cg represents the C5-N3-C6-C7-N4-C8 ring; symmetry code: *iii* = 2 - *x*, 1 - *y*, 1 - *z*). The dimers are further interconnected through $\pi\cdots\pi$ interactions between Cg and Cg^{iv} [Cg···Cg^{iv} = 4.174 (2) Å, the distance between the closest ring atom and one Cg is 3.597 (2) Å; symmetry code = 1 - *x*, 1 - *y*, 1 - *z*]. The formed chains are further connected through C3—H3A···O3ⁱⁱ(oxynitride) hydrogen bonds to form a layer almost parallel to the *ab* plane (symmetry code: *ii* = 2 - *x*, -*y*, 1 - *z*). Along the *c* axis, the formed layers are stacked and interconnected through C2—H2A···O1ⁱ(sulfonyl) interactions to form a three-dimensional framework (Fig. 3, Table 1; symmetry code: *i* = *x*, -*y* + 1/2, *z* + 1/2).

S2. Experimental

The title compound was obtained as a serendipitous byproduct as the 2,2'-dipyrazine sulfide (0.022 g, 0.1 mmol) was dissolved in a mixture of methanol 2 ml and acetonitrile 2 ml to react with Mn(ClO₄)₂·6H₂O (0.036 g, 0.1 mmol) with constantly stirring at room temperature. After three hours, the clear solution was filtrated and kept in air for about one week to yield colourless block crystals (7 mg, 29% yield). We got the the title compound as a matter of the oxidability by perchlorate acid from Mn(ClO₄)₂·6H₂O.

S3. Refinement

All hydrogen positions were calculated after each cycle of refinement using a riding model with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The highest peak (0.8 e.Å⁻³) in the difference Fourier map is located at 1.1 Å from atom N4. The refinement of a model including one H atom at this position led to lower R₁ and wR₂ values but it is chemically meaningless since there is no counter ion in the crystal structure. A positional disorder of the oxo O atom (partially on atoms O4 and O2) is surely the best solution but in this case too many restraints had to be used in the final refinements to get an acceptable model (with an site-occupancy ratio greater than 0.9:0.1).




Figure 1

The atom-numbering scheme of the title complex. Displacement ellipsoids are drawn at the 35% probability level and H atoms are shown as sticks of arbitrary radii.




Figure 2

The hydrogen-bonding (C—H...N) and π ... π stacking interactions between parallel chains along the a axis, which are respectively shown as thin red-dashed lines and thick blue-dashed lines (symmetry codes: $i = -x + 2, -y + 1, -z + 1$; $ii = x + 1, y, z$).




Figure 3

Three-dimensional structure of the title molecule viewed down the *a* direction. The red dashed lines represent hydrogen-bonding interactions.

2,2'-Sulfonyldipyrazine 4-oxide

Crystal data

$C_8H_6N_4O_3S$

$M_r = 238.23$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 7.6860$ (16) Å

$b = 15.841$ (3) Å

$c = 9.0624$ (14) Å

$\beta = 117.813$ (13)°

$V = 975.9$ (3) Å³

$Z = 4$

$F(000) = 488$

$D_x = 1.621$ Mg m⁻³

$D_m = 1.621$ Mg m⁻³

D_m measured by not measured

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 365 reflections

$\theta = 2.6$ – 28.4 °

$\mu = 0.33$ mm⁻¹

$T = 296$ K

Needle-like, colourless

$0.45 \times 0.30 \times 0.25$ mm

Data collection

Bruker APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2007)

$T_{\min} = 0.688$, $T_{\max} = 1.000$

6606 measured reflections

2429 independent reflections

1586 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.058$

$\theta_{\max} = 28.4$ °, $\theta_{\min} = 2.6$ °

$h = -10 \rightarrow 10$

$k = -21 \rightarrow 11$

$l = -12 \rightarrow 11$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.063$

$wR(F^2) = 0.203$

$S = 1.07$

2429 reflections

145 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0981P)^2 + 0.5688P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.85$ e Å⁻³

$\Delta\rho_{\min} = -0.46$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.50844 (11)	0.33579 (5)	0.25247 (12)	0.0436 (3)
N1	0.8484 (4)	0.27476 (19)	0.2912 (4)	0.0531 (8)
N2	0.8080 (4)	0.12101 (18)	0.4199 (4)	0.0520 (8)
N3	0.6907 (4)	0.4321 (2)	0.5196 (4)	0.0540 (8)
N4	0.7631 (6)	0.5635 (2)	0.3499 (6)	0.0699 (10)
O1	0.4354 (3)	0.35188 (18)	0.0788 (3)	0.0581 (7)
O2	0.3755 (4)	0.31172 (17)	0.3141 (4)	0.0627 (8)
O3	0.7852 (6)	0.0494 (2)	0.4742 (6)	0.0996 (13)
C1	0.6949 (4)	0.25624 (19)	0.3152 (4)	0.0403 (7)
C2	0.6667 (5)	0.1820 (2)	0.3776 (5)	0.0494 (9)
H2A	0.5552	0.1732	0.3910	0.059*
C3	0.9681 (5)	0.1382 (2)	0.4026 (5)	0.0535 (9)
H3A	1.0685	0.0985	0.4351	0.064*
C4	0.9844 (5)	0.2134 (2)	0.3375 (6)	0.0582 (10)
H4A	1.0959	0.2229	0.3244	0.070*
C5	0.6391 (4)	0.4281 (2)	0.3598 (4)	0.0416 (8)
C6	0.7790 (6)	0.5037 (3)	0.5944 (6)	0.0627 (11)
H6A	0.8189	0.5105	0.7076	0.075*
C7	0.8125 (6)	0.5678 (3)	0.5086 (7)	0.0677 (13)
H7A	0.8734	0.6167	0.5664	0.081*
C8	0.6737 (5)	0.4914 (2)	0.2717 (5)	0.0561 (9)
H8A	0.6355	0.4843	0.1588	0.067*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0286 (4)	0.0384 (4)	0.0661 (6)	-0.0005 (3)	0.0238 (4)	-0.0064 (4)
N1	0.0336 (13)	0.0392 (15)	0.090 (2)	-0.0016 (11)	0.0314 (14)	0.0025 (14)
N2	0.0566 (17)	0.0332 (14)	0.073 (2)	0.0047 (13)	0.0357 (16)	0.0039 (14)
N3	0.0531 (17)	0.0451 (17)	0.062 (2)	0.0018 (13)	0.0250 (15)	-0.0052 (14)
N4	0.069 (2)	0.0405 (17)	0.110 (3)	-0.0067 (16)	0.050 (2)	-0.0055 (19)
O1	0.0378 (12)	0.0647 (17)	0.0632 (17)	0.0072 (11)	0.0165 (11)	-0.0084 (13)
O2	0.0493 (14)	0.0492 (15)	0.109 (2)	-0.0070 (12)	0.0529 (15)	-0.0115 (14)
O3	0.117 (3)	0.0558 (19)	0.159 (4)	0.020 (2)	0.092 (3)	0.033 (2)
C1	0.0313 (14)	0.0330 (15)	0.0563 (19)	-0.0009 (12)	0.0203 (13)	-0.0058 (14)
C2	0.0477 (18)	0.0394 (17)	0.074 (2)	0.0008 (14)	0.0395 (18)	-0.0013 (16)

C3	0.0350 (16)	0.0456 (19)	0.072 (2)	0.0044 (14)	0.0187 (16)	-0.0014 (18)
C4	0.0315 (16)	0.0463 (19)	0.099 (3)	0.0001 (14)	0.0318 (18)	0.0007 (19)
C5	0.0311 (14)	0.0333 (15)	0.062 (2)	0.0036 (12)	0.0230 (14)	-0.0017 (14)
C6	0.055 (2)	0.056 (2)	0.068 (3)	0.0036 (18)	0.0203 (19)	-0.017 (2)
C7	0.047 (2)	0.041 (2)	0.115 (4)	-0.0084 (16)	0.038 (2)	-0.024 (2)
C8	0.055 (2)	0.045 (2)	0.072 (3)	0.0007 (16)	0.0323 (19)	-0.0001 (18)

Geometric parameters (Å, °)

S1—O1	1.425 (3)	N4—C8	1.351 (5)
S1—O2	1.426 (3)	C1—C2	1.366 (5)
S1—C5	1.784 (3)	C2—H2A	0.9300
S1—C1	1.790 (3)	C3—C4	1.362 (6)
N1—C1	1.328 (4)	C3—H3A	0.9300
N1—C4	1.343 (4)	C4—H4A	0.9300
N2—O3	1.281 (4)	C5—C8	1.382 (5)
N2—C3	1.339 (5)	C6—C7	1.374 (7)
N2—C2	1.368 (4)	C6—H6A	0.9300
N3—C5	1.312 (5)	C7—H7A	0.9300
N3—C6	1.333 (5)	C8—H8A	0.9300
N4—C7	1.306 (6)		
O1—S1—O2	119.66 (17)	N2—C3—C4	120.2 (3)
O1—S1—C5	106.68 (17)	N2—C3—H3A	119.9
O2—S1—C5	109.23 (16)	C4—C3—H3A	119.9
O1—S1—C1	108.72 (16)	N1—C4—C3	123.6 (3)
O2—S1—C1	107.53 (17)	N1—C4—H4A	118.2
C5—S1—C1	103.92 (14)	C3—C4—H4A	118.2
C1—N1—C4	114.3 (3)	N3—C5—C8	124.2 (3)
O3—N2—C3	121.5 (3)	N3—C5—S1	116.4 (3)
O3—N2—C2	120.0 (3)	C8—C5—S1	119.4 (3)
C3—N2—C2	118.5 (3)	N3—C6—C7	121.7 (4)
C5—N3—C6	114.9 (4)	N3—C6—H6A	119.1
C7—N4—C8	115.9 (4)	C7—C6—H6A	119.1
N1—C1—C2	125.6 (3)	N4—C7—C6	123.4 (4)
N1—C1—S1	115.7 (2)	N4—C7—H7A	118.3
C2—C1—S1	118.6 (2)	C6—C7—H7A	118.3
C1—C2—N2	117.7 (3)	N4—C8—C5	119.8 (4)
C1—C2—H2A	121.1	N4—C8—H8A	120.1
N2—C2—H2A	121.1	C5—C8—H8A	120.1
C4—N1—C1—C2	-1.3 (5)	N2—C3—C4—N1	1.6 (7)
C4—N1—C1—S1	-179.2 (3)	C6—N3—C5—C8	1.0 (5)
O1—S1—C1—N1	59.3 (3)	C6—N3—C5—S1	-176.7 (3)
O2—S1—C1—N1	-169.8 (3)	O1—S1—C5—N3	169.1 (2)
C5—S1—C1—N1	-54.1 (3)	O2—S1—C5—N3	38.4 (3)
O1—S1—C1—C2	-118.8 (3)	C1—S1—C5—N3	-76.1 (3)
O2—S1—C1—C2	12.1 (3)	O1—S1—C5—C8	-8.7 (3)

C5—S1—C1—C2	127.8 (3)	O2—S1—C5—C8	-139.4 (3)
N1—C1—C2—N2	0.1 (6)	C1—S1—C5—C8	106.1 (3)
S1—C1—C2—N2	178.0 (3)	C5—N3—C6—C7	-0.1 (5)
O3—N2—C2—C1	-177.9 (4)	C8—N4—C7—C6	0.5 (6)
C3—N2—C2—C1	1.9 (5)	N3—C6—C7—N4	-0.6 (6)
O3—N2—C3—C4	177.1 (4)	C7—N4—C8—C5	0.3 (6)
C2—N2—C3—C4	-2.8 (6)	N3—C5—C8—N4	-1.1 (5)
C1—N1—C4—C3	0.4 (6)	S1—C5—C8—N4	176.5 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C2—H2 <i>A</i> \cdots O1 ⁱ	0.93	2.32	3.130 (5)	146
C3—H3 <i>A</i> \cdots O3 ⁱⁱ	0.93	2.56	3.419 (4)	153
C7—H7 <i>A</i> \cdots N1 ⁱⁱⁱ	0.93	2.57	3.449 (3)	157

Symmetry codes: (i) $x, -y+1/2, z+1/2$; (ii) $-x+2, -y, -z+1$; (iii) $-x+2, -y+1, -z+1$.