

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 3-(4-Chlorobenzoyl)-6-(4-chlorophenyl)-2,4-dimethylbenzonitrile

 Xin Wang,<sup>a,b</sup> Yi-Min Zhang<sup>a\*</sup> and Xue-Fei Jia<sup>a</sup>
<sup>a</sup>School of Chemistry and Environmental Science, Henan Normal University, Xixiang, Henan 453007, People's Republic of China, and <sup>b</sup>Xinyang Agricultural College, Xinyang, Henan 464000, People's Republic of China

Correspondence e-mail: hxzhyzm@sina.com


Received 19 August 2012; accepted 21 August 2012

 Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.047;  $wR$  factor = 0.121; data-to-parameter ratio = 14.5.

In the title compound,  $\text{C}_{22}\text{H}_{15}\text{Cl}_2\text{NO}$ , the terminal chlorobenzene rings are oriented at 44.51 (15) and 86.06 (17)° with respect to the central polysubstituted benzene ring, and make a dihedral angle of 49.48 (17)° with each other. In the crystal, molecules are linked by weak  $\text{C}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{N}$  interactions.

## Related literature

For background to the title compound, see: Ma (2003, 2005, 2007); Hoffmann-Röder *et al.* (2004). For related structures, see: Zhang *et al.* (2011); Fun *et al.* (2012); Jagadeesan *et al.* (2011).



## Experimental

## Crystal data

 $\text{C}_{22}\text{H}_{15}\text{Cl}_2\text{NO}$   
 $M_r = 380.25$   
 Monoclinic,  $P2_1/c$   
 $a = 7.8102$  (12) Å  
 $b = 30.032$  (5) Å

 $c = 8.2054$  (13) Å  
 $\beta = 103.739$  (2)°  
 $V = 1869.6$  (5) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation

 $\mu = 0.36$  mm<sup>-1</sup>  
 $T = 296$  K

 $0.39 \times 0.33 \times 0.23$  mm

## Data collection

 Bruker SMART CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2007)  
 $T_{\min} = 0.873$ ,  $T_{\max} = 0.922$ 

 11684 measured reflections  
 3442 independent reflections  
 2013 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.034$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.047$   
 $wR(F^2) = 0.121$   
 $S = 1.08$   
 3442 reflections

 237 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.21$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.24$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C3}-\text{H3}\cdots\text{N1}^i$	0.93	2.61	3.305 (5)	132
$\text{C5}-\text{H5}\cdots\text{O1}^{ii}$	0.93	2.53	3.390 (5)	155

 Symmetry codes: (i)  $x - 1, y, z$ ; (ii)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

This work was supported by the National Natural Science Foundation of China (No. 21172057).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5613).

## References

- Bruker (2007). SADABS, SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fun, H.-K., Loh, W.-S., Samshuddin, S., Narayana, B. & Sarojini, B. K. (2012). *Acta Cryst.* **E68**, o1877–o1878.
- Hoffmann-Röder, A. & Krause, N. (2004). *Angew. Chem. Int. Ed.* **43**, 1196–1216.
- Jagadeesan, G., Sethusankar, K., Sivasakthikumar, R. & Mohanakrishnan, A. K. (2011). *Acta Cryst.* **E67**, o2177.
- Ma, S. (2003). *Acc. Chem. Res.* **36**, 701–712.
- Ma, S. (2005). *Chem. Rev.* **105**, 2829–2871.
- Ma, S. (2007). *Aldrichim. Acta*, **40**, 91–102.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Zhang, X., Jia, X., Fang, L., Liu, N., Wang, J. & Fan, X. (2011). *Org. Lett.* **13**, 5024–5027.

## supporting information

*Acta Cryst.* (2012). E68, o2787 [doi:10.1107/S1600536812036409]

## 3-(4-Chlorobenzoyl)-6-(4-chlorophenyl)-2,4-dimethylbenzointrile

Xin Wang, Yi-Min Zhang and Xue-Fei Jia

### S1. Comment

It has been well documented that benzenoid compounds are ubiquitous structural units in a wide variety of naturally occurring compounds and a plethora of pharmaceuticals. On the other hand, allene derivatives are powerful synthetic intermediates toward a plethora of important organic compounds and frequent building blocks of natural products (Ma, 2003, 2005, 2007; Hoffmann-Röder *et al.*, 2004). In this regard, Zhang *et al.* have developed a novel and efficient method for the preparation of polysubstituted benzenes by one-pot double Michael addition/*intramolecular* aldol reaction/ decarboxylation of 1,2-allenic ketones with cyanoacetate (Zhang *et al.*, 2011). Herein, we would like to report the structure of one of the products obtained by this method.

In the title compound (Fig. 1), all the bond lengths and bond angles are within normal ranges. The central polysubstituted benzene ring (C7—C12) forms dihedral angles of 44.51 (15)° and 86.06 (17)° with the other two chloro-substituted phenyl rings (C1—C6) and (C17—C22), respectively. And the dihedral angle between the two chloro-substituted phenyl rings, (C1—C7) and (C7—C12), is 49.48 (17)°. The mean plane of the ketone group is almost coplanar with the neighboring chloro-substituted phenyl ring (C17-C22).


In the crystal structure, the molecules are connected *via* C—H···O and C—H···N interactions.

### S2. Experimental


A mixture of 1-(4-chlorophenyl)buta-2,3-dien-1-one (1 mmol), cyanoacetate (0.5 mmol) and K<sub>2</sub>CO<sub>3</sub> (0.5 mmol) in acetone (5 ml) was refluxed for 15 min. Upon completion, the reaction mixture was cooled to room temperature, added with water (10 ml) and extracted with ethyl acetate. The combined organic phases were washed with brine, dried, filtered and concentrated under vacuum. The residue was purified by column chromatography on silica gel eluting with ethyl acetate/hexane (1:20 *v/v*) to give the title compound as Colorless solids with a yield of 80%. Single crystals, suitable for X-ray diffraction analysis, were obtained by slow evaporation of solvent from a petroleum ether-dichloromethane (2:1 *v/v*) solution.

### S3. Refinement

The H atoms were included at calculated positions and were refined as riding atoms: C—H = 0.93 and 0.96 Å for aromatic and methyl H atoms, respectively, with  $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$ , where  $x = 1.2$  for aromatic H, and  $x = 1.5$  for methyl H atoms.

**Figure 1**

Molecular structure of the title compound, with displacement ellipsoids drawn at the 30% probability level.



**Figure 2**

Crystal structure of the title compound, viewed along the *a* axis. Intermolecular C—H···O and C—H···N hydrogen bonds are shown as dashed lines, only H atoms involved in hydrogen bonds are shown.

### 3-(4-Chlorobenzoyl)-6-(4-chlorophenyl)-2,4-dimethylbenzonitrile

#### Crystal data

$C_{22}H_{15}Cl_2NO$

$M_r = 380.25$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2_1/c$

$a = 7.8102\ (12)\ \text{\AA}$

$b = 30.032\ (5)\ \text{\AA}$

$c = 8.2054\ (13)\ \text{\AA}$

$\beta = 103.739\ (2)^\circ$

$V = 1869.6\ (5)\ \text{\AA}^3$

$Z = 4$

$F(000) = 784$

$D_x = 1.351\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 3123 reflections

$\theta = 2.6\text{--}22.9^\circ$

$\mu = 0.36\ \text{mm}^{-1}$

$T = 296$  K  $0.39 \times 0.33 \times 0.23$  mm  
 Block, colourless

*Data collection*

Bruker SMART CCD area-detector diffractometer	11684 measured reflections 3442 independent reflections
Radiation source: fine-focus sealed tube	2013 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\text{int}} = 0.034$
phi and $\omega$ scans	$\theta_{\text{max}} = 25.5^\circ$ , $\theta_{\text{min}} = 2.6^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2007)	$h = -9 \rightarrow 9$ $k = -36 \rightarrow 36$ $l = -9 \rightarrow 9$
$T_{\text{min}} = 0.873$ , $T_{\text{max}} = 0.922$	

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.047$	H-atom parameters constrained
$wR(F^2) = 0.121$	$w = 1/[\sigma^2(F_o^2) + (0.0235P)^2 + 1.9216P]$
$S = 1.08$	where $P = (F_o^2 + 2F_c^2)/3$
3442 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
237 parameters	$\Delta\rho_{\text{max}} = 0.21 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.24 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7995 (4)	0.21705 (10)	0.1502 (4)	0.0402 (7)
C2	0.6497 (4)	0.20687 (10)	0.2063 (4)	0.0452 (8)
H2	0.5945	0.2292	0.2535	0.054*
C3	0.5808 (4)	0.16426 (11)	0.1934 (4)	0.0525 (9)
H3	0.4806	0.1580	0.2318	0.063*
C4	0.6621 (5)	0.13136 (10)	0.1233 (4)	0.0518 (9)
C5	0.8084 (5)	0.14040 (11)	0.0632 (4)	0.0529 (9)
H5	0.8606	0.1181	0.0131	0.063*
C6	0.8767 (4)	0.18288 (10)	0.0779 (4)	0.0483 (8)
H6	0.9767	0.1889	0.0386	0.058*
C7	0.8714 (4)	0.26288 (10)	0.1663 (4)	0.0398 (7)
C8	1.0517 (4)	0.27148 (10)	0.2239 (4)	0.0442 (8)

C9	1.1199 (4)	0.31495 (11)	0.2343 (4)	0.0483 (8)
C10	1.0020 (4)	0.35035 (10)	0.1930 (4)	0.0460 (8)
C11	0.8210 (4)	0.34288 (10)	0.1389 (4)	0.0467 (8)
C12	0.7596 (4)	0.29920 (10)	0.1238 (4)	0.0451 (8)
H12	0.6394	0.2942	0.0838	0.054*
C13	1.1720 (5)	0.23601 (12)	0.2910 (5)	0.0558 (9)
C14	1.3152 (5)	0.32286 (14)	0.2910 (5)	0.0772 (12)
H14A	1.3698	0.3164	0.2005	0.116*
H14B	1.3639	0.3038	0.3845	0.116*
H14C	1.3365	0.3534	0.3241	0.116*
C15	0.6925 (5)	0.38100 (11)	0.0981 (5)	0.0646 (10)
H15A	0.5752	0.3695	0.0592	0.097*
H15B	0.7215	0.3991	0.0122	0.097*
H15C	0.6988	0.3986	0.1969	0.097*
C16	1.0722 (5)	0.39731 (11)	0.2256 (5)	0.0547 (9)
C17	1.1277 (4)	0.42252 (11)	0.0920 (4)	0.0495 (8)
C18	1.2013 (5)	0.46432 (11)	0.1290 (5)	0.0634 (10)
H18	1.2167	0.4758	0.2367	0.076*
C19	1.2517 (5)	0.48886 (12)	0.0070 (5)	0.0693 (11)
H19	1.3029	0.5167	0.0326	0.083*
C20	1.2263 (5)	0.47210 (11)	-0.1532 (5)	0.0589 (10)
C21	1.1562 (5)	0.43075 (11)	-0.1909 (5)	0.0658 (11)
H21	1.1412	0.4194	-0.2988	0.079*
C22	1.1076 (5)	0.40580 (11)	-0.0673 (5)	0.0619 (10)
H22	1.0608	0.3774	-0.0924	0.074*
Cl1	0.57545 (16)	0.07788 (3)	0.10760 (15)	0.0838 (4)
Cl2	1.28408 (16)	0.50406 (3)	-0.30697 (14)	0.0848 (4)
N1	1.2698 (4)	0.20909 (12)	0.3525 (5)	0.0805 (11)
O1	1.0819 (4)	0.41318 (9)	0.3638 (3)	0.0857 (9)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0442 (18)	0.0436 (17)	0.0329 (18)	0.0008 (14)	0.0092 (14)	0.0012 (14)
C2	0.049 (2)	0.0429 (18)	0.045 (2)	0.0041 (15)	0.0140 (16)	-0.0021 (15)
C3	0.055 (2)	0.051 (2)	0.054 (2)	-0.0044 (17)	0.0172 (17)	0.0015 (17)
C4	0.067 (2)	0.0420 (18)	0.045 (2)	-0.0002 (17)	0.0117 (18)	0.0021 (16)
C5	0.069 (2)	0.050 (2)	0.041 (2)	0.0109 (18)	0.0164 (18)	-0.0040 (16)
C6	0.057 (2)	0.0477 (19)	0.044 (2)	0.0038 (17)	0.0204 (17)	0.0003 (16)
C7	0.0453 (19)	0.0439 (18)	0.0323 (18)	-0.0010 (15)	0.0130 (15)	-0.0001 (14)
C8	0.045 (2)	0.0508 (19)	0.040 (2)	0.0009 (16)	0.0165 (15)	-0.0008 (15)
C9	0.046 (2)	0.061 (2)	0.039 (2)	-0.0086 (17)	0.0138 (16)	-0.0054 (16)
C10	0.057 (2)	0.0502 (19)	0.0314 (19)	-0.0099 (17)	0.0125 (16)	-0.0035 (14)
C11	0.058 (2)	0.0470 (19)	0.0354 (19)	-0.0021 (16)	0.0117 (16)	0.0022 (15)
C12	0.0446 (19)	0.0485 (19)	0.042 (2)	-0.0017 (16)	0.0095 (15)	0.0012 (15)
C13	0.050 (2)	0.062 (2)	0.059 (2)	0.0008 (19)	0.0207 (19)	0.0016 (19)
C14	0.054 (2)	0.085 (3)	0.090 (3)	-0.016 (2)	0.013 (2)	-0.010 (2)
C15	0.070 (3)	0.044 (2)	0.076 (3)	0.0014 (18)	0.011 (2)	0.0032 (18)

C16	0.063 (2)	0.055 (2)	0.048 (2)	-0.0154 (18)	0.0178 (18)	-0.0095 (18)
C17	0.059 (2)	0.0467 (18)	0.044 (2)	-0.0116 (17)	0.0144 (17)	-0.0070 (16)
C18	0.083 (3)	0.053 (2)	0.057 (2)	-0.023 (2)	0.021 (2)	-0.0158 (18)
C19	0.092 (3)	0.043 (2)	0.077 (3)	-0.022 (2)	0.030 (2)	-0.010 (2)
C20	0.070 (3)	0.045 (2)	0.066 (3)	-0.0041 (18)	0.026 (2)	0.0057 (18)
C21	0.094 (3)	0.053 (2)	0.056 (2)	-0.020 (2)	0.028 (2)	-0.0097 (18)
C22	0.082 (3)	0.049 (2)	0.058 (3)	-0.0240 (19)	0.022 (2)	-0.0079 (18)
C11	0.1082 (9)	0.0465 (5)	0.1010 (9)	-0.0146 (5)	0.0337 (7)	-0.0085 (5)
C12	0.1219 (10)	0.0556 (6)	0.0896 (8)	-0.0063 (6)	0.0505 (7)	0.0130 (5)
N1	0.065 (2)	0.083 (2)	0.095 (3)	0.018 (2)	0.023 (2)	0.014 (2)
O1	0.127 (3)	0.0814 (19)	0.0601 (19)	-0.0443 (17)	0.0442 (18)	-0.0276 (15)

*Geometric parameters (Å, °)*

C1—C2	1.389 (4)	C12—H12	0.9300
C1—C6	1.393 (4)	C13—N1	1.144 (4)
C1—C7	1.481 (4)	C14—H14A	0.9600
C2—C3	1.382 (4)	C14—H14B	0.9600
C2—H2	0.9300	C14—H14C	0.9600
C3—C4	1.373 (4)	C15—H15A	0.9600
C3—H3	0.9300	C15—H15B	0.9600
C4—C5	1.374 (5)	C15—H15C	0.9600
C4—C11	1.736 (3)	C16—O1	1.216 (4)
C5—C6	1.377 (4)	C16—C17	1.479 (4)
C5—H5	0.9300	C17—C22	1.374 (4)
C6—H6	0.9300	C17—C18	1.384 (4)
C7—C12	1.389 (4)	C18—C19	1.373 (5)
C7—C8	1.398 (4)	C18—H18	0.9300
C8—C9	1.405 (4)	C19—C20	1.377 (5)
C8—C13	1.440 (5)	C19—H19	0.9300
C9—C10	1.395 (4)	C20—C21	1.363 (5)
C9—C14	1.504 (5)	C20—C12	1.729 (3)
C10—C11	1.396 (4)	C21—C22	1.384 (5)
C10—C16	1.514 (4)	C21—H21	0.9300
C11—C12	1.392 (4)	C22—H22	0.9300
C11—C15	1.507 (4)		
C2—C1—C6	117.6 (3)	C11—C12—H12	118.8
C2—C1—C7	120.4 (3)	N1—C13—C8	176.2 (4)
C6—C1—C7	122.0 (3)	C9—C14—H14A	109.5
C3—C2—C1	121.4 (3)	C9—C14—H14B	109.5
C3—C2—H2	119.3	H14A—C14—H14B	109.5
C1—C2—H2	119.3	C9—C14—H14C	109.5
C4—C3—C2	119.2 (3)	H14A—C14—H14C	109.5
C4—C3—H3	120.4	H14B—C14—H14C	109.5
C2—C3—H3	120.4	C11—C15—H15A	109.5
C3—C4—C5	121.0 (3)	C11—C15—H15B	109.5
C3—C4—C11	119.1 (3)	H15A—C15—H15B	109.5

C5—C4—C11	119.9 (3)	C11—C15—H15C	109.5
C4—C5—C6	119.3 (3)	H15A—C15—H15C	109.5
C4—C5—H5	120.3	H15B—C15—H15C	109.5
C6—C5—H5	120.3	O1—C16—C17	121.8 (3)
C5—C6—C1	121.5 (3)	O1—C16—C10	118.0 (3)
C5—C6—H6	119.3	C17—C16—C10	120.2 (3)
C1—C6—H6	119.3	C22—C17—C18	119.2 (3)
C12—C7—C8	117.5 (3)	C22—C17—C16	122.0 (3)
C12—C7—C1	120.3 (3)	C18—C17—C16	118.8 (3)
C8—C7—C1	122.2 (3)	C19—C18—C17	120.2 (3)
C7—C8—C9	121.9 (3)	C19—C18—H18	119.9
C7—C8—C13	120.4 (3)	C17—C18—H18	119.9
C9—C8—C13	117.4 (3)	C18—C19—C20	119.9 (3)
C10—C9—C8	118.4 (3)	C18—C19—H19	120.1
C10—C9—C14	121.1 (3)	C20—C19—H19	120.1
C8—C9—C14	120.5 (3)	C21—C20—C19	120.6 (3)
C9—C10—C11	121.0 (3)	C21—C20—C12	120.0 (3)
C9—C10—C16	118.5 (3)	C19—C20—C12	119.4 (3)
C11—C10—C16	120.2 (3)	C20—C21—C22	119.5 (3)
C12—C11—C10	118.8 (3)	C20—C21—H21	120.3
C12—C11—C15	119.9 (3)	C22—C21—H21	120.3
C10—C11—C15	121.3 (3)	C17—C22—C21	120.6 (3)
C7—C12—C11	122.3 (3)	C17—C22—H22	119.7
C7—C12—H12	118.8	C21—C22—H22	119.7
C6—C1—C2—C3	1.0 (5)	C16—C10—C11—C12	-174.6 (3)
C7—C1—C2—C3	-179.6 (3)	C9—C10—C11—C15	178.5 (3)
C1—C2—C3—C4	-0.2 (5)	C16—C10—C11—C15	5.1 (5)
C2—C3—C4—C5	-1.2 (5)	C8—C7—C12—C11	-0.9 (4)
C2—C3—C4—C11	179.9 (3)	C1—C7—C12—C11	179.2 (3)
C3—C4—C5—C6	1.8 (5)	C10—C11—C12—C7	2.4 (5)
C11—C4—C5—C6	-179.4 (3)	C15—C11—C12—C7	-177.3 (3)
C4—C5—C6—C1	-0.9 (5)	C7—C8—C13—N1	-119 (6)
C2—C1—C6—C5	-0.4 (5)	C9—C8—C13—N1	55 (6)
C7—C1—C6—C5	-179.9 (3)	C9—C10—C16—O1	-86.8 (4)
C2—C1—C7—C12	-44.8 (4)	C11—C10—C16—O1	86.8 (4)
C6—C1—C7—C12	134.6 (3)	C9—C10—C16—C17	92.8 (4)
C2—C1—C7—C8	135.3 (3)	C11—C10—C16—C17	-93.6 (4)
C6—C1—C7—C8	-45.3 (4)	O1—C16—C17—C22	-176.1 (4)
C12—C7—C8—C9	-1.9 (4)	C10—C16—C17—C22	4.3 (5)
C1—C7—C8—C9	178.1 (3)	O1—C16—C17—C18	3.5 (6)
C12—C7—C8—C13	171.8 (3)	C10—C16—C17—C18	-176.1 (3)
C1—C7—C8—C13	-8.3 (4)	C22—C17—C18—C19	0.7 (6)
C7—C8—C9—C10	3.0 (5)	C16—C17—C18—C19	-179.0 (4)
C13—C8—C9—C10	-170.8 (3)	C17—C18—C19—C20	1.0 (6)
C7—C8—C9—C14	-177.8 (3)	C18—C19—C20—C21	-1.9 (6)
C13—C8—C9—C14	8.4 (5)	C18—C19—C20—C12	178.0 (3)
C8—C9—C10—C11	-1.4 (4)	C19—C20—C21—C22	1.1 (6)



C14—C9—C10—C11	179.4 (3)	C12—C20—C21—C22	-178.9 (3)
C8—C9—C10—C16	172.1 (3)	C18—C17—C22—C21	-1.5 (6)
C14—C9—C10—C16	-7.1 (5)	C16—C17—C22—C21	178.1 (4)
C9—C10—C11—C12	-1.2 (5)	C20—C21—C22—C17	0.6 (6)

*Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C3—H3...N1 <sup>i</sup>	0.93	2.61	3.305 (5)	132
C5—H5...O1 <sup>ii</sup>	0.93	2.53	3.390 (5)	155

Symmetry codes: (i)  $x-1, y, z$ ; (ii)  $x, -y+1/2, z-1/2$ .