

1,1,2,2-Tetrakis[2,4-dichloro-6-(diethoxymethyl)phenoxyethyl]ethene

Yavuz Köysal,^a Sema Öztürk Yildirim,^{b,c} Ray J. Butcher^{b*} and Esra Düğdü^d

^aYeşilyurt Demir Çelik Vocational School, Ondokuz Mayıs University, Samsun, Turkey, ^bDepartment of Chemistry, Howard University, 525 College Street NW, Washington, DC 20059, USA, ^cDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, and ^dDepartment of Chemistry, Karadeniz Technical University, 61080 Trabzon, Turkey
Correspondence e-mail: rbutcher99@yahoo.com


Received 4 September 2012; accepted 6 September 2012

Key indicators: single-crystal X-ray study; $T = 123$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; disorder in main residue; R factor = 0.038; wR factor = 0.090; data-to-parameter ratio = 18.2.

In the title compound, $\text{C}_{50}\text{H}_{60}\text{Cl}_8\text{O}_{12}$, the molecules are disordered about an inversion center located at the mid-point of the central $\text{C}=\text{C}$ bond. These atoms show disorder and were modelled with two different orientations with site occupancies of 0.828 (3) and 0.172 (3). The dihedral angle between the two benzene rings in the asymmetric unit is $52.80(6)^\circ$. Intramolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{Cl}$ interactions occur and the crystal packing features inversion dimers linked by pairs of $\text{C}-\text{H}\cdots\text{O}$ bonds, generating $R_2^2(10)$ loops.

Related literature

For anti-oxidant, anti-inflammatory, chemopreventive, anti-bacterial, anticarcinogenic, antitumor and antiviral properties of sterically hindered phenols and secondary aromatic amines, see: Amorati *et al.* (2003); Torres de Pinedo *et al.* (2007); Leopoldini *et al.* (2011); Leiro *et al.* (2011); Link *et al.* (2010); Daglia (2011); Bai *et al.*, (2003); Song *et al.* (2005); Rabek (1990); Pospisil *et al.* (2003); Wolf & Kaul (1992); Thapa *et al.* (2012). For synthetic phenolic antioxidants, such as butylated hydroxytoluene (BHT), butylated hydroxyanisole (BHA) or butylated hydroxyquinone (TBHQ), which possess good anti-oxidant capacity, see: Omura (1995). For phenols capable of propagation termination due to the donation of the hydrogen atom of the phenolic OH to the free radicals, see: Kumar & Naik (2010); Findik *et al.* (2011). For bond lengths of structurally related molecules, see: Öztürk Yildirim *et al.* (2012). For a description of the Cambridge structural Database, see: Allen (2002). For details of the synthesis, see: Er *et al.* (2009).



Experimental

Crystal data

$\text{C}_{50}\text{H}_{60}\text{Cl}_8\text{O}_{12}$
 $M_r = 1136.58$
Triclinic, $P\bar{1}$
 $a = 8.0626(3)$ Å
 $b = 12.8693(5)$ Å
 $c = 13.9968(6)$ Å
 $\alpha = 97.425(3)^\circ$
 $\beta = 102.878(3)^\circ$

$\gamma = 105.391(3)^\circ$
 $V = 1337.29(9)$ Å³
 $Z = 1$
Mo $K\alpha$ radiation
 $\mu = 0.48$ mm⁻¹
 $T = 123$ K
 $0.62 \times 0.19 \times 0.07$ mm

Data collection

Agilent Xcalibur (Ruby, Gemini) diffractometer
Absorption correction: multi-scan [*CrysAlis RED* (Agilent, 2011), based on expressions derived

from Clark & Reid (1995)]
 $T_{\min} = 0.755$, $T_{\max} = 0.967$
10254 measured reflections
6125 independent reflections
5231 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.090$
 $S = 1.03$
6125 reflections
336 parameters

12 restraints
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.45$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.39$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}12-\text{H}12B\cdots\text{Cl}2$	0.99	2.66	3.180 (3)	113
$\text{C}14-\text{H}14B\cdots\text{O}4$	0.99	2.50	3.166 (3)	125
$\text{C}17-\text{H}17A\cdots\text{O}2^i$	0.95	2.47	3.3943 (18)	166

Symmetry code: (i) $-x + 2, -y + 1, -z + 1$.

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

RJB acknowledges the NSF-MRI program (grant No. CHE-0619278) for funds to purchase the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT6833).

References

- Agilent (2011). *CrysAlis PRO* and *CrysAlis RED*. Agilent Technologies, Oxfordshire, England.
- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Amorati, R., Lucarini, M., Mugnaini, V. & Pedulli, G. F. (2003). *J. Org. Chem.* **68**, 5198–5204.
- Bai, X., Cerimele, F., Fukai, M. U., Waqas, M., Campbell, P. M., Govindarajan, B., Der, C. J., Battle, T., Frank, D. A., Ye, K., Murad, E., Dubiel, W., Soff, G. & Arbiser, J. L. (2003). *J. Biol. Chem.* **278**, 35501–35507.
- Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.
- Daglia, M. (2011). *Curr. Opin. Biotechnol.* **23**, 1–8.
- Er, M., Ünver, Y., Sancak, K., Degirmencioglu, I. & Karaoglu, S. A. (2009). *Arkivoc*, **II**, 149–167.
- Findik, E., Ceylan, M. & Elmastas, M. (2011). *Eur. J. Med. Chem.* **46**, 4618–4624.
- Kumar, H. V. & Naik, N. (2010). *Eur. J. Med. Chem.* **45**, 2–10.
- Leiro, J. M., Varela, M., Piazzon, M. C., Arranz, J. A., Noya, M. & Lamas, J. (2011). *Mol. Immunol.* **47**, 1114–1120.
- Leopoldini, M., Russo, N. & Toscano, M. (2011). *Food Chem.* **125**, 288–306.
- Link, A., Balaguer, F. & Goel, A. (2010). *Biochem. Pharmacol.* **80**, 1771–1792.
- Omura, K. (1995). *J. Am. Oil Chem. Soc.* **72**, 1565–1570.
- Öztürk Yildirim, S., Butcher, R. J., Köysal, Y. & Birinci, E. (2012). *Acta Cryst.* **E68**, o2633.
- Pospisil, J., Horak, Z., Pilar, J., Billingham, N. C., Zweifel, H. & Nespurek, S. (2003). *Polym. Degrad. Stab.* **82**, 145–162.
- Rabek, J. F. (1990). In *Photostabilization of Polymers-Principles and Applications*. New York: Elsevier Applied Science.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Song, J. L., Lee, K. H. & Seong, B. L. (2005). *Antiviral Res.* **68**, 66–74.
- Thapa, P., Karki, R., Yun, M., Kadayat, T. M., Lee, E., Kwon, H. B., Na, Y., Cho, W.-J., Kim, N. D., Jeong, B.-S., Kwon, Y. & Lee, E.-S. (2012). *Eur. J. Med. Chem.* **52**, 123–136.
- Torres de Pinedo, A., Penalver, P. & Morales, J. C. (2007). *Food Chem.* **103**, 55–61.
- Wolf, R. & Kaul, B. L. (1992). Editors. *Plastics, Additives*, in *Ullmann's Encyclopedia of Industrial Chemistry*, Vol. A20, pp. 459–507. Weinheim: VCH Verlag.

supporting information

Acta Cryst. (2012). E68, o2993–o2994 [https://doi.org/10.1107/S1600536812038299]

1,1,2,2-Tetrakis[2,4-dichloro-6-(diethoxymethyl)phenoxy]ethene**Yavuz Köysal, Sema Öztürk Yildirim, Ray J. Butcher and Esra Düğdü****S1. Comment**

Many phenolic or polyphenolic compounds have been reported to have a wide range of biological activities such as antioxidant (Amorati *et al.*, 2003; Torres de Pinedo *et al.*, 2007; Leopoldini *et al.*, 2011), anti-inflammatory (Leiro *et al.*, 2011), chemoprevention (Link *et al.*, 2010), antibacterial (Daglia, 2011), anti-carcinogenic and anti-tumor (Bai *et al.*, 2003), and antiviral (Song *et al.*, 2005) activities. The most active antioxidants typically comprise sterically hindered phenols and secondary aromatic amines (Rabek, 1990; Pospisil *et al.*, 2003; Wolf & Kaul, 1992; Thapa *et al.*, 2012). In addition, phenols have been utilized extensively for food preservation. Synthetic phenolic antioxidants, such as butylated hydroxytoluene (BHT), butylated hydroxyanisole (BHA) or butylated hydroxyquinone (TBHQ) possess good antioxidant capacity (Omura, 1995). The main structural feature responsible for the anti-oxidative and free radical scavenging activity of phenolic derivatives is the phenolic hydroxyl group. Phenols are able of donating the hydrogen atom of the phenolic OH to the free radicals, thus stopping the propagation chain during the oxidation process (Kumar & Naik, 2010; Findik, *et al.*, 2011). In view of the importance of such phenolate compounds the structure of 2,2'-((2-(1,3-bis(2,4-dichloro-6-(diethoxymethyl)phenoxy)propan-2-ylidene) propane-1,3-diyl)bis(oxy))bis(1,5-dichloro-3-(diethoxymethyl)benzene) was determined.

The title compound, (Fig. 1), lies on an inversion centre, giving one half-molecule per asymmetric unit which passes through middle point of the C13=C13A double bond of the aliphatic chain. Atoms C12 C13 and C14 atoms show disorder and were modelled with two different orientations and with site occupancies of 0.828 (4):0.172 (4). The (diethoxymethyl)benzene groups adopts an all-*trans* conformation and the molecular structure is not planar. The O3 C12 C13 C14 and C12 C13 C14 O6 torsion angles are 68.6 (2) ° and 82.5 (2) ° and the dihedral angle between the planes of the benzene rings (C1/C6 to C15/C20) is 52.80 (6) ° [for the non-H atoms, maximum deviation = 0.007 (1) Å for C2]. Bond lengths and angles can be regarded as normal for such structures (Öztürk Yildirim *et al.* 2012; Allen, 2002). No classical hydrogen bonds are observed in the crystal structure.

S2. Experimental

Title compound was published methods (Er *et al.*, 2009). Crystals were grown by slow evaporation of an dimethyl-formamide/alcohol mixed solution.

S3. Refinement

The hydrogen atoms were placed in calculated positions with C—H = 0.95–0.99 Å and refined using a riding model with fixed isotropic displacement parameters [$U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for the methyl groups and $1.2U_{\text{eq}}(\text{C})$ for the other H atoms]. The molecules are disordered about an inversion center, therefore, the O1—C12/C12—C13 and C13—C14/C14—O1 distances are average values. The SIMU and DELU constraint instructions in *SHELXL97* were used atom C13 in order to model the disorder properly during the refinement. For C13B the ISOR instruction was used as otherwise it went non-

positive definite. The displacement parameters of the pairs C12/C12B and C14/C14B were set equal using the EADP instruction.




Figure 1

The molecular structure of (I) showing displacement ellipsoids at the 50% probability level [symmetry code: (A) = 1 - x, 1 - y, 1 - z].

1,1,2,2-Tetrakis[2,4-dichloro-6-(diethoxymethyl)phenoxy]ethane

Crystal data

$C_{50}H_{60}Cl_8O_{12}$

$M_r = 1136.58$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 8.0626$ (3) Å

$b = 12.8693$ (5) Å

$c = 13.9968$ (6) Å

$\alpha = 97.425$ (3)°

$\beta = 102.878$ (3)°

$\gamma = 105.391$ (3)°

$V = 1337.29$ (9) Å³

$Z = 1$

$F(000) = 592$

$D_x = 1.411$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4477 reflections

$\theta = 3.0$ – 29.4 °

$\mu = 0.48$ mm⁻¹

$T = 123$ K

Plate, colorless

$0.62 \times 0.19 \times 0.07$ mm

Data collection

Agilent Xcalibur (Ruby, Gemini)
diffractometer

Radiation source: Enhance (Cu) X-ray Source

Graphite monochromator

Detector resolution: 10.5081 pixels mm⁻¹

ω scans

Absorption correction: multi-scan
 [CrysAlis RED (Agilent, 2011), based on
 expressions derived from Clark & Reid (1995)]
 $T_{\min} = 0.755$, $T_{\max} = 0.967$
 10254 measured reflections
 6125 independent reflections

5231 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$
 $\theta_{\max} = 29.5^\circ$, $\theta_{\min} = 3.1^\circ$
 $h = -8 \rightarrow 11$
 $k = -17 \rightarrow 17$
 $l = -18 \rightarrow 17$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.090$
 $S = 1.03$
 6125 reflections
 336 parameters
 12 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0366P)^2 + 0.6175P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.45 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.39 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. Absorption correction: CrysAlis RED, (Agilent, 2011) Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm. (Clark & Reid, 1995).

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C11	-0.23088 (6)	0.18327 (3)	-0.05330 (3)	0.03455 (11)	
C12	0.27679 (5)	0.55748 (3)	0.14470 (3)	0.03249 (10)	
C13	1.09088 (5)	0.67178 (3)	0.63733 (3)	0.02509 (9)	
C14	1.49291 (5)	0.83171 (4)	0.39813 (3)	0.03397 (10)	
O1	0.06285 (14)	0.10425 (9)	0.28297 (8)	0.0261 (3)	
O2	0.36555 (14)	0.17140 (9)	0.28761 (8)	0.0224 (2)	
O3	0.38227 (14)	0.41369 (8)	0.29369 (8)	0.0246 (2)	
O4	0.69832 (14)	0.58377 (9)	0.23343 (8)	0.0247 (2)	
O5	0.83793 (15)	0.75433 (10)	0.20043 (8)	0.0284 (3)	
O6	0.78680 (12)	0.65010 (8)	0.46395 (8)	0.0199 (2)	
C1	0.23412 (18)	0.36585 (12)	0.21470 (11)	0.0181 (3)	
C2	0.17432 (19)	0.41902 (12)	0.13926 (11)	0.0207 (3)	
C3	0.03219 (19)	0.36340 (13)	0.05666 (12)	0.0219 (3)	
H3A	-0.0064	0.4001	0.0053	0.026*	
C4	-0.05207 (19)	0.25336 (13)	0.05072 (11)	0.0212 (3)	
C5	0.00081 (19)	0.19851 (12)	0.12452 (11)	0.0204 (3)	
H5A	-0.0606	0.1231	0.1193	0.024*	
C6	0.14491 (18)	0.25450 (12)	0.20676 (11)	0.0177 (3)	

C7	0.20623 (19)	0.19703 (12)	0.29041 (11)	0.0203 (3)	
H7A	0.2288	0.2474	0.3558	0.024*	
C8	0.0878 (2)	0.05009 (16)	0.36587 (14)	0.0360 (4)	
H8A	0.1526	0.1048	0.4286	0.043*	
H8B	0.1581	−0.0008	0.3557	0.043*	
C9	−0.0953 (2)	−0.01277 (15)	0.37163 (14)	0.0366 (4)	
H9A	−0.0838	−0.0623	0.4186	0.055*	
H9B	−0.1664	−0.0559	0.3052	0.055*	
H9C	−0.1545	0.0391	0.3948	0.055*	
C10	0.3566 (2)	0.10171 (17)	0.19734 (14)	0.0384 (4)	
H10A	0.3398	0.1403	0.1408	0.046*	
H10B	0.2538	0.0342	0.1828	0.046*	
C11	0.5243 (3)	0.07216 (18)	0.20881 (18)	0.0505 (6)	
H11A	0.5235	0.0302	0.1450	0.076*	
H11B	0.5340	0.0274	0.2601	0.076*	
H11C	0.6263	0.1394	0.2289	0.076*	
C12	0.3929 (3)	0.5092 (2)	0.36286 (19)	0.0173 (5)	0.828 (3)
H12A	0.2743	0.5051	0.3738	0.021*	0.828 (3)
H12B	0.4360	0.5768	0.3374	0.021*	0.828 (3)
C13	0.5229 (2)	0.50930 (13)	0.45844 (14)	0.0173 (4)	0.828 (3)
C14	0.7108 (3)	0.5318 (2)	0.44948 (19)	0.0156 (5)	0.828 (3)
H14A	0.7817	0.5029	0.5008	0.019*	0.828 (3)
H14B	0.7109	0.4957	0.3826	0.019*	0.828 (3)
C12B	0.3501 (18)	0.5066 (13)	0.3681 (11)	0.0173 (5)	0.172 (3)
H12C	0.4165	0.5802	0.3601	0.021*	0.172 (3)
H12D	0.2215	0.5003	0.3518	0.021*	0.172 (3)
C13B	0.5857 (9)	0.5064 (6)	0.5228 (6)	0.0069 (12)	0.172 (3)
C14B	0.7384 (19)	0.5280 (13)	0.4666 (12)	0.0156 (5)	0.172 (3)
H14C	0.8434	0.5096	0.5031	0.019*	0.172 (3)
H14D	0.6946	0.4831	0.3979	0.019*	0.172 (3)
C15	0.95214 (18)	0.68454 (11)	0.44614 (11)	0.0173 (3)	
C16	1.10747 (19)	0.70221 (12)	0.52180 (11)	0.0184 (3)	
C17	1.27632 (19)	0.74633 (12)	0.50857 (12)	0.0212 (3)	
H17A	1.3817	0.7582	0.5607	0.025*	
C18	1.28421 (19)	0.77210 (12)	0.41639 (12)	0.0217 (3)	
C19	1.13317 (19)	0.75490 (13)	0.33920 (12)	0.0223 (3)	
H19A	1.1437	0.7729	0.2767	0.027*	
C20	0.96562 (19)	0.71107 (12)	0.35371 (11)	0.0193 (3)	
C21	0.7944 (2)	0.69472 (13)	0.27337 (12)	0.0225 (3)	
H21A	0.7165	0.7284	0.3052	0.027*	
C22	0.7855 (2)	0.52048 (15)	0.18194 (14)	0.0321 (4)	
H22A	0.8899	0.5121	0.2295	0.039*	
H22B	0.8272	0.5574	0.1300	0.039*	
C23	0.6502 (3)	0.40991 (17)	0.13475 (16)	0.0435 (5)	
H23A	0.7047	0.3638	0.0986	0.065*	
H23B	0.5478	0.4195	0.0880	0.065*	
H23C	0.6101	0.3744	0.1870	0.065*	
C24	0.6835 (2)	0.76496 (16)	0.13328 (14)	0.0350 (4)	

H24A	0.6044	0.7868	0.1714	0.042*
H24B	0.6157	0.6938	0.0871	0.042*
C25	0.7450 (3)	0.85117 (19)	0.07503 (15)	0.0456 (5)
H25A	0.6409	0.8612	0.0300	0.068*
H25B	0.8198	0.8276	0.0357	0.068*
H25C	0.8145	0.9209	0.1214	0.068*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0297 (2)	0.0299 (2)	0.0292 (2)	0.00642 (17)	-0.01299 (17)	-0.00331 (17)
C12	0.02656 (19)	0.02333 (18)	0.0371 (2)	-0.00010 (16)	-0.00738 (17)	0.01357 (17)
C13	0.02481 (17)	0.03109 (19)	0.02040 (18)	0.00847 (15)	0.00794 (14)	0.00553 (15)
C14	0.01604 (16)	0.0430 (2)	0.0429 (2)	0.00276 (16)	0.01405 (16)	0.01116 (19)
O1	0.0259 (5)	0.0241 (5)	0.0267 (6)	0.0042 (5)	0.0045 (5)	0.0113 (5)
O2	0.0229 (5)	0.0233 (5)	0.0196 (5)	0.0105 (4)	0.0005 (4)	0.0010 (4)
O3	0.0228 (5)	0.0194 (5)	0.0229 (6)	0.0075 (4)	-0.0080 (4)	-0.0036 (4)
O4	0.0190 (5)	0.0270 (5)	0.0270 (6)	0.0078 (4)	0.0050 (4)	0.0022 (5)
O5	0.0246 (5)	0.0366 (6)	0.0276 (6)	0.0123 (5)	0.0057 (5)	0.0150 (5)
O6	0.0138 (4)	0.0155 (5)	0.0317 (6)	0.0032 (4)	0.0103 (4)	0.0041 (4)
C1	0.0154 (6)	0.0211 (7)	0.0152 (7)	0.0064 (6)	-0.0001 (5)	0.0000 (6)
C2	0.0178 (7)	0.0185 (7)	0.0231 (8)	0.0041 (6)	0.0016 (6)	0.0043 (6)
C3	0.0193 (7)	0.0258 (7)	0.0207 (7)	0.0094 (6)	0.0016 (6)	0.0064 (6)
C4	0.0170 (7)	0.0249 (7)	0.0169 (7)	0.0075 (6)	-0.0024 (6)	-0.0030 (6)
C5	0.0194 (7)	0.0183 (7)	0.0214 (7)	0.0058 (6)	0.0027 (6)	0.0012 (6)
C6	0.0173 (6)	0.0204 (7)	0.0170 (7)	0.0084 (6)	0.0053 (5)	0.0026 (6)
C7	0.0210 (7)	0.0197 (7)	0.0194 (7)	0.0064 (6)	0.0040 (6)	0.0030 (6)
C8	0.0341 (9)	0.0397 (9)	0.0376 (10)	0.0102 (8)	0.0088 (8)	0.0233 (8)
C9	0.0396 (10)	0.0340 (9)	0.0291 (9)	-0.0006 (8)	0.0074 (8)	0.0096 (8)
C10	0.0356 (9)	0.0479 (11)	0.0287 (9)	0.0231 (8)	-0.0003 (8)	-0.0092 (8)
C11	0.0298 (9)	0.0471 (12)	0.0633 (14)	0.0164 (9)	0.0021 (9)	-0.0197 (11)
C12	0.0173 (11)	0.0176 (7)	0.0158 (8)	0.0066 (8)	0.0015 (8)	0.0017 (6)
C13	0.0155 (8)	0.0130 (8)	0.0214 (10)	0.0031 (6)	0.0036 (7)	0.0016 (7)
C14	0.0124 (9)	0.0145 (7)	0.0211 (11)	0.0048 (7)	0.0049 (7)	0.0048 (7)
C12B	0.0173 (11)	0.0176 (7)	0.0158 (8)	0.0066 (8)	0.0015 (8)	0.0017 (6)
C13B	0.0069 (14)	0.0069 (14)	0.0065 (15)	0.0017 (9)	0.0020 (9)	0.0011 (9)
C14B	0.0124 (9)	0.0145 (7)	0.0211 (11)	0.0048 (7)	0.0049 (7)	0.0048 (7)
C15	0.0140 (6)	0.0137 (6)	0.0255 (7)	0.0041 (5)	0.0082 (6)	0.0033 (6)
C16	0.0190 (6)	0.0173 (7)	0.0199 (7)	0.0055 (5)	0.0077 (6)	0.0031 (6)
C17	0.0154 (6)	0.0215 (7)	0.0255 (8)	0.0061 (6)	0.0042 (6)	0.0017 (6)
C18	0.0137 (6)	0.0210 (7)	0.0314 (8)	0.0031 (6)	0.0107 (6)	0.0055 (6)
C19	0.0208 (7)	0.0258 (7)	0.0240 (8)	0.0080 (6)	0.0097 (6)	0.0095 (6)
C20	0.0170 (6)	0.0182 (7)	0.0234 (7)	0.0064 (6)	0.0053 (6)	0.0042 (6)
C21	0.0181 (7)	0.0258 (7)	0.0246 (8)	0.0070 (6)	0.0058 (6)	0.0082 (6)
C22	0.0277 (8)	0.0348 (9)	0.0354 (9)	0.0118 (7)	0.0120 (7)	0.0013 (8)
C23	0.0336 (9)	0.0421 (11)	0.0464 (12)	0.0112 (8)	0.0071 (9)	-0.0131 (9)
C24	0.0313 (9)	0.0422 (10)	0.0293 (9)	0.0134 (8)	-0.0009 (7)	0.0109 (8)
C25	0.0484 (11)	0.0707 (13)	0.0388 (10)	0.0367 (10)	0.0196 (9)	0.0309 (10)

Geometric parameters (Å, °)

C11—C4	1.7454 (15)	C11—H11C	0.9800
C12—C2	1.7376 (15)	C12—C13	1.504 (3)
C13—C16	1.7374 (15)	C12—H12A	0.9900
C14—C18	1.7442 (14)	C12—H12B	0.9900
O1—C7	1.3980 (17)	C13—C13 ⁱ	1.328 (4)
O1—C8	1.433 (2)	C13—C14	1.502 (3)
O2—C7	1.4158 (18)	C14—H14A	0.9900
O2—C10	1.428 (2)	C14—H14B	0.9900
O3—C1	1.3686 (17)	C12B—C13B ⁱ	1.546 (17)
O3—C12	1.434 (3)	C12B—H12C	0.9900
O3—C12B	1.595 (15)	C12B—H12D	0.9900
O4—C21	1.4018 (18)	C13B—C13B ⁱ	1.342 (15)
O4—C22	1.433 (2)	C13B—C12B ⁱ	1.546 (17)
O5—C21	1.4047 (19)	C13B—C14B	1.587 (18)
O5—C24	1.433 (2)	C14B—H14C	0.9900
O6—C15	1.3779 (16)	C14B—H14D	0.9900
O6—C14	1.451 (3)	C15—C16	1.392 (2)
O6—C14B	1.522 (16)	C15—C20	1.400 (2)
C1—C2	1.397 (2)	C16—C17	1.393 (2)
C1—C6	1.400 (2)	C17—C18	1.384 (2)
C2—C3	1.386 (2)	C17—H17A	0.9500
C3—C4	1.381 (2)	C18—C19	1.381 (2)
C3—H3A	0.9500	C19—C20	1.390 (2)
C4—C5	1.378 (2)	C19—H19A	0.9500
C5—C6	1.393 (2)	C20—C21	1.519 (2)
C5—H5A	0.9500	C21—H21A	1.0000
C6—C7	1.524 (2)	C22—C23	1.504 (3)
C7—H7A	1.0000	C22—H22A	0.9900
C8—C9	1.509 (2)	C22—H22B	0.9900
C8—H8A	0.9900	C23—H23A	0.9800
C8—H8B	0.9900	C23—H23B	0.9800
C9—H9A	0.9800	C23—H23C	0.9800
C9—H9B	0.9800	C24—C25	1.505 (3)
C9—H9C	0.9800	C24—H24A	0.9900
C10—C11	1.479 (3)	C24—H24B	0.9900
C10—H10A	0.9900	C25—H25A	0.9800
C10—H10B	0.9900	C25—H25B	0.9800
C11—H11A	0.9800	C25—H25C	0.9800
C11—H11B	0.9800		
C7—O1—C8	113.60 (12)	C13—C14—H14A	110.2
C7—O2—C10	115.03 (12)	O6—C14—H14B	110.2
C1—O3—C12	120.44 (13)	C13—C14—H14B	110.2
C1—O3—C12B	110.8 (5)	H14A—C14—H14B	108.5
C21—O4—C22	116.34 (12)	C13B ⁱ —C12B—O3	109.8 (10)
C21—O5—C24	112.49 (12)	C13B ⁱ —C12B—H12C	109.7

C15—O6—C14	114.90 (14)	O3—C12B—H12C	109.7
C15—O6—C14B	109.8 (6)	C13B ⁱ —C12B—H12D	109.7
O3—C1—C2	124.19 (13)	O3—C12B—H12D	109.7
O3—C1—C6	117.07 (13)	H12C—C12B—H12D	108.2
C2—C1—C6	118.61 (13)	C13B ⁱ —C13B—C12B ⁱ	123.3 (10)
C3—C2—C1	121.46 (14)	C13B ⁱ —C13B—C14B	122.6 (10)
C3—C2—C12	117.32 (12)	C12B ⁱ —C13B—C14B	113.9 (9)
C1—C2—C12	121.21 (11)	O6—C14B—C13B	105.5 (10)
C4—C3—C2	118.47 (14)	O6—C14B—H14C	110.6
C4—C3—H3A	120.8	C13B—C14B—H14C	110.6
C2—C3—H3A	120.8	O6—C14B—H14D	110.6
C5—C4—C3	121.80 (14)	C13B—C14B—H14D	110.6
C5—C4—C11	119.71 (12)	H14C—C14B—H14D	108.8
C3—C4—C11	118.50 (12)	O6—C15—C16	120.73 (13)
C4—C5—C6	119.47 (14)	O6—C15—C20	119.87 (13)
C4—C5—H5A	120.3	C16—C15—C20	119.18 (12)
C6—C5—H5A	120.3	C15—C16—C17	121.93 (14)
C5—C6—C1	120.17 (13)	C15—C16—C13	119.22 (11)
C5—C6—C7	121.03 (13)	C17—C16—C13	118.82 (11)
C1—C6—C7	118.79 (12)	C18—C17—C16	117.21 (14)
O1—C7—O2	112.96 (12)	C18—C17—H17A	121.4
O1—C7—C6	106.45 (11)	C16—C17—H17A	121.4
O2—C7—C6	113.03 (12)	C19—C18—C17	122.55 (13)
O1—C7—H7A	108.1	C19—C18—C14	118.80 (12)
O2—C7—H7A	108.1	C17—C18—C14	118.63 (11)
C6—C7—H7A	108.1	C18—C19—C20	119.52 (14)
O1—C8—C9	107.14 (14)	C18—C19—H19A	120.2
O1—C8—H8A	110.3	C20—C19—H19A	120.2
C9—C8—H8A	110.3	C19—C20—C15	119.61 (13)
O1—C8—H8B	110.3	C19—C20—C21	122.06 (13)
C9—C8—H8B	110.3	C15—C20—C21	118.30 (12)
H8A—C8—H8B	108.5	O4—C21—O5	113.06 (13)
C8—C9—H9A	109.5	O4—C21—C20	113.33 (12)
C8—C9—H9B	109.5	O5—C21—C20	108.04 (12)
H9A—C9—H9B	109.5	O4—C21—H21A	107.4
C8—C9—H9C	109.5	O5—C21—H21A	107.4
H9A—C9—H9C	109.5	C20—C21—H21A	107.4
H9B—C9—H9C	109.5	O4—C22—C23	106.83 (14)
O2—C10—C11	109.21 (15)	O4—C22—H22A	110.4
O2—C10—H10A	109.8	C23—C22—H22A	110.4
C11—C10—H10A	109.8	O4—C22—H22B	110.4
O2—C10—H10B	109.8	C23—C22—H22B	110.4
C11—C10—H10B	109.8	H22A—C22—H22B	108.6
H10A—C10—H10B	108.3	C22—C23—H23A	109.5
C10—C11—H11A	109.5	C22—C23—H23B	109.5
C10—C11—H11B	109.5	H23A—C23—H23B	109.5
H11A—C11—H11B	109.5	C22—C23—H23C	109.5
C10—C11—H11C	109.5	H23A—C23—H23C	109.5

H11A—C11—H11C	109.5	H23B—C23—H23C	109.5
H11B—C11—H11C	109.5	O5—C24—C25	108.28 (14)
O3—C12—C13	105.59 (18)	O5—C24—H24A	110.0
O3—C12—H12A	110.6	C25—C24—H24A	110.0
C13—C12—H12A	110.6	O5—C24—H24B	110.0
O3—C12—H12B	110.6	C25—C24—H24B	110.0
C13—C12—H12B	110.6	H24A—C24—H24B	108.4
H12A—C12—H12B	108.8	C24—C25—H25A	109.5
C13 ⁱ —C13—C14	123.5 (2)	C24—C25—H25B	109.5
C13 ⁱ —C13—C12	123.9 (2)	H25A—C25—H25B	109.5
C14—C13—C12	112.58 (17)	C24—C25—H25C	109.5
O6—C14—C13	107.35 (19)	H25A—C25—H25C	109.5
O6—C14—H14A	110.2	H25B—C25—H25C	109.5
C12—O3—C1—C2	60.8 (2)	C12—C13—C14—O6	82.5 (2)
C12B—O3—C1—C2	69.3 (6)	C1—O3—C12B—C13B ⁱ	133.7 (6)
C12—O3—C1—C6	-123.32 (17)	C12—O3—C12B—C13B ⁱ	-83 (3)
C12B—O3—C1—C6	-114.9 (6)	C15—O6—C14B—C13B	161.1 (6)
O3—C1—C2—C3	174.42 (14)	C14—O6—C14B—C13B	-80 (4)
C6—C1—C2—C3	-1.4 (2)	C13B ⁱ —C13B—C14B—O6	80.1 (12)
O3—C1—C2—C12	-5.0 (2)	C12B ⁱ —C13B—C14B—O6	-104.0 (10)
C6—C1—C2—C12	179.27 (11)	C14—O6—C15—C16	-87.63 (18)
C1—C2—C3—C4	0.9 (2)	C14B—O6—C15—C16	-76.5 (6)
C12—C2—C3—C4	-179.74 (12)	C14—O6—C15—C20	97.79 (17)
C2—C3—C4—C5	0.4 (2)	C14B—O6—C15—C20	108.9 (6)
C2—C3—C4—C11	-179.94 (12)	O6—C15—C16—C17	-174.14 (13)
C3—C4—C5—C6	-1.1 (2)	C20—C15—C16—C17	0.5 (2)
C11—C4—C5—C6	179.24 (12)	O6—C15—C16—C13	4.17 (19)
C4—C5—C6—C1	0.6 (2)	C20—C15—C16—C13	178.78 (11)
C4—C5—C6—C7	179.39 (14)	C15—C16—C17—C18	-0.1 (2)
O3—C1—C6—C5	-175.45 (13)	C13—C16—C17—C18	-178.42 (11)
C2—C1—C6—C5	0.6 (2)	C16—C17—C18—C19	-0.5 (2)
O3—C1—C6—C7	5.7 (2)	C16—C17—C18—C14	177.93 (11)
C2—C1—C6—C7	-178.22 (14)	C17—C18—C19—C20	0.6 (2)
C8—O1—C7—O2	64.73 (17)	C14—C18—C19—C20	-177.74 (12)
C8—O1—C7—C6	-170.66 (13)	C18—C19—C20—C15	-0.3 (2)
C10—O2—C7—O1	61.96 (17)	C18—C19—C20—C21	177.35 (14)
C10—O2—C7—C6	-58.98 (17)	O6—C15—C20—C19	174.37 (13)
C5—C6—C7—O1	-19.60 (19)	C16—C15—C20—C19	-0.3 (2)
C1—C6—C7—O1	159.23 (13)	O6—C15—C20—C21	-3.3 (2)
C5—C6—C7—O2	104.97 (15)	C16—C15—C20—C21	-177.98 (13)
C1—C6—C7—O2	-76.20 (17)	C22—O4—C21—O5	60.38 (17)
C7—O1—C8—C9	155.99 (14)	C22—O4—C21—C20	-63.00 (18)
C7—O2—C10—C11	-173.98 (16)	C24—O5—C21—O4	66.86 (17)
C1—O3—C12—C13	155.76 (13)	C24—O5—C21—C20	-166.88 (13)
C12B—O3—C12—C13	116 (3)	C19—C20—C21—O4	111.86 (16)
O3—C12—C13—C13 ⁱ	-110.9 (2)	C15—C20—C21—O4	-70.51 (18)
O3—C12—C13—C14	68.6 (2)	C19—C20—C21—O5	-14.2 (2)

C15—O6—C14—C13	-172.50 (14)	C15—C20—C21—O5	163.40 (13)
C14B—O6—C14—C13	122 (4)	C21—O4—C22—C23	-172.83 (15)
C13 ⁱ —C13—C14—O6	-98.1 (3)	C21—O5—C24—C25	166.58 (15)

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C12—H12B...C12	0.99	2.66	3.180 (3)	113
C14—H14B...O3	0.99	2.53	2.918 (2)	103
C14—H14B...O4	0.99	2.50	3.166 (3)	125
C17—H17A...O2 ⁱⁱ	0.95	2.47	3.3943 (18)	166

Symmetry code: (ii) $-x+2, -y+1, -z+1$.