

## 9-(4-Methoxyphenyl)anthracene

V. Silambarasan,<sup>a</sup> T. Srinivasan,<sup>a</sup> R. Sivasakthikumaran,<sup>b</sup>  
A. K. Mohanakrishnan<sup>b</sup> and D. Velmurugan<sup>a\*</sup>

<sup>a</sup>CAS in Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai-25, India, and <sup>b</sup>Department of Organic Chemistry, University of Madras, Guindy Campus, Chennai-25, India

Correspondence e-mail: shirai2011@gmail.com


Received 22 October 2012; accepted 16 November 2012

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.045;  $wR$  factor = 0.130; data-to-parameter ratio = 18.9.

In the title compound,  $\text{C}_{21}\text{H}_{16}\text{O}$ , the dihedral angle between the anthracene ring system and the benzene ring is  $74.3(5)^\circ$ . The anthracene ring system is essentially planar (r.m.s. deviation =  $0.0257\text{ \AA}$ ) and the methoxy group lies in the plane of the benzene ring [ $\text{C}1-\text{O}1-\text{C}2-\text{C}7$  torsion angle =  $0.5(2)^\circ$ ]. The crystal structure features  $\pi-\pi$  [centroid–centroid distance =  $3.9487(12)\text{ \AA}$ ] and  $\text{C}-\text{H}\cdots\pi$  interactions, forming a sheet running along the  $a$ -axis direction.

## Related literature

For applications of anthracene, see: Bae *et al.* (2010); Debbab *et al.* (2012). For a related structure, see: Wang *et al.* (2008).



## Experimental

## Crystal data

$\text{C}_{21}\text{H}_{16}\text{O}$   
 $M_r = 284.34$   
Monoclinic,  $P2_1/c$   
 $a = 13.5539(5)\text{ \AA}$

$b = 15.0626(5)\text{ \AA}$   
 $c = 7.6130(2)\text{ \AA}$   
 $\beta = 99.219(2)^\circ$   
 $V = 1534.17(9)\text{ \AA}^3$

$Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.07\text{ mm}^{-1}$

$T = 293\text{ K}$   
 $0.20 \times 0.20 \times 0.20\text{ mm}$

## Data collection

Bruker SMART APEXII area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Bruker, 2008)  
 $T_{\min} = 0.981$ ,  $T_{\max} = 0.985$

14917 measured reflections  
3806 independent reflections  
2317 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.033$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.130$   
 $S = 1.02$   
3806 reflections

201 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.14\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.12\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$\text{Cg}1$  and  $\text{Cg}4$  are the centroids of the C2–C7 and C16–C21 rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}7-\text{H}7\cdots\text{Cg}4^{\text{i}}$	0.93	2.77	3.566 (2)	145
$\text{C}11-\text{H}11\cdots\text{Cg}1^{\text{ii}}$	0.93	2.87	3.724 (2)	154
$\text{C}19-\text{H}19\cdots\text{Cg}1^{\text{iii}}$	0.93	2.94	3.772 (2)	150
$\text{C}21-\text{H}21\cdots\text{Cg}4^{\text{iv}}$	0.93	2.88	3.711 (2)	150

Symmetry codes: (i)  $x, y, z - 1$ ; (ii)  $-x + 1, -y, -z$ ; (iii)  $-x + 2, -y, -z + 1$ ; (iv)  $x, -y - \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 2012); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

The authors thank the TBI X-ray facility, CAS in Crystallography and Biophysics, University of Madras, India, for the data collection. VS and DV also thank the UGC SAP for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2599).

## References

- Bae, S. Y., Jung, K. H., Hoang, M. H., Kim, K. H., Lee, T. W., Cho, M. J., Jin, J., Lee, D. H., Chung, D. S., Park, C. E. & Choi, D. H. (2010). *Synth. Met.* **160**, 1022–1029.  
Bruker (2008). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Debbab, A., Aly, A. H., Edrada-Ebel, R., Wray, V., Pretsch, A., Pescitelli, G., Kurtan, T. & Proksch, P. (2012). *Eur. J. Org. Chem.* pp. 1351–1359.  
Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.  
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.  
Wang, L., You, W., Huang, W., Jiang, J.-C. & Yao, C. (2008). *Acta Cryst. E* **64**, o487.

# supporting information

*Acta Cryst.* (2012). E68, o3410 [doi:10.1107/S1600536812047149]

## **9-(4-Methoxyphenyl)anthracene**

**V. Silambarasan, T. Srinivasan, R. Sivasakthikumaran, A. K. Mohanakrishnan and D. Velmurugan**

### **S1. Comment**

Anthracene is a solid polycyclic aromatic hydrocarbon consisting of three fused benzene rings. Its derivatives possess antimicrobial activity (Debbab *et al.*, 2012). It is used in the production of dyes and organic semiconductors (Bae *et al.*, 2010). In order to obtain detailed information on molecular conformations in the solid state, an X-ray crystallographic study of the title compound was carried out.

The anthracene ring system in the title molecule (Fig. 1) is essentially planar ( $\text{rmsd} = 0.0257 \text{ \AA}$ ) and the methoxy group lies in the plane of the benzene ring with the torsion angle  $C1—O1—C2—C7 = 0.5 (2)^\circ$ . The dihedral angle between the mean-planes of the anthracene and benzene rings is  $74.3 (5)^\circ$  showing that both the ring systems are almost perpendicular to each other. The bond lengths and angles in the title compound are comparable to those observed in a closely related compound (Wang *et al.*, 2008).


The  $\pi\cdots\pi$  electron interaction is observed between the rings ( $C9—C14$ ) [at  $x, y, z$ ] and ( $C9—C14$ ) [at  $1 - x, -y, 1 - z$ ] with the centroid-centroid distance  $3.9487 (12) \text{ \AA}$ . In addition, the crystal packing is stabilized by C–H $\cdots\pi$  (Table. 1) types of interaction.

### **S2. Experimental**


(4-Methoxyphenyl)(2-(phenyl(pivaloyloxy)methyl)phenyl)methyl)pivalate (0.5 g, 0.94 mmol) upon interaction with  $ZnBr_2$  (0.02 g, 0.13 mmol) followed by removal of solvent and column chromatographic purification (silica gel; hexane–ethyl acetate, 99:1) led to the isolation of product as a pale yellow solid (0.32 g, 87%). The compound was recrystallized from chloroform. Single crystals suitable for X-ray diffraction analysis were obtained by slow evaporation of a solution of the title compound in acetone at room temperature.

### **S3. Refinement**

All H atoms were fixed geometrically and allowed to ride on their parent C atoms, with C–H distances 0.93 and  $0.96 \text{ \AA}$  for aryl and methyl H-atoms with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{methyl-C})$  and  $1.2U_{\text{eq}}(\text{aryl-C})$ .

**Figure 1**

The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are presented as small spheres of arbitrary radius.

**Figure 2**

A view of the  $\pi\cdots\pi$  and C—H··· $\pi$  interactions (dotted lines) in the crystal structure of the title compound. H atoms non-participating in hydrogen-bonding were omitted for clarity.

### **9-(4-Methoxyphenyl)anthracene**

#### *Crystal data*

$C_{21}H_{16}O$

$M_r = 284.34$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 13.5539 (5)$  Å

$b = 15.0626 (5)$  Å

$c = 7.6130 (2)$  Å

$\beta = 99.219 (2)^\circ$

$V = 1534.17 (9)$  Å<sup>3</sup>

$Z = 4$

$F(000) = 600$

$D_x = 1.231$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 3806 reflections

$\theta = 2.0\text{--}28.3^\circ$

$\mu = 0.07$  mm<sup>-1</sup>

$T = 293\text{ K}$   
Block, colourless

$0.20 \times 0.20 \times 0.20\text{ mm}$

#### Data collection

Bruker SMART APEXII area-detector  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  and  $\varphi$  scans  
Absorption correction: multi-scan  
(*SADABS*; Bruker, 2008)  
 $T_{\min} = 0.981$ ,  $T_{\max} = 0.985$

14917 measured reflections  
3806 independent reflections  
2317 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.033$   
 $\theta_{\max} = 28.3^\circ$ ,  $\theta_{\min} = 2.0^\circ$   
 $h = -18 \rightarrow 16$   
 $k = -19 \rightarrow 19$   
 $l = -10 \rightarrow 10$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.130$   
 $S = 1.02$   
3806 reflections  
201 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods  
Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0556P)^2 + 0.1555P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.14\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.12\text{ e \AA}^{-3}$   
Extinction correction: *SHELXL*,  
 $Fc^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$   
Extinction coefficient: 0.019 (2)

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.84535 (8)	-0.16715 (6)	-0.12133 (14)	0.0625 (3)
C5	0.75001 (9)	0.00335 (9)	0.24854 (16)	0.0444 (3)
C8	0.71622 (11)	0.06367 (9)	0.38283 (16)	0.0469 (3)
C4	0.73498 (10)	-0.08803 (9)	0.25059 (19)	0.0513 (4)
H4	0.7022	-0.1127	0.3373	0.062*
C6	0.79836 (11)	0.03687 (9)	0.11597 (18)	0.0543 (4)
H6	0.8091	0.0978	0.1112	0.065*
C2	0.81595 (9)	-0.10761 (9)	-0.00379 (17)	0.0471 (3)
C17	0.78712 (11)	0.10166 (9)	0.51685 (17)	0.0484 (3)
C9	0.61419 (11)	0.08466 (10)	0.37219 (18)	0.0542 (4)
C3	0.76772 (11)	-0.14246 (9)	0.12656 (19)	0.0526 (4)
H3	0.7572	-0.2034	0.1309	0.063*

C16	0.75482 (13)	0.16124 (9)	0.64257 (18)	0.0571 (4)
C18	0.89164 (11)	0.08456 (10)	0.53287 (18)	0.0553 (4)
H18	0.9144	0.0464	0.4521	0.066*
C14	0.58323 (13)	0.14597 (11)	0.4959 (2)	0.0633 (4)
C7	0.83127 (11)	-0.01703 (9)	-0.00955 (18)	0.0535 (4)
H7	0.8635	0.0074	-0.0972	0.064*
C19	0.95862 (14)	0.12230 (11)	0.6623 (2)	0.0687 (5)
H19	1.0264	0.1101	0.6690	0.082*
C15	0.65393 (14)	0.18180 (10)	0.6282 (2)	0.0681 (5)
H15	0.6333	0.2208	0.7099	0.082*
C21	0.82821 (16)	0.19878 (11)	0.7773 (2)	0.0730 (5)
H21	0.8080	0.2371	0.8605	0.088*
C10	0.53910 (12)	0.04832 (12)	0.2391 (2)	0.0710 (5)
H10	0.5573	0.0077	0.1581	0.085*
C20	0.92599 (16)	0.18010 (12)	0.7871 (2)	0.0772 (5)
H20	0.9723	0.2054	0.8765	0.093*
C1	0.89552 (14)	-0.13330 (12)	-0.2567 (2)	0.0788 (5)
H1A	0.8533	-0.0912	-0.3275	0.118*
H1B	0.9110	-0.1812	-0.3309	0.118*
H1C	0.9562	-0.1046	-0.2035	0.118*
C13	0.47972 (16)	0.16896 (14)	0.4771 (3)	0.0847 (6)
H13	0.4587	0.2095	0.5555	0.102*
C11	0.44201 (14)	0.07163 (15)	0.2281 (3)	0.0893 (6)
H11	0.3945	0.0468	0.1402	0.107*
C12	0.41231 (16)	0.13311 (17)	0.3484 (3)	0.0960 (7)
H12	0.3454	0.1491	0.3386	0.115*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0721 (7)	0.0500 (6)	0.0713 (7)	0.0036 (5)	0.0293 (6)	-0.0138 (5)
C5	0.0457 (7)	0.0464 (7)	0.0423 (6)	0.0033 (6)	0.0101 (6)	-0.0003 (6)
C8	0.0587 (9)	0.0420 (7)	0.0433 (7)	0.0043 (6)	0.0183 (6)	0.0026 (6)
C4	0.0539 (8)	0.0488 (8)	0.0546 (8)	-0.0021 (6)	0.0190 (7)	0.0041 (6)
C6	0.0739 (10)	0.0397 (7)	0.0537 (8)	-0.0032 (7)	0.0235 (7)	-0.0027 (6)
C2	0.0444 (7)	0.0456 (7)	0.0526 (7)	0.0043 (6)	0.0112 (6)	-0.0073 (6)
C17	0.0664 (9)	0.0384 (7)	0.0439 (7)	-0.0020 (6)	0.0198 (6)	0.0027 (6)
C9	0.0628 (9)	0.0552 (9)	0.0485 (7)	0.0108 (7)	0.0207 (7)	0.0060 (7)
C3	0.0572 (9)	0.0384 (7)	0.0645 (9)	-0.0013 (6)	0.0169 (7)	-0.0005 (6)
C16	0.0865 (11)	0.0409 (7)	0.0494 (8)	-0.0037 (7)	0.0269 (8)	-0.0005 (6)
C18	0.0656 (10)	0.0521 (8)	0.0504 (8)	-0.0068 (7)	0.0163 (7)	-0.0022 (7)
C14	0.0776 (11)	0.0613 (10)	0.0580 (9)	0.0177 (8)	0.0322 (8)	0.0073 (8)
C7	0.0692 (10)	0.0459 (8)	0.0507 (7)	-0.0032 (7)	0.0258 (7)	-0.0015 (6)
C19	0.0736 (11)	0.0719 (11)	0.0605 (9)	-0.0202 (9)	0.0111 (8)	-0.0043 (8)
C15	0.1015 (14)	0.0520 (9)	0.0599 (9)	0.0122 (9)	0.0405 (9)	-0.0029 (7)
C21	0.1192 (16)	0.0489 (9)	0.0561 (9)	-0.0182 (10)	0.0303 (10)	-0.0120 (7)
C10	0.0626 (10)	0.0904 (13)	0.0613 (9)	0.0193 (9)	0.0136 (8)	-0.0032 (9)
C20	0.1028 (15)	0.0692 (11)	0.0601 (10)	-0.0349 (11)	0.0147 (10)	-0.0097 (8)

C1	0.0935 (13)	0.0728 (12)	0.0811 (11)	-0.0002 (10)	0.0480 (10)	-0.0192 (10)
C13	0.0924 (14)	0.0910 (14)	0.0807 (12)	0.0338 (11)	0.0441 (11)	0.0065 (11)
C11	0.0623 (11)	0.1274 (18)	0.0782 (11)	0.0241 (11)	0.0109 (9)	-0.0017 (12)
C12	0.0735 (13)	0.1313 (19)	0.0882 (14)	0.0398 (13)	0.0287 (12)	0.0091 (14)

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

O1—C2	1.3704 (15)	C18—H18	0.9300
O1—C1	1.4176 (18)	C14—C15	1.384 (2)
C5—C6	1.3838 (17)	C14—C13	1.430 (3)
C5—C4	1.3919 (18)	C7—H7	0.9300
C5—C8	1.4935 (17)	C19—C20	1.411 (2)
C8—C17	1.4063 (19)	C19—H19	0.9300
C8—C9	1.408 (2)	C15—H15	0.9300
C4—C3	1.3766 (19)	C21—C20	1.345 (3)
C4—H4	0.9300	C21—H21	0.9300
C6—C7	1.3813 (18)	C10—C11	1.352 (2)
C6—H6	0.9300	C10—H10	0.9300
C2—C3	1.3768 (18)	C20—H20	0.9300
C2—C7	1.3818 (19)	C1—H1A	0.9600
C17—C18	1.426 (2)	C1—H1B	0.9600
C17—C16	1.4309 (19)	C1—H1C	0.9600
C9—C10	1.425 (2)	C13—C12	1.341 (3)
C9—C14	1.429 (2)	C13—H13	0.9300
C3—H3	0.9300	C11—C12	1.406 (3)
C16—C15	1.389 (2)	C11—H11	0.9300
C16—C21	1.426 (2)	C12—H12	0.9300
C18—C19	1.353 (2)		
C2—O1—C1	117.58 (11)	C6—C7—C2	119.44 (12)
C6—C5—C4	117.14 (12)	C6—C7—H7	120.3
C6—C5—C8	120.68 (12)	C2—C7—H7	120.3
C4—C5—C8	122.17 (11)	C18—C19—C20	120.22 (17)
C17—C8—C9	120.07 (12)	C18—C19—H19	119.9
C17—C8—C5	119.69 (12)	C20—C19—H19	119.9
C9—C8—C5	120.21 (13)	C14—C15—C16	121.90 (14)
C3—C4—C5	121.16 (12)	C14—C15—H15	119.1
C3—C4—H4	119.4	C16—C15—H15	119.1
C5—C4—H4	119.4	C20—C21—C16	121.62 (16)
C7—C6—C5	122.23 (13)	C20—C21—H21	119.2
C7—C6—H6	118.9	C16—C21—H21	119.2
C5—C6—H6	118.9	C11—C10—C9	121.45 (17)
O1—C2—C3	116.26 (12)	C11—C10—H10	119.3
O1—C2—C7	124.36 (12)	C9—C10—H10	119.3
C3—C2—C7	119.38 (12)	C21—C20—C19	120.27 (16)
C8—C17—C18	122.87 (12)	C21—C20—H20	119.9
C8—C17—C16	119.58 (13)	C19—C20—H20	119.9
C18—C17—C16	117.54 (13)	O1—C1—H1A	109.5

C8—C9—C10	122.52 (13)	O1—C1—H1B	109.5
C8—C9—C14	119.77 (14)	H1A—C1—H1B	109.5
C10—C9—C14	117.70 (14)	O1—C1—H1C	109.5
C4—C3—C2	120.65 (13)	H1A—C1—H1C	109.5
C4—C3—H3	119.7	H1B—C1—H1C	109.5
C2—C3—H3	119.7	C12—C13—C14	121.20 (17)
C15—C16—C21	122.17 (15)	C12—C13—H13	119.4
C15—C16—C17	119.34 (15)	C14—C13—H13	119.4
C21—C16—C17	118.48 (15)	C10—C11—C12	120.5 (2)
C19—C18—C17	121.86 (14)	C10—C11—H11	119.7
C19—C18—H18	119.1	C12—C11—H11	119.7
C17—C18—H18	119.1	C13—C12—C11	120.56 (18)
C15—C14—C9	119.31 (14)	C13—C12—H12	119.7
C15—C14—C13	122.16 (16)	C11—C12—H12	119.7
C9—C14—C13	118.53 (17)		
C6—C5—C8—C17	-72.83 (17)	C8—C17—C18—C19	179.58 (13)
C4—C5—C8—C17	106.98 (15)	C16—C17—C18—C19	0.5 (2)
C6—C5—C8—C9	105.19 (16)	C8—C9—C14—C15	-2.0 (2)
C4—C5—C8—C9	-74.99 (17)	C10—C9—C14—C15	179.23 (14)
C6—C5—C4—C3	0.5 (2)	C8—C9—C14—C13	177.33 (14)
C8—C5—C4—C3	-179.30 (13)	C10—C9—C14—C13	-1.4 (2)
C4—C5—C6—C7	-0.3 (2)	C5—C6—C7—C2	-0.1 (2)
C8—C5—C6—C7	179.51 (13)	O1—C2—C7—C6	179.96 (13)
C1—O1—C2—C3	-179.79 (13)	C3—C2—C7—C6	0.3 (2)
C1—O1—C2—C7	0.5 (2)	C17—C18—C19—C20	0.3 (2)
C9—C8—C17—C18	-178.74 (12)	C9—C14—C15—C16	1.3 (2)
C5—C8—C17—C18	-0.71 (19)	C13—C14—C15—C16	-178.05 (15)
C9—C8—C17—C16	0.35 (19)	C21—C16—C15—C14	179.21 (14)
C5—C8—C17—C16	178.38 (11)	C17—C16—C15—C14	0.3 (2)
C17—C8—C9—C10	179.89 (13)	C15—C16—C21—C20	-178.28 (15)
C5—C8—C9—C10	1.9 (2)	C17—C16—C21—C20	0.7 (2)
C17—C8—C9—C14	1.2 (2)	C8—C9—C10—C11	-177.86 (16)
C5—C8—C9—C14	-176.82 (12)	C14—C9—C10—C11	0.9 (2)
C5—C4—C3—C2	-0.3 (2)	C16—C21—C20—C19	0.1 (3)
O1—C2—C3—C4	-179.78 (13)	C18—C19—C20—C21	-0.6 (2)
C7—C2—C3—C4	-0.1 (2)	C15—C14—C13—C12	-179.66 (18)
C8—C17—C16—C15	-1.1 (2)	C9—C14—C13—C12	1.0 (3)
C18—C17—C16—C15	178.03 (13)	C9—C10—C11—C12	0.2 (3)
C8—C17—C16—C21	179.92 (12)	C14—C13—C12—C11	0.0 (3)
C18—C17—C16—C21	-0.94 (19)	C10—C11—C12—C13	-0.6 (3)

*Hydrogen-bond geometry (Å, °)*

Cg1 and Cg4 are the centroids of the C2—C7 and C16—C21 rings, respectively.

D—H···A	D—H	H···A	D···A	D—H···A
C7—H7···Cg4 <sup>i</sup>	0.93	2.77	3.566 (2)	145
C11—H11···Cg1 <sup>ii</sup>	0.93	2.87	3.724 (2)	154

---

C19—H19···Cg1 <sup>iii</sup>	0.93	2.94	3.772 (2)	150
C21—H21···Cg4 <sup>iv</sup>	0.93	2.88	3.711 (2)	150

---

Symmetry codes: (i)  $x, y, z-1$ ; (ii)  $-x+1, -y, -z$ ; (iii)  $-x+2, -y, -z+1$ ; (iv)  $x, -y-1/2, z-1/2$ .