

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

catena-Poly[[[bis(4-pyridinealdoxime- κN^1)zinc]- μ -benzene-1,4-dicarboxylato- $\kappa^2 O^1:O^4$] 4-pyridinealdoxime monosolvate]

Hitoshi Kumagai,^{a*} Satoshi Kawata^b and Yoshiyuki Sakamoto^a

^aToyota Central R&D Labs., Inc., Nagakute 41-1, Aichi, Japan, and ^bDepartment of Chemistry, Fukuoka University, Fukuoka 814-0180, Japan
Correspondence e-mail: e1254@mosk.tytlabs.co.jp


Received 18 February 2013; accepted 4 March 2013

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.030; wR factor = 0.078; data-to-parameter ratio = 15.9.

In the title compound, $\{[Zn(C_8H_4O_4)(C_6H_6N_2O)_2] \cdot C_6H_6N_2O\}_n$, the Zn^{II} ion exhibits a tetrahedral coordination environment defined by two benzene-1,4-dicarboxylate dianions and two 4-pyridinealdoxime ligands. The dianions bridge the Zn^{II} ions, giving a zigzag chain along the b axis. Adjacent chains are connected by $O-H \cdots O$ hydrogen bonds, forming a cavity in which an uncoordinating 4-pyridinealdoxime molecule is located; this molecule is linked by $O-H \cdots O$ and $O-H \cdots N$ hydrogen bonds to the zigzag chain.

Related literature

For coordination polymers, see: Cheetham *et al.* (1999); Furukawa *et al.* (2010). For related host-guest systems, see: Kitagawa & Kawata (2002); Lehn (1995).



Experimental

Crystal data

$[Zn(C_8H_4O_4)(C_6H_6N_2O)_2] \cdot C_6H_6N_2O$
 $M_r = 595.88$
 Monoclinic, $P2_1/c$
 $a = 7.583$ (3) Å
 $b = 15.831$ (6) Å
 $c = 21.906$ (8) Å
 $\beta = 98.516$ (3)°
 $V = 2600.7$ (16) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.00$ mm⁻¹
 $T = 293$ K
 $0.60 \times 0.40 \times 0.15$ mm

Data collection

Rigaku Mercury70 diffractometer
 Absorption correction: multi-scan (*REQAB*; Rigaku, 1998)
 $T_{min} = 0.714$, $T_{max} = 0.860$
 24675 measured reflections
 5922 independent reflections
 5402 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.078$
 $S = 1.06$
 5922 reflections
 373 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{max} = 0.34$ e Å⁻³
 $\Delta\rho_{min} = -0.27$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$O5-H3 \cdots N5^i$	0.89 (5)	1.81 (4)	2.692 (3)	172 (4)
$O6-H9 \cdots O4^{ii}$	0.81 (3)	1.94 (3)	2.752 (3)	177 (3)
$O7-H13 \cdots O4^i$	0.77 (4)	2.07 (4)	2.800 (3)	158 (4)

Symmetry codes: (i) $x + 1, y, z$; (ii) $x + 1, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku/MSC, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SIR2008* (Burla *et al.*, 2007); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *CrystalStructure* (Rigaku, 2010); software used to prepare material for publication: *CrystalStructure*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5248).

References

- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giovacazzo, C., Polidori, G., Siliqi, D. & Spagna, R. (2007). *J. Appl. Cryst.* **40**, 609–613.
- Cheetham, A. K., Ferey, G. & Loiseau, T. (1999). *Angew. Chem. Int. Ed.* **38**, 3268–3292.
- Furukawa, H., Ko, N., Go, Y. B., Aratani, N., Choi, S. B., Choi, E., Yazaydin, A. O., Snurr, R. Q., O'Keeffe, M., Kim, J. & Yaghi, O. M. (2010). *Science*, **329**, 424–428.
- Kitagawa, S. & Kawata, S. (2002). *Coord. Chem. Rev.* **224**, 11–34.
- Lehn, J.-M. (1995). In *Supramolecular Chemistry: Concepts and Perspectives*. VCH: Weinheim.
- Rigaku (1998). *REQAB*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (2010). *CrystalStructure*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2013). E69, m216 [doi:10.1107/S1600536813006107]

catena-Poly[[[bis(4-pyridinealdoxime- κN^1)zinc]- μ -benzene-1,4-dicarboxylato- $\kappa^2 O^1:O^4$] 4-pyridinealdoxime monosolvate]**Hitoshi Kumagai, Satoshi Kawata and Yoshiyuki Sakamoto****S1. Comment**


The design and synthesis of coordination polymers (CPs) have received considerable attention in recent years due to potential applications for magnetic materials, sorption and host–guest materials (Cheetham *et al.*, 1999; Furukawa *et al.*, 2010; Lehn, 1995). Intermolecular interactions, such as coordination bonding, hydrogen bonding and van der Waals forces, have been used to produce such network materials. Among them, hydrogen bonding interaction is an important interaction to realize self-assemblies of molecules with novel properties such as proton-transfer-mediated electron transfer, non-linear optics and thermochromic properties of crystalline forms. We have focused on the synthesis and characterization of one-dimensional coordination polymers as host materials and have found a number of host–guest systems using hydrogen bonding interactions (Kitagawa & Kawata, 2002). Here we report synthesis and single-crystal structure of a new one-dimensional coordination polymer which consists of tetrahedral Zn(II) ion, 1,4-benzene-dicarboxylate as a bridging ligand and 4-pyridineoxime (4-pyNOH) as terminal ligands. Two types of 4-pyridinealdoxime are found in the crystal. One is uncoordinated and the other 4-pyNOH coordinates to Zn(II) ion. Uncoordinated 4-pyNOH molecules are stabilized by intermolecular hydrogen bonds in the cavity formed by hydrogen bonds between chains.

S2. Experimental

An aqueous solution (5 ml) of zinc nitrate hexahydrate (0.29 g) was transferred to a glass tube, then an ethanol-water mixture (5 ml) of terephthalic acid (0.17 g), NaOH (0.08 g) and 4-pyNOH (0.24 g) was poured into the glass tube without mixing the two solutions. Colorless crystals began to form at ambient temperature in 2 months. One of these crystals was used for X-ray crystallography.

S3. Refinement

Hydrogen atoms bonded to carbon atoms were introduced at the positions calculated theoretically and treated with riding models with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. Other hydrogen atoms (H3, H9, H13) were located in a Fourier difference map and refined freely [O—H = 0.77 (4)–0.89 (5) Å].

**Figure 1**

View of the title compound with atomic numbering scheme. Hydrogen atoms are omitted for clarity.




Figure 2

Hydrogen bonding interactions between chains (blue and red) and intercalated 4-pyNOH molecules. Hydrogen atoms and hydrogen bonding interactions are shown as purple color and dashed line, respectively.

catena-Poly[[[bis(4-pyridinealdoxime- κN^1)zinc]- μ -benzene-1,4-dicarboxylato- $\kappa^2 O^1:O^4$] 4-pyridinealdoxime monosolvate]

Crystal data

$[\text{Zn}(\text{C}_8\text{H}_4\text{O}_4)(\text{C}_6\text{H}_6\text{N}_2\text{O})_2] \cdot \text{C}_6\text{H}_6\text{N}_2\text{O}$

$M_r = 595.88$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 7.583 (3) \text{ \AA}$

$b = 15.831 (6) \text{ \AA}$

$c = 21.906 (8) \text{ \AA}$

$\beta = 98.516 (3)^\circ$

$V = 2600.7 (16) \text{ \AA}^3$

$Z = 4$

$F(000) = 1224.00$

$D_x = 1.522 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71070 \text{ \AA}$

Cell parameters from 5655 reflections

$\theta = 3.0\text{--}27.5^\circ$

$\mu = 1.00 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Platelet, colorless

$0.60 \times 0.40 \times 0.15 \text{ mm}$

Data collection

Rigaku Mercury70

diffractometer

Detector resolution: $7.314 \text{ pixels mm}^{-1}$

ω scans

Absorption correction: multi-scan

(*REQAB*; Rigaku, 1998)

$T_{\min} = 0.714$, $T_{\max} = 0.860$

24675 measured reflections

5922 independent reflections

5402 reflections with $F^2 > 2\sigma(F^2)$

$R_{\text{int}} = 0.019$

$\theta_{\max} = 27.5^\circ$

$h = -9 \rightarrow 9$

$k = -20 \rightarrow 20$

$l = -28 \rightarrow 28$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.078$
 $S = 1.06$
 5922 reflections
 373 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0395P)^2 + 0.8505P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.34 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

Special details

Refinement. Refinement was performed using all reflections. The weighted R -factor (wR) and goodness of fit (S) are based on F^2 . R -factor (gt) are based on F . The threshold expression of $F^2 > 2.0 \sigma(F^2)$ is used only for calculating R -factor (gt).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	-0.06451 (2)	0.622158 (11)	0.714672 (8)	0.03146 (7)
O1	-0.21883 (16)	0.54529 (7)	0.74818 (6)	0.0431 (3)
O2	0.00514 (16)	0.45411 (8)	0.76070 (7)	0.0509 (4)
O3	-0.79388 (15)	0.22499 (7)	0.79980 (6)	0.0421 (3)
O4	-0.58272 (17)	0.16416 (8)	0.86719 (6)	0.0474 (3)
O5	0.9256 (2)	0.74280 (12)	0.91104 (8)	0.0654 (5)
O6	0.3534 (2)	0.50254 (9)	0.39196 (7)	0.0525 (4)
O7	0.4650 (4)	0.24025 (15)	0.98388 (10)	0.0887 (7)
N1	0.15225 (17)	0.66275 (8)	0.77281 (6)	0.0339 (3)
N2	0.7742 (2)	0.70378 (11)	0.88055 (7)	0.0503 (4)
N3	0.02911 (17)	0.59469 (8)	0.63468 (6)	0.0330 (3)
N4	0.30194 (19)	0.49616 (9)	0.45026 (7)	0.0415 (4)
N5	0.1156 (3)	0.60383 (14)	0.94885 (9)	0.0614 (5)
N6	0.3640 (3)	0.30885 (14)	0.95959 (9)	0.0680 (5)
C1	0.2999 (3)	0.61485 (10)	0.78018 (9)	0.0417 (4)
C2	0.4614 (3)	0.64161 (11)	0.81024 (9)	0.0427 (4)
C3	0.4744 (3)	0.72219 (11)	0.83570 (7)	0.0381 (4)
C4	0.3214 (3)	0.77116 (11)	0.82951 (8)	0.0407 (4)
C5	0.1641 (3)	0.74000 (10)	0.79765 (8)	0.0376 (4)
C6	0.6445 (3)	0.75430 (13)	0.86763 (9)	0.0491 (5)
C7	-0.0049 (3)	0.65017 (10)	0.58845 (8)	0.0369 (4)
C8	0.0624 (3)	0.64167 (11)	0.53383 (8)	0.0409 (4)
C9	0.1692 (3)	0.57299 (10)	0.52502 (8)	0.0376 (4)
C10	0.2016 (3)	0.51452 (11)	0.57240 (8)	0.0414 (4)
C11	0.1296 (3)	0.52711 (11)	0.62583 (8)	0.0405 (4)
C12	0.2375 (3)	0.56456 (12)	0.46607 (9)	0.0474 (5)
C13	-0.2764 (2)	0.41148 (9)	0.78608 (7)	0.0318 (3)
C14	-0.4586 (2)	0.42012 (9)	0.76680 (8)	0.0356 (4)
C15	-0.5772 (2)	0.36017 (10)	0.78215 (8)	0.0357 (4)

C16	-0.5140 (2)	0.29122 (9)	0.81793 (7)	0.0314 (3)
C17	-0.3329 (2)	0.28438 (10)	0.83923 (8)	0.0363 (4)
C18	-0.2141 (2)	0.34347 (10)	0.82278 (8)	0.0359 (4)
C19	-0.1499 (2)	0.47331 (10)	0.76378 (7)	0.0344 (4)
C20	-0.6376 (2)	0.22133 (9)	0.83015 (8)	0.0348 (4)
C21	0.1206 (3)	0.54202 (17)	0.90740 (10)	0.0629 (6)
C22	0.2009 (3)	0.46555 (16)	0.92026 (9)	0.0603 (6)
C23	0.2809 (3)	0.44822 (15)	0.98042 (9)	0.0547 (5)
C24	0.2726 (3)	0.51129 (17)	1.02396 (10)	0.0637 (6)
C25	0.1912 (3)	0.58644 (18)	1.00653 (11)	0.0670 (7)
C26	0.3730 (4)	0.36927 (16)	0.99751 (10)	0.0631 (6)
H1	0.2921	0.5605	0.7640	0.0500*
H2	0.5607	0.6065	0.8136	0.0512*
H3	0.990 (6)	0.699 (3)	0.9271 (18)	0.131 (15)*
H4	0.3247	0.8249	0.8468	0.0488*
H5	0.0629	0.7739	0.7933	0.0451*
H6	0.6572	0.8111	0.8783	0.0590*
H7	-0.0770	0.6964	0.5934	0.0443*
H8	0.0364	0.6819	0.5029	0.0490*
H9	0.376 (4)	0.4540 (18)	0.3841 (12)	0.075 (9)*
H10	0.2715	0.4672	0.5682	0.0497*
H11	0.1514	0.4871	0.6571	0.0485*
H12	0.2331	0.6109	0.4398	0.0568*
H13	0.456 (5)	0.208 (3)	0.9572 (16)	0.105 (13)*
H14	-0.5012	0.4668	0.7433	0.0427*
H15	-0.6987	0.3660	0.7685	0.0429*
H17	-0.2909	0.2396	0.8648	0.0436*
H18	-0.0926	0.3376	0.8364	0.0431*
H21	0.0658	0.5518	0.8672	0.0755*
H22	0.2023	0.4255	0.8892	0.0723*
H24	0.3222	0.5025	1.0649	0.0764*
H25	0.1883	0.6278	1.0365	0.0804*
H26	0.4388	0.3632	1.0366	0.0757*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.02747 (10)	0.02856 (10)	0.03936 (11)	0.00125 (6)	0.00831 (7)	0.00241 (7)
O1	0.0429 (7)	0.0306 (6)	0.0598 (8)	-0.0038 (5)	0.0205 (6)	0.0039 (5)
O2	0.0326 (6)	0.0496 (8)	0.0730 (9)	-0.0027 (6)	0.0165 (6)	0.0068 (7)
O3	0.0330 (6)	0.0317 (6)	0.0612 (8)	-0.0067 (5)	0.0050 (6)	0.0005 (5)
O4	0.0488 (7)	0.0361 (7)	0.0571 (8)	-0.0058 (6)	0.0077 (6)	0.0126 (6)
O5	0.0478 (8)	0.0793 (12)	0.0652 (10)	-0.0185 (8)	-0.0049 (7)	-0.0220 (9)
O6	0.0630 (9)	0.0467 (8)	0.0536 (8)	-0.0017 (7)	0.0281 (7)	-0.0102 (6)
O7	0.1144 (17)	0.0847 (14)	0.0620 (12)	0.0132 (12)	-0.0030 (11)	0.0018 (11)
N1	0.0327 (7)	0.0311 (7)	0.0382 (7)	0.0005 (5)	0.0066 (6)	0.0001 (6)
N2	0.0416 (9)	0.0650 (11)	0.0428 (9)	-0.0147 (8)	0.0011 (7)	-0.0138 (8)
N3	0.0293 (7)	0.0324 (7)	0.0377 (7)	-0.0008 (5)	0.0062 (6)	-0.0013 (6)

N4	0.0367 (8)	0.0466 (8)	0.0422 (8)	-0.0037 (6)	0.0091 (6)	-0.0074 (7)
N5	0.0432 (9)	0.0849 (14)	0.0554 (11)	-0.0078 (9)	0.0050 (8)	-0.0060 (10)
N6	0.0698 (13)	0.0799 (14)	0.0540 (11)	-0.0014 (11)	0.0079 (9)	0.0014 (10)
C1	0.0360 (9)	0.0327 (8)	0.0553 (11)	0.0017 (7)	0.0033 (8)	-0.0092 (7)
C2	0.0339 (9)	0.0408 (9)	0.0524 (11)	0.0025 (7)	0.0029 (8)	-0.0083 (8)
C3	0.0400 (9)	0.0413 (9)	0.0335 (8)	-0.0079 (7)	0.0069 (7)	-0.0032 (7)
C4	0.0525 (10)	0.0340 (8)	0.0365 (9)	-0.0024 (7)	0.0098 (8)	-0.0069 (7)
C5	0.0423 (9)	0.0345 (8)	0.0381 (9)	0.0054 (7)	0.0129 (7)	-0.0010 (7)
C6	0.0511 (11)	0.0491 (11)	0.0462 (11)	-0.0129 (9)	0.0037 (9)	-0.0101 (8)
C7	0.0375 (9)	0.0322 (8)	0.0410 (9)	0.0028 (7)	0.0061 (7)	-0.0008 (7)
C8	0.0492 (10)	0.0351 (8)	0.0388 (9)	0.0023 (7)	0.0081 (8)	0.0037 (7)
C9	0.0370 (9)	0.0373 (9)	0.0394 (9)	-0.0036 (7)	0.0086 (7)	-0.0043 (7)
C10	0.0399 (9)	0.0395 (9)	0.0454 (10)	0.0090 (7)	0.0083 (8)	-0.0012 (7)
C11	0.0413 (9)	0.0385 (9)	0.0416 (9)	0.0070 (7)	0.0063 (7)	0.0046 (7)
C12	0.0593 (12)	0.0405 (10)	0.0461 (10)	-0.0002 (8)	0.0204 (9)	-0.0002 (8)
C13	0.0309 (8)	0.0272 (7)	0.0385 (8)	-0.0034 (6)	0.0094 (6)	-0.0041 (6)
C14	0.0328 (8)	0.0256 (7)	0.0491 (9)	0.0022 (6)	0.0085 (7)	0.0036 (7)
C15	0.0274 (8)	0.0311 (8)	0.0490 (10)	0.0002 (6)	0.0066 (7)	0.0018 (7)
C16	0.0313 (8)	0.0270 (7)	0.0371 (8)	-0.0027 (6)	0.0085 (6)	-0.0013 (6)
C17	0.0348 (8)	0.0334 (8)	0.0402 (9)	0.0005 (6)	0.0037 (7)	0.0061 (7)
C18	0.0287 (8)	0.0367 (8)	0.0417 (9)	-0.0014 (6)	0.0029 (7)	0.0006 (7)
C19	0.0357 (8)	0.0313 (8)	0.0373 (8)	-0.0059 (6)	0.0091 (7)	-0.0045 (6)
C20	0.0351 (8)	0.0279 (8)	0.0431 (9)	-0.0031 (6)	0.0111 (7)	-0.0022 (7)
C21	0.0549 (12)	0.0891 (17)	0.0432 (11)	-0.0147 (12)	0.0025 (9)	-0.0012 (11)
C22	0.0608 (13)	0.0814 (16)	0.0392 (10)	-0.0156 (12)	0.0096 (9)	-0.0062 (10)
C23	0.0451 (11)	0.0778 (14)	0.0426 (10)	-0.0173 (10)	0.0114 (8)	-0.0015 (10)
C24	0.0549 (13)	0.0956 (18)	0.0397 (11)	-0.0098 (12)	0.0045 (9)	-0.0071 (11)
C25	0.0539 (13)	0.0950 (19)	0.0515 (13)	-0.0069 (13)	0.0057 (10)	-0.0186 (12)
C26	0.0637 (14)	0.0843 (17)	0.0420 (11)	-0.0127 (12)	0.0102 (10)	0.0050 (11)

Geometric parameters (Å, °)

Zn1—O1	1.9088 (14)	C14—C15	1.383 (3)
Zn1—O3 ⁱ	1.9501 (13)	C15—C16	1.387 (3)
Zn1—N1	2.0289 (13)	C16—C17	1.388 (3)
Zn1—N3	2.0327 (15)	C16—C20	1.500 (3)
O1—C19	1.279 (2)	C17—C18	1.383 (3)
O2—C19	1.226 (2)	C21—C22	1.366 (4)
O3—C20	1.2713 (19)	C22—C23	1.394 (3)
O4—C20	1.245 (2)	C23—C24	1.389 (4)
O5—N2	1.385 (3)	C23—C26	1.453 (4)
O6—N4	1.393 (3)	C24—C25	1.368 (4)
O7—N6	1.388 (4)	O5—H3	0.89 (4)
N1—C1	1.342 (3)	O6—H9	0.81 (3)
N1—C5	1.336 (2)	O7—H13	0.78 (4)
N2—C6	1.267 (3)	C1—H1	0.930
N3—C7	1.336 (3)	C2—H2	0.930
N3—C11	1.344 (3)	C4—H4	0.930

N4—C12	1.258 (3)	C5—H5	0.930
N5—C21	1.339 (4)	C6—H6	0.930
N5—C25	1.337 (3)	C7—H7	0.930
N6—C26	1.262 (4)	C8—H8	0.930
C1—C2	1.369 (3)	C10—H10	0.930
C2—C3	1.390 (3)	C11—H11	0.930
C3—C4	1.385 (3)	C12—H12	0.930
C3—C6	1.464 (3)	C14—H14	0.930
C4—C5	1.381 (3)	C15—H15	0.930
C7—C8	1.375 (3)	C17—H17	0.930
C8—C9	1.386 (3)	C18—H18	0.930
C9—C10	1.385 (3)	C21—H21	0.930
C9—C12	1.467 (3)	C22—H22	0.930
C10—C11	1.378 (3)	C24—H24	0.930
C13—C14	1.390 (3)	C25—H25	0.930
C13—C18	1.384 (3)	C26—H26	0.930
C13—C19	1.502 (3)		
O1—Zn1—O3 ⁱ	103.92 (6)	N5—C21—C22	124.4 (2)
O1—Zn1—N1	116.51 (6)	C21—C22—C23	119.2 (2)
O1—Zn1—N3	120.49 (6)	C22—C23—C24	116.8 (3)
O3 ⁱ —Zn1—N1	102.30 (6)	C22—C23—C26	122.5 (2)
O3 ⁱ —Zn1—N3	107.02 (6)	C24—C23—C26	120.69 (19)
N1—Zn1—N3	104.77 (6)	C23—C24—C25	119.9 (2)
Zn1—O1—C19	114.91 (12)	N5—C25—C24	123.7 (3)
Zn1 ⁱⁱ —O3—C20	119.68 (10)	N6—C26—C23	120.4 (2)
Zn1—N1—C1	118.72 (11)	N2—O5—H3	102 (3)
Zn1—N1—C5	123.01 (11)	N4—O6—H9	103 (2)
C1—N1—C5	117.62 (14)	N6—O7—H13	104 (3)
O5—N2—C6	112.76 (18)	N1—C1—H1	118.183
Zn1—N3—C7	117.37 (12)	C2—C1—H1	118.181
Zn1—N3—C11	124.92 (11)	C1—C2—H2	120.577
C7—N3—C11	117.63 (15)	C3—C2—H2	120.575
O6—N4—C12	111.28 (16)	C3—C4—H4	120.076
C21—N5—C25	116.0 (3)	C5—C4—H4	120.063
O7—N6—C26	111.6 (2)	N1—C5—H5	118.876
N1—C1—C2	123.64 (16)	C4—C5—H5	118.881
C1—C2—C3	118.85 (16)	N2—C6—H6	120.410
C2—C3—C4	117.76 (16)	C3—C6—H6	120.420
C2—C3—C6	121.03 (17)	N3—C7—H7	118.651
C4—C3—C6	121.21 (17)	C8—C7—H7	118.656
C3—C4—C5	119.86 (16)	C7—C8—H8	120.074
N1—C5—C4	122.24 (16)	C9—C8—H8	120.082
N2—C6—C3	119.17 (18)	C9—C10—H10	120.301
N3—C7—C8	122.69 (16)	C11—C10—H10	120.286
C7—C8—C9	119.84 (16)	N3—C11—H11	118.577
C8—C9—C10	117.54 (17)	C10—C11—H11	118.577
C8—C9—C12	118.92 (16)	N4—C12—H12	119.234

C10—C9—C12	123.51 (16)	C9—C12—H12	119.230
C9—C10—C11	119.41 (17)	C13—C14—H14	119.625
N3—C11—C10	122.85 (16)	C15—C14—H14	119.627
N4—C12—C9	121.54 (18)	C14—C15—H15	120.158
C14—C13—C18	119.43 (15)	C16—C15—H15	120.177
C14—C13—C19	119.41 (14)	C16—C17—H17	119.676
C18—C13—C19	121.06 (14)	C18—C17—H17	119.670
C13—C14—C15	120.75 (14)	C13—C18—H18	120.059
C14—C15—C16	119.66 (15)	C17—C18—H18	120.071
C15—C16—C17	119.55 (15)	N5—C21—H21	117.814
C15—C16—C20	120.58 (14)	C22—C21—H21	117.818
C17—C16—C20	119.74 (14)	C21—C22—H22	120.395
C16—C17—C18	120.65 (15)	C23—C22—H22	120.411
C13—C18—C17	119.87 (15)	C23—C24—H24	120.064
O1—C19—O2	124.31 (16)	C25—C24—H24	120.069
O1—C19—C13	114.32 (14)	N5—C25—H25	118.127
O2—C19—C13	121.36 (15)	C24—C25—H25	118.128
O3—C20—O4	124.72 (15)	N6—C26—H26	119.823
O3—C20—C16	115.58 (14)	C23—C26—H26	119.820
O4—C20—C16	119.68 (14)		
O1—Zn1—O3 ⁱ —C20 ⁱ	-51.57 (11)	C2—C3—C6—N2	-11.0 (3)
O3 ⁱ —Zn1—O1—C19	-177.30 (8)	C4—C3—C6—N2	169.38 (16)
O1—Zn1—N1—C1	84.68 (11)	C6—C3—C4—C5	178.20 (15)
O1—Zn1—N1—C5	-104.80 (11)	C3—C4—C5—N1	1.0 (3)
N1—Zn1—O1—C19	-65.65 (10)	N3—C7—C8—C9	0.5 (3)
O1—Zn1—N3—C7	122.99 (9)	C7—C8—C9—C10	0.8 (3)
O1—Zn1—N3—C11	-60.34 (11)	C7—C8—C9—C12	179.15 (14)
N3—Zn1—O1—C19	62.98 (10)	C8—C9—C10—C11	-0.7 (3)
O3 ⁱ —Zn1—N1—C1	-162.75 (9)	C8—C9—C12—N4	-164.73 (16)
O3 ⁱ —Zn1—N1—C5	7.77 (12)	C10—C9—C12—N4	13.5 (3)
N1—Zn1—O3 ⁱ —C20 ⁱ	-173.22 (10)	C12—C9—C10—C11	-179.00 (15)
O3 ⁱ —Zn1—N3—C7	4.81 (10)	C9—C10—C11—N3	-0.6 (3)
O3 ⁱ —Zn1—N3—C11	-178.51 (9)	C14—C13—C18—C17	0.8 (3)
N3—Zn1—O3 ⁱ —C20 ⁱ	76.92 (11)	C18—C13—C14—C15	-2.2 (3)
N1—Zn1—N3—C7	-103.31 (9)	C14—C13—C19—O1	25.0 (2)
N1—Zn1—N3—C11	73.36 (11)	C14—C13—C19—O2	-154.27 (14)
N3—Zn1—N1—C1	-51.20 (10)	C19—C13—C14—C15	174.36 (13)
N3—Zn1—N1—C5	119.32 (10)	C18—C13—C19—O1	-158.55 (14)
Zn1—O1—C19—O2	3.1 (2)	C18—C13—C19—O2	22.2 (3)
Zn1—O1—C19—C13	-176.10 (8)	C19—C13—C18—C17	-175.65 (13)
Zn1 ⁱⁱ —O3—C20—O4	-22.5 (3)	C13—C14—C15—C16	0.9 (3)
Zn1 ⁱⁱ —O3—C20—C16	156.16 (9)	C14—C15—C16—C17	1.7 (3)
Zn1—N1—C1—C2	169.39 (12)	C14—C15—C16—C20	-174.19 (14)
Zn1—N1—C5—C4	-170.06 (10)	C15—C16—C17—C18	-3.0 (3)
C1—N1—C5—C4	0.6 (3)	C15—C16—C20—O3	8.3 (3)
C5—N1—C1—C2	-1.6 (3)	C15—C16—C20—O4	-172.96 (15)
O5—N2—C6—C3	-179.56 (15)	C17—C16—C20—O3	-167.49 (14)

Zn1—N3—C7—C8	175.10 (10)	C17—C16—C20—O4	11.2 (3)
Zn1—N3—C11—C10	-174.78 (10)	C20—C16—C17—C18	172.87 (14)
C7—N3—C11—C10	1.9 (3)	C16—C17—C18—C13	1.8 (3)
C11—N3—C7—C8	-1.8 (3)	N5—C21—C22—C23	1.4 (4)
O6—N4—C12—C9	178.33 (14)	C21—C22—C23—C24	0.2 (4)
C21—N5—C25—C24	0.9 (4)	C21—C22—C23—C26	-178.39 (19)
C25—N5—C21—C22	-1.9 (4)	C22—C23—C24—C25	-1.0 (4)
O7—N6—C26—C23	178.2 (2)	C22—C23—C26—N6	-9.1 (4)
N1—C1—C2—C3	1.1 (3)	C24—C23—C26—N6	172.4 (2)
C1—C2—C3—C4	0.4 (3)	C26—C23—C24—C25	177.53 (19)
C1—C2—C3—C6	-179.20 (16)	C23—C24—C25—N5	0.5 (4)
C2—C3—C4—C5	-1.4 (3)		

Symmetry codes: (i) $-x-1, y+1/2, -z+3/2$; (ii) $-x-1, y-1/2, -z+3/2$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O5—H3...N5 ⁱⁱⁱ	0.89 (5)	1.81 (4)	2.692 (3)	172 (4)
O6—H9...O4 ^{iv}	0.81 (3)	1.94 (3)	2.752 (3)	177 (3)
O7—H13...O4 ⁱⁱⁱ	0.77 (4)	2.07 (4)	2.800 (3)	158 (4)

Symmetry codes: (iii) $x+1, y, z$; (iv) $x+1, -y+1/2, z-1/2$.