

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,2'-[(4-Methyl-2-phenylimidazolidine-1,3-diyl)bis(methylene)]diphenol

Augusto Rivera,^{a*} Lorena Cárdenas,^a Jaime Ríos-Motta,^a
Václav Eigner,^{b,c} and Michal Dušek^c

^aUniversidad Nacional de Colombia, Sede Bogotá, Facultad de Ciencias, Departamento de Química, Cra 30 No.45-03, Bogotá, Código Postal 111321, Colombia, ^bDepartment of Solid State Chemistry, Institute of Chemical Technology, Technická 5, 166 28 Prague, Czech Republic, and ^cInstitute of Physics AS CR, v.v.i., Na Slovance 2, 182 21 Prague 8, Czech Republic
Correspondence e-mail: ariverau@unal.edu.co


Received 27 June 2013; accepted 28 June 2013

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; disorder in main residue; R factor = 0.042; wR factor = 0.113; data-to-parameter ratio = 12.6.

The methyl-substituted imidazolidine ring of the title compound, $\text{C}_{24}\text{H}_{26}\text{N}_2\text{O}_2$, adopts an envelope conformation with the N atom adjacent to the methylene group as the flap. The methyl-ethylene fragment in this ring is disordered over two positions with an occupancy ratio of 0.899 (4):0.101 (4). The hydroxybenzyl groups are inclined at 71.57 (15) and 69.97 (19)° to the mean plane of major disorder component of the heterocyclic ring with an interplanar angle between the two hydroxybenzyl groups of 66.00 (5)°. The phenyl substituent approaches a nearly perpendicular orientation relative to the mean plane of the imidazolidine ring, making a dihedral angle of 75.60 (12)°. This conformation is stabilized by two intramolecular $\text{O}-\text{H}\cdots\text{N}$ bonds, which generate $S(6)$ ring motifs.

Related literature

For related structures, see: Rivera *et al.* (2012a,b). For the synthesis of the precursor, see: Rivera *et al.* (2013). For bond-length data, see: Allen *et al.* (1987). For puckering parameters, see: Cremer & Pople (1975). For hydrogen-bond graph-set motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{24}\text{H}_{26}\text{N}_2\text{O}_2$
 $M_r = 374.5$
Monoclinic, $P2_1/c$
 $a = 16.8974$ (8) Å
 $b = 9.4893$ (5) Å
 $c = 12.5287$ (6) Å
 $\beta = 92.928$ (4)°
 $V = 2006.29$ (17) Å³
 $Z = 4$
Cu $K\alpha$ radiation
 $\mu = 0.62$ mm⁻¹
 $T = 120$ K
 $0.35 \times 0.25 \times 0.09$ mm

Data collection

Agilent Xcalibur (Atlas, Gemini ultra) diffractometer
Absorption correction: analytical (*CrysAlis PRO*; Agilent, 2010)
 $T_{\min} = 0.886$, $T_{\max} = 0.952$
8259 measured reflections
3491 independent reflections
2704 reflections with $I > 3\sigma(I)$
 $R_{\text{int}} = 0.03$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.113$
 $S = 1.51$
3491 reflections
276 parameters
5 restraints
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.25$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.18$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H101}\cdots\text{N9}$	0.98 (2)	1.88 (2)	2.7569 (18)	147.7 (19)
$\text{O19}-\text{H1019}\cdots\text{N11}$	1.00 (2)	1.79 (2)	2.709 (2)	152 (2)

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SUPERFLIP* (Palatinus & Chapuis 2007); program(s) used to refine structure: *JANA2006* (Petříček *et al.* 2006); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2005); software used to prepare material for publication: *JANA2006*.

We acknowledge the Dirección de Investigaciones, Sede Bogotá (DIB), and the Institutional research plan No. AVOZ10100521 of the Institute of Physics and the Praemium Academiae project of the Academy of Sciences (ASCR).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5339).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Brandenburg, K. & Putz, H. (2005). *DIAMOND*. Crystal Impact, Bonn, Germany.
Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
Palatinus, L. & Chapuis, G. (2007). *J. Appl. Cryst.* **40**, 786–790.
Petříček, V., Dusek, M. & Palatinus, L. (2006). *JANA2006*. Institute of Physics, Praha, Czech Republic.
Rivera, A., Cárdenas, L. & Ríos-Motta, J. (2013). *Curr. Org. Chem.* Accepted.
Rivera, A., Nerio, L. S., Ríos-Motta, J., Fejfarová, K. & Dušek, M. (2012a). *Acta Cryst. E* **68**, o170–o171.
Rivera, A., Pacheco, D., Ríos-Motta, J., Fejfarová, K. & Dusek, M. (2012b). *Tetrahedron Lett.* **53**, 6132–6135.

supporting information

Acta Cryst. (2013). E69, o1198 [doi:10.1107/S1600536813017893]

2,2'-[(4-Methyl-2-phenylimidazolidine-1,3-diyl)bis(methylene)]diphenol**Augusto Rivera, Lorena Cárdenas, Jaime Ríos-Motta, Václav Eigner and Michal Dušek****S1. Comment**

We have recently synthesized for the first time a series of tetrasubstituted imidazolidines by the reaction of *N,N'*-dibenzylpropane-1,2-diamine with substituted benzaldehydes, and we have provided evidence that they display intramolecular O—H \cdots N hydrogen bonds (Rivera *et al.*, 2013). In this article, we further advance these investigations by presenting the crystal structure of the title compound.

In the title compound, C₂₄H₂₆N₂O₂, the methyl-substituted imidazolidine ring exhibits molecular disorder over two orientations, with a refined occupancy ratio of 0.899 (4):0.101 (4) for atoms (C26, C27, C28):(C26', C27', C28'). For the major component, the imidazolidine ring (N9/C10/N11/C26/C27) adopts an envelope conformation with atom N11 as the flap: puckering parameters (Cremer & Pople, 1975) being Q₂ = 0.4429 (19) Å and Φ = 75.0 (2)°. The bond lengths (Allen *et al.*, 1987) and angles are close to standard values. Within the imidazolidine ring, bond distances and angles are comparable to those found in related structures (Rivera *et al.*, 2012*a,b*). The C2—O1 and C18—O19 bond lengths are 1.374 (2) and 1.369 (2) Å, respectively, which are comparable to other C—O bond lengths in di-Mannich bases reported by us (Rivera *et al.*, 2012*a,b*). The hydroxybenzyl groups makes an angle of 71.57 (15)° and 69.97 (19)° with the mean plane of heterocyclic ring defined by N9,C10,C26 and C27. With reference to this plane, the phenyl ring is almost perpendicular with a dihedral angle of 75.60 (12)°. The interplanar angle between the two hydroxybenzyl groups is 66.00 (5)°.

Single-crystal X-ray diffraction analysis reveals the existence of intramolecular O—H \cdots N hydrogen-bonding interactions between the two N atoms of the imidazolidine ring and the hydroxyl groups in the aromatic rings with S(6) set graph motifs (Bernstein *et al.* 1995) (Table 1). However, the two observed intramolecular hydrogen bond distances were different (Table 1). It is then surprising to observe the difference in O \cdots N distances between O1 \cdots N9 [O \cdots N = 2.757 (2) Å] and O19 \cdots N11 [O \cdots N = 2.709 (2) Å], which are longer than those observed in a related structure (Rivera *et al.*, 2012*a*). Because the length of hydrogen bonds depends on bond strength, these results indicated that the nitrogen atoms in the title compound are weaker hydrogen bond acceptors than the nitrogen in related structure (Rivera *et al.*, 2012*a*). This fact can be explained by the presence of the aryl group at the aminal carbon, where the imidazolidine ring adopts a conformation that causes an increase in unfavorable through-space interactions between the two nitrogen lone pairs. Thus, one of the possible consequences of these structural features is the reduction in electronic density around the nitrogen atoms.

S2. Experimental

For the originally reported synthesis, see: Rivera *et al.* (2013). The title compound (**I**) were recrystallized from ethanol.

S3. Refinement

All hydrogen atoms were discernible in difference Fourier maps and could be refined to reasonable geometry. According to common practice H atoms bonded to C were kept in ideal positions with $C-H = 0.96 \text{ \AA}$ while the coordinates of the H atoms bonded to O were refined freely. In both cases $U_{iso}(H)$ was set to $1.2U_{eq}(C,O)$. All non-hydrogen atoms were refined using harmonic refinement. The disordered part of molecule was refined with bond lengths kept equal for both groups. The resulting occupancy ratio was 0.899 (4):0.101 (4).




Figure 1

A perspective view of the title compound. Displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small spheres of arbitrary radii. Hydrogen bonds are drawn as dashed lines and only the major disorder component of the disordered section of the imidazolidine ring is shown.

2,2'-[(4-Methyl-2-phenylimidazolidine-1,3-diyl)bis(methylene)]diphenol

Crystal data

$C_{24}H_{26}N_2O_2$

$M_r = 374.5$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ycb

$a = 16.8974 (8) \text{ \AA}$

$b = 9.4893 (5) \text{ \AA}$

$c = 12.5287 (6) \text{ \AA}$

$\beta = 92.928 (4)^\circ$

$V = 2006.29 (17) \text{ \AA}^3$

$Z = 4$

$F(000) = 800$

$D_x = 1.239 \text{ Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.5418 \text{ \AA}$

Cell parameters from 3676 reflections

$\theta = 3.5\text{--}66.9^\circ$

$\mu = 0.62 \text{ mm}^{-1}$

$T = 120 \text{ K}$

Polygon shape, colorless

$0.35 \times 0.25 \times 0.09 \text{ mm}$

Data collection

Agilent Xcalibur (Atlas, Gemini ultra)
diffractometer

Radiation source: Enhance Ultra (Cu) X-ray
Source

Mirror monochromator

Detector resolution: $10.3784 \text{ pixels mm}^{-1}$

ω scans

Absorption correction: analytical
(*CrysAlis PRO*; Agilent, 2010)

$T_{min} = 0.886$, $T_{max} = 0.952$

8259 measured reflections

3491 independent reflections

2704 reflections with $I > 3\sigma(I)$

$R_{int} = 0.03$

$\theta_{\max} = 67.0^\circ$, $\theta_{\min} = 5.2^\circ$
 $h = -16 \rightarrow 19$

$k = -11 \rightarrow 11$
 $l = -14 \rightarrow 12$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.113$
 $S = 1.51$
 3491 reflections
 276 parameters
 5 restraints
 151 constraints

H atoms treated by a mixture of independent
 and constrained refinement
 Weighting scheme based on measured s.u.'s $w =$
 $1/(\sigma^2(I) + 0.0016I^2)$
 $(\Delta/\sigma)_{\max} = 0.049$
 $\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$
 Extinction correction: B-C type 1 Gaussian
 isotropic (Becker & Coppens, 1974)
 Extinction coefficient: 3200 (500)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.10968 (7)	0.77621 (13)	0.14452 (10)	0.0335 (4)	
O19	0.40788 (9)	1.04207 (16)	0.27300 (11)	0.0460 (5)	
N11	0.28125 (8)	0.99956 (14)	0.39327 (11)	0.0262 (4)	
N9	0.16113 (8)	0.91720 (14)	0.32755 (11)	0.0262 (4)	
C3	-0.01906 (11)	0.84850 (19)	0.08234 (15)	0.0323 (5)	
C17	0.49554 (10)	1.2217 (2)	0.33593 (15)	0.0335 (6)	
C22	0.33481 (10)	0.58293 (19)	0.28115 (14)	0.0303 (5)	
C2	0.03749 (10)	0.83400 (18)	0.16509 (14)	0.0282 (5)	
C18	0.43614 (10)	1.12529 (19)	0.35565 (13)	0.0301 (5)	
C21	0.29810 (10)	0.71116 (18)	0.29487 (13)	0.0277 (5)	
C4	-0.09159 (11)	0.91012 (19)	0.10146 (15)	0.0355 (6)	
C16	0.52487 (10)	1.3093 (2)	0.41629 (15)	0.0360 (6)	
C13	0.40514 (10)	1.11650 (18)	0.45702 (13)	0.0276 (5)	
C25	0.26334 (10)	0.63967 (18)	0.46963 (13)	0.0274 (5)	
C10	0.22524 (10)	0.88368 (17)	0.40711 (13)	0.0261 (5)	
C20	0.26239 (9)	0.74138 (17)	0.39014 (12)	0.0241 (5)	
C12	0.34266 (11)	1.00921 (19)	0.47984 (13)	0.0311 (5)	
C14	0.43528 (11)	1.2068 (2)	0.53588 (14)	0.0340 (6)	
C7	0.02229 (10)	0.87866 (17)	0.26891 (13)	0.0275 (5)	
C28	0.17720 (14)	1.1089 (2)	0.19419 (16)	0.0394 (7)	0.899 (4)
C23	0.33604 (10)	0.48228 (19)	0.36133 (14)	0.0315 (5)	
C24	0.29998 (10)	0.51088 (19)	0.45540 (14)	0.0310 (5)	
C8	0.08435 (10)	0.86041 (19)	0.35794 (13)	0.0311 (5)	
C6	-0.05061 (10)	0.94177 (19)	0.28506 (15)	0.0333 (6)	
C5	-0.10736 (11)	0.95685 (19)	0.20251 (15)	0.0357 (6)	
C27	0.16255 (12)	1.07345 (19)	0.30988 (16)	0.0280 (6)	0.899 (4)
C26	0.22944 (12)	1.1238 (2)	0.38684 (17)	0.0288 (6)	0.899 (4)
C26'	0.1443 (9)	1.0661 (5)	0.3488 (14)	0.0280 (6)	0.101 (4)
C27'	0.2300 (7)	1.1160 (12)	0.3475 (10)	0.0288 (6)	0.101 (4)
C15	0.49529 (11)	1.3021 (2)	0.51673 (15)	0.0383 (6)	
H1c3	-0.008331	0.816175	0.011928	0.0387*	
H1c17	0.516222	1.227294	0.266103	0.0402*	

H1c22	0.359666	0.563345	0.21556	0.0364*	
H1c21	0.29715	0.779845	0.238562	0.0333*	
H1c4	-0.130988	0.920237	0.044006	0.0426*	
H1c16	0.566013	1.375547	0.402306	0.0432*	
H1c25	0.238354	0.658669	0.535225	0.0329*	
H1c10	0.207032	0.876559	0.478241	0.0313*	
H1c12	0.367021	0.918641	0.490924	0.0373*	
H2c12	0.31869	1.033102	0.545288	0.0373*	
H1c14	0.414141	1.203482	0.60552	0.0408*	
H1c28	0.177434	1.20937	0.185208	0.0473*	0.899 (4)
H2c28	0.227476	1.071195	0.175972	0.0473*	0.899 (4)
H3c28	0.135948	1.068532	0.148357	0.0473*	0.899 (4)
H1c23	0.361769	0.393343	0.351577	0.0378*	
H1c24	0.300297	0.441351	0.511081	0.0373*	
H1c8	0.06796	0.908229	0.420618	0.0373*	
H2c8	0.090006	0.762114	0.374792	0.0373*	
H1c6	-0.061615	0.975461	0.35501	0.04*	
H1c5	-0.157476	0.9996	0.215351	0.0429*	
H1c27	0.113089	1.118473	0.322895	0.0336*	0.899 (4)
H1c26	0.256822	1.200587	0.355095	0.0346*	0.899 (4)
H2c26	0.208953	1.141634	0.455632	0.0346*	0.899 (4)
H1c15	0.516029	1.362475	0.572938	0.046*	
C28'	0.2525 (10)	1.162 (2)	0.2369 (11)	0.042 (6)	0.101 (4)
H1o1	0.1465 (13)	0.812 (2)	0.2010 (16)	0.0402*	
H1o19	0.3584 (15)	1.002 (2)	0.3010 (17)	0.0552*	
H1c26'	0.125922	1.075665	0.419702	0.0336*	0.101 (4)
H2c26'	0.114337	1.105645	0.288888	0.0336*	0.101 (4)
H1c27'	0.237554	1.19909	0.390453	0.0346*	0.101 (4)
H1c28'	0.252753	1.082083	0.190266	0.0509*	0.101 (4)
H2c28'	0.214686	1.230137	0.208928	0.0509*	0.101 (4)
H3c28'	0.304301	1.204201	0.241575	0.0509*	0.101 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0253 (6)	0.0372 (7)	0.0383 (7)	0.0031 (5)	0.0046 (5)	-0.0046 (5)
O19	0.0470 (8)	0.0509 (9)	0.0409 (8)	-0.0076 (7)	0.0101 (7)	-0.0169 (6)
N11	0.0290 (7)	0.0204 (7)	0.0290 (7)	-0.0013 (6)	-0.0014 (6)	0.0009 (6)
N9	0.0251 (7)	0.0219 (7)	0.0314 (7)	-0.0004 (6)	0.0008 (6)	0.0031 (6)
C3	0.0309 (9)	0.0308 (9)	0.0353 (9)	-0.0024 (7)	0.0035 (7)	0.0023 (7)
C17	0.0241 (8)	0.0393 (11)	0.0373 (10)	0.0043 (8)	0.0043 (7)	0.0022 (8)
C22	0.0260 (8)	0.0305 (9)	0.0348 (9)	-0.0043 (7)	0.0045 (7)	-0.0092 (7)
C2	0.0250 (8)	0.0234 (8)	0.0366 (9)	-0.0016 (7)	0.0048 (7)	0.0012 (7)
C18	0.0269 (8)	0.0295 (9)	0.0336 (9)	0.0071 (7)	-0.0008 (7)	-0.0054 (7)
C21	0.0282 (8)	0.0271 (9)	0.0278 (8)	-0.0047 (7)	0.0003 (7)	-0.0002 (7)
C4	0.0286 (9)	0.0318 (10)	0.0455 (11)	-0.0007 (8)	-0.0045 (8)	0.0070 (8)
C16	0.0252 (9)	0.0362 (10)	0.0459 (11)	-0.0027 (8)	-0.0037 (8)	0.0094 (9)
C13	0.0283 (8)	0.0252 (9)	0.0287 (8)	0.0028 (7)	-0.0034 (7)	0.0024 (7)

C25	0.0243 (8)	0.0265 (9)	0.0315 (9)	-0.0021 (7)	0.0024 (7)	0.0013 (7)
C10	0.0285 (8)	0.0250 (9)	0.0248 (8)	-0.0010 (7)	0.0022 (7)	0.0013 (7)
C20	0.0222 (8)	0.0228 (8)	0.0271 (8)	-0.0028 (7)	-0.0003 (6)	-0.0018 (6)
C12	0.0377 (10)	0.0288 (9)	0.0263 (8)	-0.0036 (8)	-0.0031 (7)	0.0019 (7)
C14	0.0382 (10)	0.0365 (10)	0.0269 (9)	-0.0061 (8)	-0.0034 (7)	0.0017 (7)
C7	0.0244 (8)	0.0245 (9)	0.0339 (9)	-0.0008 (7)	0.0042 (7)	0.0018 (7)
C28	0.0519 (14)	0.0285 (11)	0.0372 (11)	-0.0055 (10)	-0.0041 (10)	0.0045 (9)
C23	0.0228 (8)	0.0243 (9)	0.0473 (10)	-0.0008 (7)	0.0011 (7)	-0.0053 (8)
C24	0.0272 (8)	0.0257 (9)	0.0402 (10)	-0.0012 (7)	0.0014 (7)	0.0053 (7)
C8	0.0265 (9)	0.0350 (10)	0.0322 (9)	-0.0002 (8)	0.0066 (7)	0.0009 (7)
C6	0.0278 (9)	0.0306 (10)	0.0420 (10)	0.0023 (7)	0.0069 (8)	-0.0016 (8)
C5	0.0256 (9)	0.0319 (10)	0.0499 (11)	0.0037 (8)	0.0041 (8)	0.0004 (8)
C27	0.0277 (11)	0.0219 (9)	0.0344 (13)	0.0017 (8)	0.0008 (9)	0.0021 (8)
C26	0.0352 (10)	0.0208 (9)	0.0304 (13)	0.0011 (8)	0.0013 (9)	-0.0022 (8)
C26'	0.0277 (11)	0.0219 (9)	0.0344 (13)	0.0017 (8)	0.0008 (9)	0.0021 (8)
C27'	0.0352 (10)	0.0208 (9)	0.0304 (13)	0.0011 (8)	0.0013 (9)	-0.0022 (8)
C15	0.0405 (10)	0.0361 (11)	0.0372 (10)	-0.0099 (9)	-0.0084 (8)	0.0007 (8)
C28'	0.036 (10)	0.052 (13)	0.039 (11)	-0.021 (9)	-0.002 (8)	0.016 (9)

Geometric parameters (Å, °)

O1—C2	1.374 (2)	C10—H1c10	0.96
O19—C18	1.369 (2)	C12—H1c12	0.96
N11—C10	1.467 (2)	C12—H2c12	0.96
N11—C12	1.465 (2)	C14—C15	1.389 (3)
N11—C26	1.469 (2)	C14—H1c14	0.96
N11—C27'	1.500 (12)	C7—C8	1.501 (2)
N9—C10	1.469 (2)	C7—C6	1.394 (2)
N9—C8	1.473 (2)	C28—C27	1.521 (3)
N9—C27	1.500 (2)	C28—H1c28	0.96
N9—C26'	1.469 (6)	C28—H2c28	0.96
C3—C2	1.380 (2)	C28—H3c28	0.96
C3—C4	1.390 (3)	C23—C24	1.381 (3)
C3—H1c3	0.96	C23—H1c23	0.96
C17—C18	1.389 (2)	C24—H1c24	0.96
C17—C16	1.378 (3)	C8—H1c8	0.96
C17—H1c17	0.96	C8—H2c8	0.96
C22—C21	1.381 (2)	C6—C5	1.381 (3)
C22—C23	1.385 (3)	C6—H1c6	0.96
C22—H1c22	0.96	C5—H1c5	0.96
C2—C7	1.404 (2)	C27—C26	1.525 (3)
C18—C13	1.401 (2)	C27—H1c27	0.96
C21—C20	1.395 (2)	C26—H1c26	0.96
C21—H1c21	0.96	C26—H2c26	0.96
C4—C5	1.380 (3)	C26'—C27'	1.525 (19)
C4—H1c4	0.96	C26'—H1c26'	0.96
C16—C15	1.379 (3)	C26'—H2c26'	0.96
C16—H1c16	0.96	C27'—C28'	1.52 (2)

C13—C12	1.505 (2)	C27'—H1c27'	0.96
C13—C14	1.385 (2)	C15—H1c15	0.96
C25—C20	1.386 (2)	C28'—H1c28'	0.96
C25—C24	1.386 (2)	C28'—H2c28'	0.96
C25—H1c25	0.96	C28'—H3c28'	0.96
C10—C20	1.509 (2)		
C10—N11—C12	113.34 (13)	C2—C7—C6	117.97 (15)
C10—N11—C26	102.83 (13)	C8—C7—C6	122.06 (16)
C10—N11—C27'	103.5 (5)	C27—C28—H1c28	109.47
C12—N11—C26	112.84 (13)	C27—C28—H2c28	109.47
C12—N11—C27'	127.6 (5)	C27—C28—H3c28	109.47
C10—N9—C8	111.98 (13)	H1c28—C28—H2c28	109.47
C10—N9—C27	107.30 (13)	H1c28—C28—H3c28	109.47
C10—N9—C26'	103.2 (6)	H2c28—C28—H3c28	109.47
C8—N9—C27	114.89 (14)	C22—C23—C24	119.49 (16)
C8—N9—C26'	97.1 (6)	C22—C23—H1c23	120.25
C2—C3—C4	119.68 (17)	C24—C23—H1c23	120.25
C2—C3—H1c3	120.16	C25—C24—C23	120.18 (16)
C4—C3—H1c3	120.16	C25—C24—H1c24	119.91
C18—C17—C16	120.22 (17)	C23—C24—H1c24	119.91
C18—C17—H1c17	119.89	N9—C8—C7	110.91 (14)
C16—C17—H1c17	119.89	N9—C8—H1c8	109.47
C21—C22—C23	120.50 (16)	N9—C8—H2c8	109.47
C21—C22—H1c22	119.75	C7—C8—H1c8	109.47
C23—C22—H1c22	119.75	C7—C8—H2c8	109.47
O1—C2—C3	119.08 (15)	H1c8—C8—H2c8	108
O1—C2—C7	120.03 (14)	C7—C6—C5	121.38 (17)
C3—C2—C7	120.89 (16)	C7—C6—H1c6	119.31
O19—C18—C17	118.21 (16)	C5—C6—H1c6	119.31
O19—C18—C13	121.29 (15)	C4—C5—C6	119.63 (17)
C17—C18—C13	120.49 (16)	C4—C5—H1c5	120.19
C22—C21—C20	120.31 (15)	C6—C5—H1c5	120.19
C22—C21—H1c21	119.85	N9—C27—C28	111.35 (15)
C20—C21—H1c21	119.85	N9—C27—C26	103.51 (14)
C3—C4—C5	120.43 (17)	N9—C27—H1c27	113.22
C3—C4—H1c4	119.78	C28—C27—C26	112.46 (17)
C5—C4—H1c4	119.78	C28—C27—H1c27	104.42
C17—C16—C15	120.21 (17)	C26—C27—H1c27	112.13
C17—C16—H1c16	119.9	N11—C26—C27	101.74 (14)
C15—C16—H1c16	119.9	N11—C26—H1c26	109.47
C18—C13—C12	120.74 (15)	N11—C26—H2c26	109.47
C18—C13—C14	117.97 (16)	C27—C26—H1c26	109.47
C12—C13—C14	121.27 (15)	C27—C26—H2c26	109.47
C20—C25—C24	120.70 (16)	H1c26—C26—H2c26	116.23
C20—C25—H1c25	119.65	N9—C26'—H1c26'	109.47
C24—C25—H1c25	119.65	N9—C26'—H2c26'	109.47
N11—C10—N9	102.44 (12)	C27'—C26'—H1c26'	109.47

N11—C10—C20	112.26 (13)	C27'—C26'—H2c26'	109.47
N11—C10—H1c10	113.69	H1c26'—C26'—H2c26'	120.28
N9—C10—C20	113.32 (13)	N11—C27'—C26'	107.3 (8)
N9—C10—H1c10	112.64	N11—C27'—C28'	113.3 (11)
C20—C10—H1c10	102.9	N11—C27'—H1c27'	109.44
C21—C20—C25	118.82 (15)	C26'—C27'—C28'	112.5 (12)
C21—C20—C10	120.40 (14)	C26'—C27'—H1c27'	110.35
C25—C20—C10	120.77 (14)	C28'—C27'—H1c27'	103.96
N11—C12—C13	112.27 (13)	C16—C15—C14	119.45 (17)
N11—C12—H1c12	109.47	C16—C15—H1c15	120.27
N11—C12—H2c12	109.47	C14—C15—H1c15	120.27
C13—C12—H1c12	109.47	C27'—C28'—H1c28'	109.47
C13—C12—H2c12	109.47	C27'—C28'—H2c28'	109.47
H1c12—C12—H2c12	106.52	C27'—C28'—H3c28'	109.47
C13—C14—C15	121.65 (17)	H1c28'—C28'—H2c28'	109.47
C13—C14—H1c14	119.18	H1c28'—C28'—H3c28'	109.47
C15—C14—H1c14	119.17	H2c28'—C28'—H3c28'	109.47
C2—C7—C8	119.96 (15)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1o1...N9	0.98 (2)	1.88 (2)	2.7569 (18)	147.7 (19)
O19—H1o19...N11	1.00 (2)	1.79 (2)	2.709 (2)	152 (2)
O1—H1o1...C8	0.98 (2)	2.32 (2)	2.844 (2)	112.5 (16)
O19—H1o19...C12	1.00 (2)	2.27 (2)	2.884 (2)	118.6 (16)
O19—H1o19...C28'	1.00 (2)	2.45 (3)	2.877 (18)	105.1 (16)
C26'—H2c26'...C28	0.96	1.63	2.082 (17)	103.83
C28'—H3c28'...O19	0.96	2.35	2.877 (18)	114.12
C26—H1c26...C28'	0.96	1.52	1.973 (15)	102.84