

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(2E)-2-(1,3-Benzothiazol-2-yl)-3-(dimethylamino)prop-2-enitrileShaaban K. Mohamed,^{a,b} Mehmet Akkurt,^c Benson M. Kariuki,^d Ali M. Ali^e and Mustafa R. Albayati^{f*}

^aChemistry and Environmental Division, Manchester Metropolitan University, Manchester M1 5GD, England, ^bChemistry Department, Faculty of Science, Minia University, 61519 El-Minia, Egypt, ^cDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^dSchool of Chemistry, Cardiff University, Main Building, Park Place, Cardiff, CF10 3AT, Wales, ^eDepartment of Chemistry, Faculty of Science, Sohag University, 82524 Sohag, Egypt, and ^fKirkuk University, College of Science, Department of Chemistry, Kirkuk, Iraq
Correspondence e-mail: shaabankamel@yahoo.com


Received 7 December 2013; accepted 8 December 2013

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.037; wR factor = 0.107; data-to-parameter ratio = 14.9.

The molecular conformation of title compound, $\text{C}_{12}\text{H}_{11}\text{N}_3\text{S}$, is almost planar [maximum deviation = 0.063 (2) Å]; an intramolecular $\text{C}-\text{H}\cdots\text{N}$ hydrogen bond is noted. In the crystal, molecules interact with each other *via* $\pi-\pi$ stacking interactions between thiazole rings [centroid-centroid distance = 3.7475 (9) Å] and methyl- $\text{H}\cdots\pi(\text{C}_6)$ interactions, forming columns along the a axis.

Related literature

For various biological activities (*e.g.* anti-tumour, anti-inflammatory, anti-viral, *etc.*) of benzothiazole compounds, see: Selvam *et al.* (2011); Sanja & Cvetkovic (2011); Alang *et al.* (2010); Pal *et al.* (2011); Sharma *et al.* (2010); El-Shaer *et al.* (1997); Gupta & Raat (2010); Hutchinson *et al.* (2002); Gong *et al.* (2004); Hutchinson *et al.* (2003); Geronikaki & Theophilidis (1992); Vicini *et al.* (1990); Das *et al.* (2003); Klose *et al.* (1983); Satsangi *et al.* (1983).



Experimental

Crystal data

$\text{C}_{12}\text{H}_{11}\text{N}_3\text{S}$
 $M_r = 229.31$
Monoclinic, $P2_1/c$

$a = 7.3785$ (2) Å
 $b = 20.1801$ (4) Å
 $c = 8.2706$ (2) Å

$\beta = 112.947$ (4)°
 $V = 1134.03$ (6) Å³
 $Z = 4$
Cu $K\alpha$ radiation

$\mu = 2.32$ mm⁻¹
 $T = 293$ K
 $0.26 \times 0.20 \times 0.09$ mm

Data collection

Oxford Diffraction SuperNova
(Dual, Cu at zero, Atlas)
diffractometer
Absorption correction: multi-scan
(*CrysAlis PRO*; Oxford

Diffraction, 2013)
 $T_{\min} = 0.584$, $T_{\max} = 0.818$
4062 measured reflections
2195 independent reflections
1971 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.107$
 $S = 1.05$
2195 reflections

147 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.25$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C1–C6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C9–H9 \cdots N1	0.93	2.44	2.851 (2)	106
C10–H10A \cdots Cg1 ¹	0.96	2.77	3.549 (2)	138

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2013); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *PLATON* (Spek, 2009).

We thank Manchester Metropolitan University, Erciyes University and Cardiff University for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK5279).

References

- Alang, G., Kaur, G., Kaur, R., Singh, A. & Tiwari, R. (2010). *J. Young Pharmacists*, **2**, 394–398.
Das, J. J., Moquin, R. V., Liu, C., Doweiko, A. M., Defex, H. F., Fang, Q., Pang, S., Pitt, S., Shen, D. R., Schieven, G. L., Barrish, J. C. & Wityak, J. (2003). *Bioorg. Med. Chem. Lett.* **13**, 2587–2590.
El-Shaer, H. M., Abdel-Aziz, S. A., Allimony, H. A., Ali, U. F. & Abdel-Rahman, R. M. (1997). *Pharmazie*, **52**, 285–289.
Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
Geronikaki, A. & Theophilidis, G. (1992). *Eur. J. Med. Chem.* **27**, 709–716.
Gong, B., Hong, F., Kohm, C., Bonham, L. & Klin, P. (2004). *Bioorg. Med. Chem. Lett.* **14**, 1455–1459.
Gupta, A. & Raat, S. (2010). *J. Curr. Pharm. Res.* **3**, 13–23.
Hutchinson, I., Bradshaw, T. D., Matthew, C. S., Stevens, M. F. G. & Westwell, A. D. (2003). *Bioorg. Med. Chem. Lett.* **13**, 471–474.
Hutchinson, I., Jennings, S. A., Vishnuvajjala, B. R., Westwell, A. D. & Stevens, M. F. G. (2002). *J. Med. Chem.* **45**, 744–747.
Klose, N., Niedbolla, K., Schwartz, K. & Bottcher, I. (1983). *Arch. Pharm.* **316**, 941–951.
Oxford Diffraction (2013). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
Pal, N., Kumar, M. & Seth, G. (2011). *Eur. J. Chem.* **8**, 1174–1189.

- Sanja, O. & Cvetkovic, D. D. P.-K. (2011). *Chem. Ind. Chem. Eng. Q.* **17**, 9–15.
- Satsangi, R. K., Zaidi, S. M. & Misra, V. C. (1983). *Pharmazie*, **38**, 341–342.
- Selvam, T. P., Radhika, P., Janagaraj, S. & Kumar, A. S. (2011). *Res. Biotech.* **2**, 50–57.
- Sharma, N. K., Priyanka, M. S. & Jha, K. K. (2010). *J. Pharm. Res.* **3**, 969–970.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Vicini, P., Amoretti, L., Chiavarini, M. & Impicciatore, M. (1990). *Il Farmaco*, **45**, 933–943.

supporting information

Acta Cryst. (2014). E70, o52–o53 [https://doi.org/10.1107/S1600536813033266]

(2E)-2-(1,3-Benzothiazol-2-yl)-3-(dimethylamino)prop-2-enenitrile**Shaaban K. Mohamed, Mehmet Akkurt, Benson M. Kariuki, Ali M. Ali and Mustafa R. Albayati****S1. Comment**

It is well known that the thiazolyl group is of great importance in biological systems. In recent years, there has been considerable interest in the synthesis of substituted benzothiazolyl compounds due to their pharmacological properties such as anti-fungal (Selvam *et al.*, 2011; Sanja & Cvetkovic, 2011), anti-viral (Alang *et al.*, 2010; Pal *et al.*, 2011), anti-bacterial (Sharma *et al.*, 2010; El-Shaar *et al.*, 1997), analgesic (Gupta & Raat, 2010), anti-tumour (Hutchinson *et al.*, 2002) and anti-tuberculosis (Gong *et al.*, 2004; Hutchinson *et al.*, 2003) activities. Moreover, such compounds have been also found to have a potent local anaesthetic activity (Geronikaki & Theophilidis, 1992; Vicini *et al.*, 1990). Anti-inflammatory, analgesic, and anti-pyretic activities for some thiazolyl and benzothiazolyl derivatives are also known (Das *et al.*, 2003; Klose *et al.*, 1983; Satsangi *et al.*, 1983). Based on the above and following to our ongoing studies in synthesis of bio-active heterocyclic compounds the title compound has been prepared as a precursor for further study.

As shown in Fig. 1, title compound (I) has an almost planar conformation for non-hydrogen atoms with a maximum deviation of $-0.063(2)$ Å for C6.

The crystal structure is stabilized by π - π stacking interactions [$Cg1 \cdots Cg1(1-x, 1-y, 1-z) = 3.7475(9)$ Å; where Cg1 is a centroid of the five-membered (S1/N1/C1/C2/C7) thiazole ring of the 1,3-benzothiazole ring system] between the centroids of thiazole rings of the adjacent molecules. The crystal structure has no classical hydrogen bonds. Figs. 2 show the molecular packing of (I) viewed along the *a* direction.

S2. Experimental

A mixture of 1,3-benzothiazol-2-ylacetonitrile (174 mg, 1 mmol) and dimethylformamide-dimethylacetal (119 mg, 1 mmol) were taken in acetic acid (10 ml). The reaction mixture was refluxed and monitored by TLC until completion after 8 h. The reaction mixture was allowed to cool at ambient temperature and poured into ice water. The solid product was collected by filtration and recrystallized from ethanol to afford the title compound in 88% yield. Single crystals suitable for X-ray analysis were grown up on slow evaporation of ethanolic solution of the title compound at room temperature over three days. *M.pt.*: 441–443 K.

S3. Refinement

All H-atoms were refined using a riding model with C—H = 0.93 Å and $U_{iso} = 1.2U_{eq}(C)$ for aromatic-H atoms, and C—H = 0.96 Å and $U_{iso} = 1.5U_{eq}(C)$ for methyl-H atoms.




Figure 1

The structure of the title compound with the atom numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 50% probability level.




Figure 2

The molecular packing of the title compound viewing along the *a* axis. All H atoms are omitted for clarity.

(2E)-2-(1,3-Benzothiazol-2-yl)-3-(dimethylamino)prop-2-enitrile

Crystal data

$C_{12}H_{11}N_3S$

$M_r = 229.31$

Monoclinic, $P2_1/c$

Hall symbol: $-P 2_1/c$

$a = 7.3785$ (2) Å

$b = 20.1801$ (4) Å

$c = 8.2706$ (2) Å

$\beta = 112.947$ (4)°

$V = 1134.03$ (6) Å³

$Z = 4$

$F(000) = 480$

$D_x = 1.343$ Mg m⁻³

Cu $K\alpha$ radiation, $\lambda = 1.54184$ Å

Cell parameters from 1971 reflections

$\theta = 4.4\text{--}72.7^\circ$

$\mu = 2.32$ mm⁻¹

$T = 293$ K

Block, purple

$0.26 \times 0.20 \times 0.09$ mm

Data collection

Oxford Diffraction SuperNova (Dual, Cu at zero, Atlas) diffractometer
 Radiation source: SuperNova (Cu) X-ray Source
 Mirror monochromator
 ω scans
 Absorption correction: multi-scan (CrysAlis PRO; Oxford Diffraction, 2013)

$T_{\min} = 0.584$, $T_{\max} = 0.818$
 4062 measured reflections
 2195 independent reflections
 1971 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$
 $\theta_{\max} = 72.7^\circ$, $\theta_{\min} = 4.4^\circ$
 $h = -7 \rightarrow 9$
 $k = -16 \rightarrow 25$
 $l = -10 \rightarrow 8$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.107$
 $S = 1.05$
 2195 reflections
 147 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0678P)^2 + 0.1211P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.11365 (6)	0.50492 (2)	0.23607 (5)	0.0475 (1)
N1	0.24456 (18)	0.46002 (6)	0.55620 (16)	0.0427 (4)
N2	0.3152 (3)	0.66200 (8)	0.3024 (2)	0.0704 (6)
N3	0.5370 (2)	0.62896 (8)	0.81666 (18)	0.0544 (4)
C1	0.0527 (2)	0.42447 (7)	0.26748 (18)	0.0417 (4)
C2	0.1371 (2)	0.40931 (7)	0.44773 (19)	0.0403 (4)
C3	0.1078 (2)	0.34624 (8)	0.5022 (2)	0.0491 (5)
C4	-0.0040 (3)	0.30080 (8)	0.3794 (2)	0.0518 (5)
C5	-0.0886 (2)	0.31692 (9)	0.2020 (2)	0.0516 (5)
C6	-0.0608 (2)	0.37875 (9)	0.1440 (2)	0.0501 (5)
C7	0.2450 (2)	0.51233 (7)	0.46445 (18)	0.0388 (4)
C8	0.3437 (2)	0.57454 (7)	0.53589 (19)	0.0411 (4)
C9	0.4385 (2)	0.57944 (8)	0.7160 (2)	0.0459 (5)
C10	0.5596 (3)	0.69358 (10)	0.7510 (3)	0.0643 (6)
C11	0.6221 (4)	0.62098 (14)	1.0073 (3)	0.0817 (8)

C12	0.3318 (2)	0.62455 (8)	0.4120 (2)	0.0491 (5)
H3	0.16340	0.33490	0.62050	0.0590*
H4	-0.02320	0.25870	0.41570	0.0620*
H5	-0.16480	0.28570	0.12150	0.0620*
H6	-0.11670	0.38950	0.02530	0.0600*
H9	0.43210	0.54140	0.77700	0.0550*
H10A	0.62640	0.68910	0.67250	0.0960*
H10B	0.63510	0.72160	0.84750	0.0960*
H10C	0.43220	0.71290	0.68920	0.0960*
H11A	0.60670	0.57590	1.03660	0.1230*
H11B	0.55600	0.64990	1.05890	0.1230*
H11C	0.75950	0.63200	1.05180	0.1230*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0628 (3)	0.0411 (2)	0.0349 (2)	-0.0012 (2)	0.0150 (2)	0.0049 (1)
N1	0.0501 (7)	0.0382 (6)	0.0408 (6)	0.0023 (5)	0.0188 (5)	0.0008 (5)
N2	0.0975 (13)	0.0468 (8)	0.0584 (9)	-0.0053 (8)	0.0213 (8)	0.0088 (7)
N3	0.0544 (7)	0.0575 (8)	0.0464 (7)	-0.0001 (6)	0.0144 (6)	-0.0116 (6)
C1	0.0489 (7)	0.0408 (7)	0.0378 (7)	0.0023 (6)	0.0194 (6)	0.0037 (6)
C2	0.0457 (7)	0.0413 (7)	0.0365 (7)	0.0036 (6)	0.0189 (6)	0.0014 (5)
C3	0.0618 (9)	0.0453 (8)	0.0428 (8)	0.0007 (7)	0.0231 (7)	0.0065 (6)
C4	0.0642 (9)	0.0417 (8)	0.0567 (9)	-0.0033 (7)	0.0315 (7)	0.0019 (7)
C5	0.0575 (9)	0.0502 (8)	0.0518 (9)	-0.0093 (7)	0.0263 (7)	-0.0104 (7)
C6	0.0576 (9)	0.0546 (9)	0.0377 (7)	-0.0041 (7)	0.0180 (6)	-0.0027 (6)
C7	0.0443 (7)	0.0394 (7)	0.0336 (7)	0.0066 (5)	0.0162 (6)	0.0018 (5)
C8	0.0457 (7)	0.0369 (7)	0.0412 (7)	0.0041 (6)	0.0175 (6)	-0.0008 (6)
C9	0.0486 (8)	0.0454 (8)	0.0432 (8)	0.0039 (6)	0.0173 (6)	-0.0026 (6)
C10	0.0652 (10)	0.0537 (10)	0.0699 (11)	-0.0082 (8)	0.0219 (9)	-0.0178 (9)
C11	0.0848 (15)	0.0984 (17)	0.0464 (10)	-0.0053 (13)	0.0087 (9)	-0.0160 (11)
C12	0.0570 (9)	0.0379 (7)	0.0485 (8)	0.0002 (6)	0.0165 (7)	-0.0029 (6)

Geometric parameters (Å, °)

S1—C1	1.7311 (15)	C7—C8	1.456 (2)
S1—C7	1.7612 (14)	C8—C9	1.380 (2)
N1—C2	1.3878 (19)	C8—C12	1.416 (2)
N1—C7	1.3008 (19)	C3—H3	0.9300
N2—C12	1.150 (2)	C4—H4	0.9300
N3—C9	1.322 (2)	C5—H5	0.9300
N3—C10	1.447 (3)	C6—H6	0.9300
N3—C11	1.461 (3)	C9—H9	0.9300
C1—C2	1.407 (2)	C10—H10A	0.9600
C1—C6	1.389 (2)	C10—H10B	0.9600
C2—C3	1.395 (2)	C10—H10C	0.9600
C3—C4	1.378 (2)	C11—H11A	0.9600
C4—C5	1.391 (2)	C11—H11B	0.9600

C5—C6	1.381 (3)	C11—H11C	0.9600
C1—S1—C7	89.15 (7)	C2—C3—H3	120.00
C2—N1—C7	110.61 (12)	C4—C3—H3	120.00
C9—N3—C10	124.08 (15)	C3—C4—H4	120.00
C9—N3—C11	119.72 (17)	C5—C4—H4	119.00
C10—N3—C11	116.13 (18)	C4—C5—H5	120.00
S1—C1—C2	109.23 (10)	C6—C5—H5	120.00
S1—C1—C6	129.06 (12)	C1—C6—H6	121.00
C2—C1—C6	121.71 (14)	C5—C6—H6	121.00
N1—C2—C1	115.46 (13)	N3—C9—H9	114.00
N1—C2—C3	125.84 (13)	C8—C9—H9	115.00
C1—C2—C3	118.70 (13)	N3—C10—H10A	109.00
C2—C3—C4	119.52 (14)	N3—C10—H10B	109.00
C3—C4—C5	121.02 (16)	N3—C10—H10C	109.00
C4—C5—C6	120.77 (15)	H10A—C10—H10B	109.00
C1—C6—C5	118.27 (14)	H10A—C10—H10C	109.00
S1—C7—N1	115.55 (11)	H10B—C10—H10C	110.00
S1—C7—C8	119.17 (10)	N3—C11—H11A	109.00
N1—C7—C8	125.28 (13)	N3—C11—H11B	109.00
C7—C8—C9	117.55 (13)	N3—C11—H11C	109.00
C7—C8—C12	116.15 (13)	H11A—C11—H11B	109.00
C9—C8—C12	126.30 (14)	H11A—C11—H11C	110.00
N3—C9—C8	131.03 (15)	H11B—C11—H11C	109.00
N2—C12—C8	175.21 (17)		
C7—S1—C1—C2	-0.33 (12)	S1—C1—C2—N1	0.48 (18)
C7—S1—C1—C6	179.26 (16)	S1—C1—C2—C3	-179.33 (12)
C1—S1—C7—C8	-179.85 (13)	N1—C2—C3—C4	179.54 (17)
C1—S1—C7—N1	0.14 (13)	C1—C2—C3—C4	-0.7 (2)
C2—N1—C7—C8	-179.91 (15)	C2—C3—C4—C5	-0.2 (3)
C7—N1—C2—C3	179.41 (16)	C3—C4—C5—C6	0.8 (3)
C2—N1—C7—S1	0.10 (19)	C4—C5—C6—C1	-0.4 (3)
C7—N1—C2—C1	-0.4 (2)	S1—C7—C8—C12	-2.3 (2)
C11—N3—C9—C8	179.2 (2)	N1—C7—C8—C9	-2.6 (2)
C10—N3—C9—C8	2.5 (3)	S1—C7—C8—C9	177.44 (12)
C6—C1—C2—C3	1.1 (2)	N1—C7—C8—C12	177.71 (15)
C6—C1—C2—N1	-179.15 (15)	C12—C8—C9—N3	0.5 (3)
S1—C1—C6—C5	179.96 (13)	C7—C8—C9—N3	-179.24 (17)
C2—C1—C6—C5	-0.5 (2)		

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C1—C6 ring.

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C9—H9···N1	0.93	2.44	2.851 (2)	106

C10—H10A \cdots Cg1 ⁱ	0.96	2.77	3.549 (2)	138
------------------------------------	------	------	-----------	-----

Symmetry code: (i) $-x+1, -y+1, -z+1$.