

Crystal structure of 3-[4-(pyrimidin-2-yl)-piperazin-1-ium-1-yl]butanoate

Thammarse S. Yamuna,^a Jerry P. Jasinski,^{b*} Manpreet Kaur,^a Brian J. Anderson^b and H.S. Yathirajan^a^aDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, and ^bDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA. *Correspondence e-mail: jjasinski@keene.edu

Received 15 August 2014; accepted 21 August 2014

Edited by M. Bolte, Goethe-Universität Frankfurt Germany


The title compound, C₁₂H₁₈N₄O₂, crystallizes in the zwitterionic form with protonation at the N atom of the piperazine ring bearing the carboxylate group. The piperazine ring adopts a slightly distorted chair conformation. In the crystal, N—H...O hydrogen bonds are observed, forming chains along [010]. The packing is consolidated by C—H...O interactions, which generate a three-dimensional network.

Keywords: crystal structure; 3-(piperazin-1-ium-1-yl)butanoate; zwitterionic form; fused heterocyclic derivatives; aza-Michael reactions.

CCDC reference: 1020635

1. Related literature

For general background and pharmacological properties of fused heterocyclic derivatives, see: Amin *et al.* (2009); Ibrahim & El-Metwally (2010); Kuyper *et al.* (1996); Onal & Yildirim (2007); Padmaja *et al.* (2009); Tollefson *et al.* (1991). For pharmacological properties of pyrimidines, see: Burdge (2000). For background to aza-Michael reactions, see: Arend *et al.* (1998); Vicario *et al.* (2005); Xu & Xia (2005). For related structures, see: Jin *et al.* (2012); Parvez *et al.* (2004); Yamuna *et al.* (2014a,b).



2. Experimental

2.1. Crystal data

C₁₂H₁₈N₄O₂M_r = 250.30

Monoclinic, *P*2₁/*c*
a = 13.5157 (6) Å
b = 7.8454 (3) Å
c = 12.2147 (5) Å
 β = 106.884 (5)°
V = 1239.36 (9) Å³

Z = 4
 Cu *K*α radiation
 μ = 0.77 mm⁻¹
T = 173 K
 0.32 × 0.22 × 0.06 mm

2.2. Data collection

Agilent Eos Gemini diffractometer
 Absorption correction: multi-scan
 (*CrysAlis PRO* and *CrysAlis RED*; Agilent, 2012)
*T*_{min} = 0.854, *T*_{max} = 1.000

3995 measured reflections
 3995 independent reflections
 3668 reflections with *I* > 2σ(*I*)

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.149$
S = 1.06
 3995 reflections
 169 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.36 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.27 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, °).

| <i>D</i> —H... <i>A</i> | <i>D</i> —H | H... <i>A</i> | <i>D</i> ... <i>A</i> | <i>D</i> —H... <i>A</i> |
|-----------------------------|-------------|---------------|-----------------------|-------------------------|
| N1—H1...O2 ⁱ | 1.00 | 1.67 | 2.653 (2) | 168 |
| C2—H2B...O2 ⁱ | 0.99 | 2.51 | 3.277 (3) | 134 |
| C4—H4B...O1 ⁱⁱ | 0.99 | 2.58 | 3.428 (3) | 144 |
| C7—H7B...O1 ⁱ | 0.99 | 2.53 | 3.147 (3) | 120 |
| C11—H11...O1 ⁱⁱⁱ | 0.95 | 2.47 | 3.352 (3) | 155 |

Symmetry codes: (i) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x, -y + 1, -z + 1$; (iii) $x + 1, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED*; program(s) used to solve structure: *SUPERFLIP* (Palatinus & Chapuis, 2007); program(s) used to refine structure: *SHELXL2012* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2*.

Acknowledgements

TSY thanks University of Mysore for research facilities and also grateful to the Principal, Maharani's Science College for Women, Mysore, for giving permission to do research. JPI acknowledges the NSF-MRI program (grant No. CHE-1039027) for funds to purchase the X-ray diffractometer.

Supporting information for this paper is available from the IUCr electronic archives (Reference: BT6993).

References

- Agilent (2012). *CrysAlis PRO* and *CrysAlis RED*. Agilent Technologies, Yarnton, Oxfordshire, England.
 Amin, K. M., Hanna, M. M., Abo-Youssef, H. E., Riham, F. & George, R. F. (2009). *Eur. J. Med. Chem.*, **44**, 4572–4584.
 Arend, M., Westermann, B. & Risch, N. (1998). *Angew. Chem. Int. Ed.* **37**, 1045–1070.
 Burdge, E. L. (2000). *Pest Manag. Sci.* **56**, 245–248.
 Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
 Ibrahim, D. A. & El-Metwally, A. M. (2010). *Eur. J. Med. Chem.*, **45**, 1158–1166.

- Jin, S., Huang, Y., Fang, H., Wang, T. & Ding, L. (2012). *Acta Cryst.* **E68**, o2827.
- Kuyper, L. F., Garvey, J. M., Baccanari, D. P., Champness, J. N., Stammers, D. K. & Beddell, C. R. (1996). *Bioorg. Med. Chem. Lett.*, **4**, 593–602.
- Onal, Z. & Yildirim, I. (2007). *Heterocycl. Commun.* **13**, 113–120.
- Padmaja, A., Payani, T., Reddy, G. D., Dinneswara Reddy, G. & Padmavathi, V. (2009). *Eur. J. Med. Chem.* **44**, 4557–4566.
- Palatinus, L. & Chapuis, G. (2007). *J. Appl. Cryst.* **40**, 786–790.
- Parvez, M., Arayne, S., Sultana, N. & Siddiqi, A. Z. (2004). *Acta Cryst.* **C60**, o281–o283.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tollefson, G. D., Lancaster, S. P. & Montague-Clouse, J. (1991). *Psychopharmacol. Bull.*, **27**, 163–170.
- Vicario, J. L., Badia, D. & Carrillo, L. (2005). *Org. Prep. Proced. Int.* **37**, 513–538.
- Xu, L.-W. & Xia, C.-G. (2005). *Eur. J. Org. Chem.* pp. 633–639.
- Yamuna, T. S., Kaur, M., Anderson, B. J., Jasinski, J. P. & Yathirajan, H. S. (2014b). *Acta Cryst.* **E70**, o206–o207.
- Yamuna, T. S., Kaur, M., Jasinski, J. P. & Yathirajan, H. S. (2014a). *Acta Cryst.* **E70**, o702–o703.

supporting information

Acta Cryst. (2014). E70, o1063–o1064 [doi:10.1107/S1600536814018972]

Crystal structure of 3-[4-(pyrimidin-2-yl)piperazin-1-ium-1-yl]butanoate

Thammarse S. Yamuna, Jerry P. Jasinski, Manpreet Kaur, Brian J. Anderson and H.S. Yathirajan

S1. Comment

Pyrimidines in general have been of much interest for biological and medical reasons and thus their chemistry has been investigated extensively (Onal & Yildirim, 2007) and in many drugs used for the treatment of hypothyroidism and hypertension, in cancer chemotherapy or HIV infections (Burdge, 2000) and with related fused heterocyclic compounds that exhibit biological activities such as anticancer (Amin *et al.*, 2009), antiviral (Ibrahim & El-Metwally, 2010), antibacterial (Kuyper *et al.*, 1996) and antioxidant (Padmaja *et al.*, 2009). Some pyrimidinylpiperazinyl compounds like buspirone and BuSpar (Tollefson *et al.*, 1991) are used to treat anxiety. The incorporation of two moieties increases biological activity of both the molecules. Aza-Michael addition reaction has been extensively studied using a variety of catalysts as well as solvents and various researchers have also reported the utility of aza-Michael addition towards the synthesis of various pharmacological active compounds and proved useful in the synthesis of core intermediates of many natural products (Arend *et al.*, 1998). The role of aza-Michael reaction in the synthesis of pharmacologically important families of β -amino carbonyl compounds and its derivatives is well documented in the literature (Vicario *et al.*, 2005; Xu & Xia, 2005). Our research group has published many papers on incorporated heterocyclic ring structures, viz; 4-(pyrimidin-2-yl)piperazin-1-ium (E)-3-carboxyprop-2-enoate (Yamuna *et al.*, 2014a); flupentixol tartarate (Yamuna *et al.*, 2014b). Some related zwitterion structures are: 3,3'-(piperazine-1,4-diium-1,4-diyl)dipropionate dihydrate (Jin *et al.*, 2012), enoxacin trihydrate[1-ethyl-6-fluoro-1,4-dihydro-4-oxo-7-(piperazin-4-ium-1-yl)-1,8-naphthyridine-3-carboxylate trihydrate] (Parvez *et al.*, 2004). In view of the importance of derivatives of the incorporated heterocyclic pyrimidinylpiperazines, this paper reports the crystal structure of the title zwitterionic compound, (I), 3-(4-Pyrimidin-2-ylpiperazin-1-ium-1-yl)-butanoate, $C_{12}H_{18}N_4O_2$ prepared from 2-(piperazin-1-yl)pyrimidine and but-2-enoic acid by aza-Michael addition reaction.

The title compound, (I), $C_{12}H_{18}N_4O_2$ crystallizes in the zwitterionic form with protonation on the N1 nitrogen atom of the piperazine ring (Fig. 1). In the compound, the piperazine ring adopts a slightly disordered chair conformation (puckering parameters Q , θ , and $\varphi = 0.576$ (2) Å, 3.0 (2)° and 282 (4)°, respectively. Bond lengths are in normal ranges. In the crystal, N—H \cdots O intermolecular hydrogen bonds are observed forming 1D chains along [0 1 0] (Fig. 2). The packing is consolidated by weak C—H \cdots O interactions which generate a three-dimensional network.

S2. Experimental

A mixture of 1-(2-pyrimidinyl)piperazine, from sigma-aldrich (0.2 g, 1.22 mmol) and crotonic acid (but-2-enoic acid) (0.1048 g, 1.22 mmol) were dissolved in DMSO, stirred well and warmed at 343 K for 20 minutes. After few days, X-ray quality crystals were obtained on slow evaporation (m.p.: 411–418 K).

S3. Refinement

H3 was refined isotropically and all of the remaining H atoms were placed in their calculated positions and then refined using the riding model with Atom—H lengths of 0.95 Å (CH); 0.99 Å (CH₂); 0.98 Å (CH₃) or 1.00 Å (NH). Isotropic displacement parameters for these atoms were set to 1.2 (CH, CH₂, NH) or 1.5 (CH₃) times U_{eq} of the parent atom. Idealised Me was refined as a rotating group. The title compound was refined as a twin with BASF = 0.41572.





Figure 1

ORTEP drawing of (C₁₂H₁₈N₄O₂) showing the labeling scheme of the asymmetric unit of the title compound with 30% probability displacement ellipsoids.

**Figure 2**

Molecular packing for (I), viewed along the *b* axis. Dashed lines indicate weak C—H···O intermolecular interactions in addition to N—H···O intermolecular hydrogen bonds which together form an extended three-dimensional supramolecular network structure. H atoms not involved in hydrogen bonding have been removed for clarity.

3-[4-(Pyrimidin-2-yl)piperazin-1-ium-1-yl]butanoate

Crystal data

$C_{12}H_{18}N_4O_2$
 $M_r = 250.30$
 Monoclinic, $P2_1/c$
 $a = 13.5157(6) \text{ \AA}$
 $b = 7.8454(3) \text{ \AA}$
 $c = 12.2147(5) \text{ \AA}$
 $\beta = 106.884(5)^\circ$
 $V = 1239.36(9) \text{ \AA}^3$
 $Z = 4$

$F(000) = 536$
 $D_x = 1.341 \text{ Mg m}^{-3}$
 Cu $K\alpha$ radiation, $\lambda = 1.54184 \text{ \AA}$
 Cell parameters from 5404 reflections
 $\theta = 3.8\text{--}71.4^\circ$
 $\mu = 0.77 \text{ mm}^{-1}$
 $T = 173 \text{ K}$
 Irregular, colourless
 $0.32 \times 0.22 \times 0.06 \text{ mm}$

Data collection

Agilent Eos Gemini
 diffractometer
 Radiation source: Enhance (Cu) X-ray Source
 Detector resolution: $16.0416 \text{ pixels mm}^{-1}$
 ω scans
 Absorption correction: multi-scan
 (*CrysAlis PRO* and *CrysAlis RED*; Agilent,
 2012)

$T_{\min} = 0.854$, $T_{\max} = 1.000$
 3995 measured reflections
 3995 independent reflections
 3668 reflections with $I > 2\sigma(I)$
 $\theta_{\max} = 71.2^\circ$, $\theta_{\min} = 3.4^\circ$
 $h = -16 \rightarrow 16$
 $k = -9 \rightarrow 9$
 $l = -13 \rightarrow 14$

*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.053$ $wR(F^2) = 0.149$ $S = 1.06$

3995 reflections

169 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0768P)^2 + 0.6386P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.36 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refined as a 2-component twin.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|---------------|------------|--------------|----------------------------------|
| O1 | -0.15848 (13) | 0.4680 (3) | 0.29107 (16) | 0.0449 (5) |
| O2 | -0.05978 (12) | 0.4109 (2) | 0.17822 (15) | 0.0339 (4) |
| N1 | 0.16438 (12) | 0.1694 (2) | 0.44116 (14) | 0.0217 (4) |
| H1 | 0.1313 | 0.0618 | 0.4040 | 0.026* |
| N2 | 0.32869 (12) | 0.0043 (3) | 0.60849 (16) | 0.0287 (4) |
| N3 | 0.40034 (14) | 0.0836 (3) | 0.79721 (17) | 0.0324 (5) |
| N4 | 0.50457 (13) | 0.0549 (3) | 0.66957 (17) | 0.0336 (5) |
| C1 | -0.08258 (15) | 0.4025 (3) | 0.2726 (2) | 0.0277 (5) |
| C2 | -0.00940 (16) | 0.2979 (3) | 0.3691 (2) | 0.0308 (5) |
| H2A | -0.0174 | 0.3357 | 0.4434 | 0.037* |
| H2B | -0.0296 | 0.1763 | 0.3588 | 0.037* |
| C3 | 0.10322 (15) | 0.3150 (3) | 0.3722 (2) | 0.0260 (5) |
| H3 | 0.098 (2) | 0.289 (4) | 0.289 (3) | 0.041 (8)* |
| C4 | 0.16425 (15) | 0.1604 (3) | 0.56333 (18) | 0.0261 (5) |
| H4A | 0.0922 | 0.1532 | 0.5671 | 0.031* |
| H4B | 0.1959 | 0.2650 | 0.6040 | 0.031* |
| C5 | 0.22479 (15) | 0.0048 (3) | 0.6211 (2) | 0.0290 (5) |
| H5A | 0.2291 | 0.0053 | 0.7034 | 0.035* |
| H5B | 0.1880 | -0.1001 | 0.5866 | 0.035* |
| C6 | 0.32772 (15) | 0.0092 (3) | 0.48941 (19) | 0.0276 (5) |
| H6A | 0.2918 | -0.0928 | 0.4491 | 0.033* |
| H6B | 0.3995 | 0.0084 | 0.4845 | 0.033* |
| C7 | 0.27285 (14) | 0.1685 (3) | 0.43312 (18) | 0.0243 (4) |
| H7A | 0.3102 | 0.2707 | 0.4716 | 0.029* |
| H7B | 0.2719 | 0.1716 | 0.3518 | 0.029* |
| C8 | 0.41389 (15) | 0.0508 (3) | 0.69526 (18) | 0.0248 (4) |
| C9 | 0.48642 (19) | 0.1182 (3) | 0.8810 (2) | 0.0359 (5) |
| H9 | 0.4802 | 0.1456 | 0.9545 | 0.043* |

| | | | | |
|------|--------------|------------|------------|------------|
| C10 | 0.58388 (18) | 0.1162 (3) | 0.8665 (2) | 0.0372 (6) |
| H10 | 0.6444 | 0.1349 | 0.9284 | 0.045* |
| C11 | 0.58837 (17) | 0.0857 (3) | 0.7574 (2) | 0.0376 (6) |
| H11 | 0.6540 | 0.0864 | 0.7436 | 0.045* |
| C12 | 0.14735 (18) | 0.4895 (3) | 0.4145 (2) | 0.0329 (5) |
| H12A | 0.1545 | 0.4997 | 0.4964 | 0.049* |
| H12B | 0.1007 | 0.5784 | 0.3725 | 0.049* |
| H12C | 0.2153 | 0.5026 | 0.4019 | 0.049* |

Atomic displacement parameters (Å²)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|-------------|-------------|-------------|--------------|--------------|
| O1 | 0.0322 (9) | 0.0605 (12) | 0.0415 (11) | 0.0190 (8) | 0.0099 (8) | 0.0110 (9) |
| O2 | 0.0282 (8) | 0.0318 (8) | 0.0409 (10) | 0.0066 (6) | 0.0087 (7) | 0.0019 (7) |
| N1 | 0.0143 (7) | 0.0280 (9) | 0.0214 (9) | 0.0003 (6) | 0.0028 (6) | 0.0038 (7) |
| N2 | 0.0138 (8) | 0.0488 (11) | 0.0234 (9) | 0.0065 (7) | 0.0053 (7) | 0.0076 (8) |
| N3 | 0.0284 (9) | 0.0427 (11) | 0.0266 (10) | 0.0100 (8) | 0.0088 (8) | 0.0026 (8) |
| N4 | 0.0184 (8) | 0.0522 (12) | 0.0303 (10) | 0.0007 (8) | 0.0070 (8) | −0.0065 (9) |
| C1 | 0.0200 (9) | 0.0303 (11) | 0.0295 (11) | −0.0011 (8) | 0.0018 (8) | 0.0025 (9) |
| C2 | 0.0222 (10) | 0.0364 (12) | 0.0335 (12) | 0.0054 (9) | 0.0078 (9) | 0.0073 (10) |
| C3 | 0.0200 (9) | 0.0287 (10) | 0.0259 (10) | 0.0018 (8) | 0.0014 (8) | 0.0052 (9) |
| C4 | 0.0166 (9) | 0.0379 (12) | 0.0245 (10) | 0.0041 (8) | 0.0072 (8) | 0.0033 (9) |
| C5 | 0.0158 (9) | 0.0441 (13) | 0.0282 (11) | 0.0042 (8) | 0.0084 (8) | 0.0112 (10) |
| C6 | 0.0169 (9) | 0.0424 (13) | 0.0233 (10) | 0.0046 (8) | 0.0059 (8) | 0.0012 (9) |
| C7 | 0.0154 (9) | 0.0373 (11) | 0.0204 (10) | −0.0038 (8) | 0.0055 (7) | 0.0012 (9) |
| C8 | 0.0184 (9) | 0.0304 (10) | 0.0254 (11) | 0.0077 (8) | 0.0062 (8) | 0.0044 (9) |
| C9 | 0.0359 (12) | 0.0428 (13) | 0.0264 (12) | 0.0102 (10) | 0.0051 (10) | −0.0020 (10) |
| C10 | 0.0297 (11) | 0.0408 (13) | 0.0336 (13) | 0.0035 (10) | −0.0027 (10) | −0.0052 (10) |
| C11 | 0.0199 (10) | 0.0495 (14) | 0.0407 (14) | 0.0000 (10) | 0.0046 (9) | −0.0103 (11) |
| C12 | 0.0322 (11) | 0.0287 (11) | 0.0337 (13) | −0.0001 (9) | 0.0032 (10) | 0.0024 (10) |

Geometric parameters (Å, °)

| | | | |
|--------|-----------|---------|-----------|
| O1—C1 | 1.226 (3) | C4—H4A | 0.9900 |
| O2—C1 | 1.278 (3) | C4—H4B | 0.9900 |
| N1—H1 | 1.0000 | C4—C5 | 1.524 (3) |
| N1—C3 | 1.514 (2) | C5—H5A | 0.9900 |
| N1—C4 | 1.494 (3) | C5—H5B | 0.9900 |
| N1—C7 | 1.498 (2) | C6—H6A | 0.9900 |
| N2—C5 | 1.457 (2) | C6—H6B | 0.9900 |
| N2—C6 | 1.451 (3) | C6—C7 | 1.513 (3) |
| N2—C8 | 1.369 (3) | C7—H7A | 0.9900 |
| N3—C8 | 1.335 (3) | C7—H7B | 0.9900 |
| N3—C9 | 1.335 (3) | C9—H9 | 0.9500 |
| N4—C8 | 1.351 (3) | C9—C10 | 1.379 (3) |
| N4—C11 | 1.336 (3) | C10—H10 | 0.9500 |
| C1—C2 | 1.538 (3) | C10—C11 | 1.373 (4) |
| C2—H2A | 0.9900 | C11—H11 | 0.9500 |

| | | | |
|---------------|--------------|---------------|-------------|
| C2—H2B | 0.9900 | C12—H12A | 0.9800 |
| C2—C3 | 1.518 (3) | C12—H12B | 0.9800 |
| C3—H3 | 1.01 (3) | C12—H12C | 0.9800 |
| C3—C12 | 1.523 (3) | | |
| C3—N1—H1 | 106.5 | N2—C5—H5B | 109.4 |
| C4—N1—H1 | 106.5 | C4—C5—H5A | 109.4 |
| C4—N1—C3 | 115.63 (17) | C4—C5—H5B | 109.4 |
| C4—N1—C7 | 110.51 (15) | H5A—C5—H5B | 108.0 |
| C7—N1—H1 | 106.5 | N2—C6—H6A | 109.7 |
| C7—N1—C3 | 110.70 (15) | N2—C6—H6B | 109.7 |
| C6—N2—C5 | 112.18 (17) | N2—C6—C7 | 109.75 (18) |
| C8—N2—C5 | 122.52 (19) | H6A—C6—H6B | 108.2 |
| C8—N2—C6 | 121.99 (18) | C7—C6—H6A | 109.7 |
| C8—N3—C9 | 115.4 (2) | C7—C6—H6B | 109.7 |
| C11—N4—C8 | 115.6 (2) | N1—C7—C6 | 109.51 (16) |
| O1—C1—O2 | 125.4 (2) | N1—C7—H7A | 109.8 |
| O1—C1—C2 | 118.0 (2) | N1—C7—H7B | 109.8 |
| O2—C1—C2 | 116.65 (19) | C6—C7—H7A | 109.8 |
| C1—C2—H2A | 109.0 | C6—C7—H7B | 109.8 |
| C1—C2—H2B | 109.0 | H7A—C7—H7B | 108.2 |
| H2A—C2—H2B | 107.8 | N3—C8—N2 | 117.45 (18) |
| C3—C2—C1 | 112.91 (18) | N3—C8—N4 | 126.3 (2) |
| C3—C2—H2A | 109.0 | N4—C8—N2 | 116.26 (19) |
| C3—C2—H2B | 109.0 | N3—C9—H9 | 118.3 |
| N1—C3—C2 | 109.19 (17) | N3—C9—C10 | 123.5 (2) |
| N1—C3—H3 | 105.9 (16) | C10—C9—H9 | 118.3 |
| N1—C3—C12 | 113.11 (17) | C9—C10—H10 | 122.0 |
| C2—C3—H3 | 100.5 (16) | C11—C10—C9 | 116.1 (2) |
| C2—C3—C12 | 112.28 (19) | C11—C10—H10 | 122.0 |
| C12—C3—H3 | 115.0 (17) | N4—C11—C10 | 123.0 (2) |
| N1—C4—H4A | 109.6 | N4—C11—H11 | 118.5 |
| N1—C4—H4B | 109.6 | C10—C11—H11 | 118.5 |
| N1—C4—C5 | 110.12 (17) | C3—C12—H12A | 109.5 |
| H4A—C4—H4B | 108.1 | C3—C12—H12B | 109.5 |
| C5—C4—H4A | 109.6 | C3—C12—H12C | 109.5 |
| C5—C4—H4B | 109.6 | H12A—C12—H12B | 109.5 |
| N2—C5—C4 | 110.96 (17) | H12A—C12—H12C | 109.5 |
| N2—C5—H5A | 109.4 | H12B—C12—H12C | 109.5 |
| O1—C1—C2—C3 | -144.6 (2) | C6—N2—C5—C4 | -57.3 (3) |
| O2—C1—C2—C3 | 36.8 (3) | C6—N2—C8—N3 | 163.5 (2) |
| N1—C4—C5—N2 | 54.5 (2) | C6—N2—C8—N4 | -18.1 (3) |
| N2—C6—C7—N1 | -59.2 (2) | C7—N1—C3—C2 | 172.41 (18) |
| N3—C9—C10—C11 | 3.7 (4) | C7—N1—C3—C12 | -61.8 (2) |
| C1—C2—C3—N1 | -161.91 (18) | C7—N1—C4—C5 | -55.5 (2) |
| C1—C2—C3—C12 | 71.8 (3) | C8—N2—C5—C4 | 102.5 (2) |
| C3—N1—C4—C5 | 177.77 (16) | C8—N2—C6—C7 | -100.4 (2) |

| | | | |
|--------------|--------------|---------------|------------|
| C3—N1—C7—C6 | -172.54 (17) | C8—N3—C9—C10 | -1.8 (4) |
| C4—N1—C3—C2 | -61.0 (2) | C8—N4—C11—C10 | -1.8 (4) |
| C4—N1—C3—C12 | 64.8 (2) | C9—N3—C8—N2 | 176.1 (2) |
| C4—N1—C7—C6 | 58.0 (2) | C9—N3—C8—N4 | -2.2 (3) |
| C5—N2—C6—C7 | 59.5 (2) | C9—C10—C11—N4 | -1.7 (4) |
| C5—N2—C8—N3 | 5.7 (3) | C11—N4—C8—N2 | -174.3 (2) |
| C5—N2—C8—N4 | -175.9 (2) | C11—N4—C8—N3 | 4.0 (4) |

Hydrogen-bond geometry (Å, °)

| <i>D—H...A</i> | <i>D—H</i> | <i>H...A</i> | <i>D...A</i> | <i>D—H...A</i> |
|-----------------------------|------------|--------------|--------------|----------------|
| N1—H1...O2 ⁱ | 1.00 | 1.67 | 2.653 (2) | 168 |
| C2—H2B...O2 ⁱ | 0.99 | 2.51 | 3.277 (3) | 134 |
| C4—H4B...O1 ⁱⁱ | 0.99 | 2.58 | 3.428 (3) | 144 |
| C7—H7B...O1 ⁱ | 0.99 | 2.53 | 3.147 (3) | 120 |
| C11—H11...O1 ⁱⁱⁱ | 0.95 | 2.47 | 3.352 (3) | 155 |

Symmetry codes: (i) $-x, y-1/2, -z+1/2$; (ii) $-x, -y+1, -z+1$; (iii) $x+1, -y+1/2, z+1/2$.