

# Crystal structure of 4-methyl-2-oxo-2H-chromen-7-yl ferrocenecarboxylate

Juan Yu, Lei-Lei Gao, Peng Huang\* and Dian-Lei Wang\*

School of Pharmacy, Anhui University of Chinese Medicine, Hefei 230038, People's Republic of China. \*Correspondence e-mail: great7701@126.com, wdl8105@126.com

Received 24 September 2014; accepted 7 October 2014

Edited by M. Weil, Vienna University of Technology, Austria


The title molecule,  $[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_{16}\text{H}_{11}\text{O}_4)]$ , consists of a ferrocenyl moiety and a 4-methylcoumarin group linked through an ester unit to one of the cyclopentadienyl (Cp) rings. The two Cp rings are virtually parallel, with an angle between the two least-squares planes of  $0.74(16)^\circ$ . The distances between the  $\text{Fe}^{\text{II}}$  atom and the centroids of the two Cp rings are 1.639 (2) and 1.652 (2) Å. The conformation of the ferrocenyl moiety is slightly away from eclipsed. The dihedral angle between the coumarin ring system and the ferrocenyl ester moiety is  $69.17(19)^\circ$ .  $\pi$ - $\pi$  stacking interactions involving the benzene rings of neighbouring coumarin moieties, with centroid-centroid distances of 3.739 (2) Å, consolidate the crystal packing.

**Keywords:** crystal structure; ferrocene; coumarin; pharmacological activity 4-methyl-2-oxo-2H-chromene-7-yl ferrocenecarboxylate.

**CCDC reference:** 1027955

## 1. Related literature

For background to ferrocene and its derivatives, see: Štěpnička (2002). For coumarin and its pharmacological activities, see: Peng *et al.* (2013). For the crystal structures of related ferrocenyl derivatives, see: Chen & Lu (2004); Imrie *et al.* (2002, 2005).



## 2. Experimental

### 2.1. Crystal data

$[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_{16}\text{H}_{11}\text{O}_4)]$   
 $M_r = 388.19$   
 Monoclinic,  $P2_1/c$   
 $a = 7.8678(11)$  Å  
 $b = 20.294(4)$  Å  
 $c = 11.1455(18)$  Å  
 $\beta = 108.243(14)^\circ$

$V = 1690.1(5)$  Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.92$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.20 \times 0.10 \times 0.10$  mm

### 2.2. Data collection

Bruker APEXII CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2007)  
 $T_{\text{min}} = 0.838$ ,  $T_{\text{max}} = 0.914$

11857 measured reflections  
 2979 independent reflections  
 2398 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.031$

### 2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$   
 $wR(F^2) = 0.087$   
 $S = 1.16$   
 2979 reflections

236 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.23$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.18$  e Å<sup>-3</sup>

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

## Acknowledgements

This work was supported by a grant from the National Natural Science Foundation of China (No. 81403318) and the Natural Science Foundation of Education Committee of Anhui Province (No. KJ2014A134).

Supporting information for this paper is available from the IUCr electronic archives (Reference: WM5068).

## References

- Bruker (2007). APEX2, SADABS and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Chen, W. Y. & Lu, J. (2004). *Chin. Chem. Lett.* **15**, 1146–1148.  
 Imrie, C., Elago, E. R. T., McClelland, C. W. & Williams, N. (2002). *Green Chem.* **4**, 159–160.

Imrie, C., Elago, E. R. T., Williams, N., McClelland, C. W. & Engelbrecht, P. (2005). *J. Organomet. Chem.* **690**, 4959–4966.  
Peng, X. M. L. V., Damu, G. & Zhou, H. (2013). *Curr. Pharm. Des.* **19**, 3884–3930.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Štěpnička, P. (2002). *New J. Chem.* **26**, 567–575.

## supporting information

*Acta Cryst.* (2014). E70, m369–m370 [doi:10.1107/S1600536814022120]

## Crystal structure of 4-methyl-2-oxo-2*H*-chromen-7-yl ferrocenecarboxylate


Juan Yu, Lei-Lei Gao, Peng Huang and Dian-Lei Wang

### S1. Experimental

To a CH<sub>2</sub>Cl<sub>2</sub> solution (30 ml) of ferrocenylcarboxylic acid (1.0 g, 4.3 mmol), 7-hydroxy-4-methylcoumarin (0.76 g, 4.3 mmol), 1-ethyl-3-(dimethylaminopropyl)carbodiimide hydrochloride (EDCI) (0.89 g, 4.5 mmol) and *N,N*-dimethylamino-pyridine (DMAP) (0.26 g, 2.2 mmol) were added into a 50 ml round bottom flask. Then the reaction mixture was stirred at room temperature. The reaction process was monitored by thin layer chromatography (TLC). After completion of the reaction, the product was extracted with ethyl acetate. The extracts were combined, washed with water and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. After removing ethyl acetate under reduced pressure, the crude product was purified by column chromatography using petroleum ether/EtOAc (8:1) as eluent. Yellow crystals were obtained by slow evaporation of an ether/EtOAc (8:1) solution to give 1.03 g (62% yield) of the title compound, <sup>1</sup>H-NMR (CDCl<sub>3</sub>, δ p.p.m.), 2.48 (s, 2H, CH<sub>3</sub>), 4.34 (s, 5H, C<sub>5</sub>H<sub>5</sub>), 4.57 (s, 2H, C<sub>5</sub>H<sub>2</sub>), 5.00 (s, 2H, C<sub>5</sub>H<sub>2</sub>), 7.18–7.22 (m, 2H, ArH), 7.68 (d, 1H, ArH).

### S2. Refinement

All hydrogen atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H = 0.93 Å and  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}$ .



**Figure 1**

The molecular structure of the title compound showing atoms as ellipsoids at the 30% probability level.

## 4-Methyl-2-oxo-2H-chromen-7-yl ferrocenecarboxylate

## Crystal data

[Fe(C<sub>5</sub>H<sub>5</sub>)(C<sub>16</sub>H<sub>11</sub>O<sub>4</sub>)] $M_r = 388.19$ Monoclinic,  $P2_1/c$ 

Hall symbol: -P 2ybc

 $a = 7.8678$  (11) Å $b = 20.294$  (4) Å $c = 11.1455$  (18) Å $\beta = 108.243$  (14)° $V = 1690.1$  (5) Å<sup>3</sup> $Z = 4$  $F(000) = 800$  $D_x = 1.526$  Mg m<sup>-3</sup>Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 3613 reflections

 $\theta = 2.7$ – $23.4$ ° $\mu = 0.92$  mm<sup>-1</sup> $T = 293$  K

Block, yellow

 $0.20 \times 0.10 \times 0.10$  mm

## Data collection

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(SADABS; Bruker, 2007)

 $T_{\min} = 0.838$ ,  $T_{\max} = 0.914$ 

11857 measured reflections

2979 independent reflections

2398 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.031$  $\theta_{\text{max}} = 25.0$ °,  $\theta_{\text{min}} = 2.0$ ° $h = -9 \rightarrow 9$  $k = -24 \rightarrow 24$  $l = -13 \rightarrow 12$ 

## Refinement

Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.031$  $wR(F^2) = 0.087$  $S = 1.16$ 

2979 reflections

236 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0432P)^2]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\text{max}} = 0.001$  $\Delta\rho_{\text{max}} = 0.23$  e Å<sup>-3</sup> $\Delta\rho_{\text{min}} = -0.18$  e Å<sup>-3</sup>

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.74731 (4)	0.157566 (14)	0.42513 (3)	0.04477 (13)
O1	0.4801 (2)	0.07407 (8)	0.61150 (16)	0.0717 (5)
O2	0.7574 (2)	0.09846 (8)	0.74074 (15)	0.0695 (5)
O3	0.7829 (2)	-0.13167 (8)	0.81894 (15)	0.0654 (5)

O4	0.7893 (3)	-0.23952 (9)	0.8434 (2)	0.0910 (6)
C1	0.7620 (3)	0.25205 (11)	0.4903 (2)	0.0620 (7)
H1	0.8464	0.2857	0.4811	0.074*
C2	0.5913 (3)	0.24019 (11)	0.4035 (2)	0.0616 (6)
H2	0.5372	0.2640	0.3242	0.074*
C3	0.5129 (3)	0.18719 (11)	0.4494 (2)	0.0549 (6)
H3	0.3943	0.1683	0.4083	0.066*
C4	0.6368 (3)	0.16608 (10)	0.5653 (2)	0.0490 (5)
C5	0.7925 (3)	0.20659 (10)	0.5906 (2)	0.0560 (6)
H5	0.9001	0.2037	0.6644	0.067*
C6	0.9160 (4)	0.15047 (13)	0.3194 (3)	0.0734 (8)
H6	0.9804	0.1869	0.2955	0.088*
C7	0.7462 (4)	0.12856 (15)	0.2511 (2)	0.0771 (8)
H7	0.6698	0.1464	0.1707	0.093*
C8	0.7034 (4)	0.07557 (13)	0.3160 (3)	0.0764 (8)
H8	0.5928	0.0497	0.2897	0.092*
C9	0.8462 (4)	0.06495 (12)	0.4251 (3)	0.0758 (9)
H9	0.8544	0.0310	0.4892	0.091*
C10	0.9783 (4)	0.11104 (14)	0.4274 (3)	0.0737 (8)
H10	1.0946	0.1155	0.4929	0.088*
C11	0.6091 (3)	0.10858 (11)	0.6366 (2)	0.0533 (6)
C12	0.7538 (3)	0.04467 (11)	0.8189 (2)	0.0573 (6)
C13	0.7677 (3)	-0.01842 (12)	0.7792 (2)	0.0586 (6)
H13	0.7736	-0.0268	0.6986	0.070*
C14	0.7727 (3)	-0.06931 (11)	0.8629 (2)	0.0497 (5)
C15	0.7663 (3)	-0.05849 (10)	0.9842 (2)	0.0478 (5)
C16	0.7540 (3)	0.00656 (11)	1.0201 (2)	0.0609 (6)
H16	0.7509	0.0155	1.1012	0.073*
C17	0.7463 (3)	0.05785 (12)	0.9382 (2)	0.0633 (7)
H17	0.7361	0.1010	0.9631	0.076*
C18	0.7724 (3)	-0.11504 (12)	1.0658 (2)	0.0528 (6)
C19	0.7797 (3)	-0.17514 (12)	1.0191 (3)	0.0624 (7)
H19	0.7826	-0.2113	1.0709	0.075*
C20	0.7833 (3)	-0.18689 (13)	0.8926 (3)	0.0674 (7)
C21	0.7700 (4)	-0.10441 (13)	1.1988 (2)	0.0731 (8)
H21A	0.7715	-0.1463	1.2392	0.110*
H21B	0.6636	-0.0808	1.1972	0.110*
H21C	0.8733	-0.0794	1.2449	0.110*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Fe1	0.0529 (2)	0.03700 (19)	0.0492 (2)	0.00541 (13)	0.02285 (16)	0.00617 (13)
O1	0.0629 (11)	0.0796 (12)	0.0749 (12)	-0.0128 (10)	0.0250 (10)	0.0156 (10)
O2	0.0814 (12)	0.0671 (11)	0.0539 (10)	-0.0207 (9)	0.0125 (9)	0.0164 (8)
O3	0.0824 (13)	0.0564 (10)	0.0612 (10)	0.0045 (9)	0.0279 (9)	-0.0112 (8)
O4	0.1009 (15)	0.0562 (11)	0.1085 (16)	0.0149 (10)	0.0219 (13)	-0.0211 (11)
C1	0.0792 (19)	0.0351 (11)	0.0761 (18)	0.0034 (11)	0.0306 (16)	0.0030 (12)

C2	0.0711 (17)	0.0467 (13)	0.0711 (17)	0.0193 (12)	0.0282 (14)	0.0145 (12)
C3	0.0530 (14)	0.0545 (13)	0.0617 (15)	0.0134 (11)	0.0244 (12)	0.0002 (12)
C4	0.0602 (15)	0.0440 (12)	0.0487 (13)	0.0060 (10)	0.0253 (12)	-0.0004 (10)
C5	0.0695 (16)	0.0426 (12)	0.0558 (14)	-0.0003 (11)	0.0196 (12)	-0.0036 (11)
C6	0.089 (2)	0.0639 (16)	0.090 (2)	0.0025 (15)	0.0609 (19)	0.0054 (15)
C7	0.100 (2)	0.0832 (19)	0.0534 (16)	0.0245 (17)	0.0307 (16)	-0.0026 (15)
C8	0.082 (2)	0.0568 (16)	0.106 (2)	-0.0087 (14)	0.052 (2)	-0.0307 (16)
C9	0.106 (2)	0.0499 (14)	0.096 (2)	0.0350 (16)	0.067 (2)	0.0242 (14)
C10	0.0569 (16)	0.089 (2)	0.0793 (19)	0.0225 (15)	0.0268 (15)	-0.0039 (16)
C11	0.0627 (16)	0.0550 (14)	0.0498 (14)	0.0015 (12)	0.0285 (13)	0.0003 (11)
C12	0.0632 (16)	0.0573 (14)	0.0481 (14)	-0.0121 (11)	0.0128 (12)	0.0100 (11)
C13	0.0683 (16)	0.0676 (16)	0.0429 (13)	-0.0056 (13)	0.0218 (12)	0.0003 (12)
C14	0.0499 (13)	0.0500 (12)	0.0511 (14)	-0.0007 (10)	0.0187 (11)	-0.0044 (11)
C15	0.0503 (13)	0.0486 (12)	0.0468 (13)	-0.0011 (10)	0.0184 (11)	-0.0001 (10)
C16	0.0886 (18)	0.0535 (14)	0.0466 (14)	-0.0041 (13)	0.0298 (13)	-0.0036 (11)
C17	0.092 (2)	0.0468 (13)	0.0555 (16)	-0.0043 (12)	0.0292 (14)	-0.0028 (11)
C18	0.0517 (14)	0.0540 (14)	0.0550 (14)	0.0031 (10)	0.0199 (12)	0.0082 (11)
C19	0.0591 (16)	0.0518 (14)	0.0778 (18)	0.0053 (12)	0.0234 (14)	0.0106 (13)
C20	0.0589 (16)	0.0546 (15)	0.085 (2)	0.0083 (12)	0.0177 (14)	-0.0042 (15)
C21	0.092 (2)	0.0718 (17)	0.0631 (17)	0.0049 (15)	0.0353 (15)	0.0209 (13)

*Geometric parameters (Å, °)*

Fe1—C4	2.019 (2)	C6—C10	1.400 (4)
Fe1—C7	2.024 (2)	C6—H6	0.9793
Fe1—C8	2.026 (2)	C7—C8	1.395 (4)
Fe1—C5	2.027 (2)	C7—H7	0.9791
Fe1—C9	2.034 (2)	C8—C9	1.390 (4)
Fe1—C6	2.037 (3)	C8—H8	0.9793
Fe1—C3	2.037 (2)	C9—C10	1.393 (4)
Fe1—C1	2.041 (2)	C9—H9	0.9794
Fe1—C10	2.042 (2)	C10—H10	0.9796
Fe1—C2	2.047 (2)	C12—C13	1.370 (3)
O1—C11	1.192 (3)	C12—C17	1.376 (3)
O2—C11	1.379 (3)	C13—C14	1.384 (3)
O2—C12	1.402 (3)	C13—H13	0.9300
O3—C14	1.368 (3)	C14—C15	1.387 (3)
O3—C20	1.388 (3)	C15—C16	1.391 (3)
O4—C20	1.208 (3)	C15—C18	1.456 (3)
C1—C2	1.408 (3)	C16—C17	1.373 (3)
C1—C5	1.411 (3)	C16—H16	0.9300
C1—H1	0.9799	C17—H17	0.9300
C2—C3	1.413 (3)	C18—C19	1.335 (3)
C2—H2	0.9799	C18—C21	1.503 (3)
C3—C4	1.419 (3)	C19—C20	1.439 (4)
C3—H3	0.9800	C19—H19	0.9300
C4—C5	1.428 (3)	C21—H21A	0.9600
C4—C11	1.466 (3)	C21—H21B	0.9600

C5—H5	0.9800	C21—H21C	0.9600
C6—C7	1.387 (4)		
C4—Fe1—C7	153.19 (12)	C1—C5—Fe1	70.25 (13)
C4—Fe1—C8	120.07 (10)	C4—C5—Fe1	69.05 (13)
C7—Fe1—C8	40.29 (11)	C1—C5—H5	126.3
C4—Fe1—C5	41.35 (9)	C4—C5—H5	126.4
C7—Fe1—C5	164.06 (12)	Fe1—C5—H5	126.4
C8—Fe1—C5	153.89 (12)	C7—C6—C10	107.9 (3)
C4—Fe1—C9	109.67 (9)	C7—C6—Fe1	69.54 (16)
C7—Fe1—C9	67.55 (11)	C10—C6—Fe1	70.10 (15)
C8—Fe1—C9	40.03 (12)	C7—C6—H6	125.7
C5—Fe1—C9	119.71 (11)	C10—C6—H6	126.4
C4—Fe1—C6	165.92 (12)	Fe1—C6—H6	126.2
C7—Fe1—C6	39.94 (11)	C6—C7—C8	108.1 (3)
C8—Fe1—C6	67.30 (11)	C6—C7—Fe1	70.52 (15)
C5—Fe1—C6	126.92 (12)	C8—C7—Fe1	69.91 (15)
C9—Fe1—C6	67.43 (10)	C6—C7—H7	126.4
C4—Fe1—C3	40.94 (9)	C8—C7—H7	125.5
C7—Fe1—C3	118.74 (12)	Fe1—C7—H7	125.9
C8—Fe1—C3	109.32 (11)	C9—C8—C7	108.2 (3)
C5—Fe1—C3	69.04 (10)	C9—C8—Fe1	70.30 (15)
C9—Fe1—C3	129.41 (11)	C7—C8—Fe1	69.80 (15)
C6—Fe1—C3	151.50 (12)	C9—C8—H8	125.3
C4—Fe1—C1	68.54 (9)	C7—C8—H8	126.5
C7—Fe1—C1	126.66 (11)	Fe1—C8—H8	126.0
C8—Fe1—C1	164.88 (13)	C8—C9—C10	107.8 (2)
C5—Fe1—C1	40.59 (9)	C8—C9—Fe1	69.67 (14)
C9—Fe1—C1	152.77 (13)	C10—C9—Fe1	70.30 (14)
C6—Fe1—C1	107.33 (10)	C8—C9—H9	126.7
C3—Fe1—C1	68.16 (10)	C10—C9—H9	125.5
C4—Fe1—C10	128.88 (11)	Fe1—C9—H9	126.0
C7—Fe1—C10	67.28 (12)	C9—C10—C6	108.0 (3)
C8—Fe1—C10	67.12 (12)	C9—C10—Fe1	69.73 (14)
C5—Fe1—C10	108.31 (11)	C6—C10—Fe1	69.76 (15)
C9—Fe1—C10	39.97 (11)	C9—C10—H10	126.3
C6—Fe1—C10	40.14 (11)	C6—C10—H10	125.6
C3—Fe1—C10	167.07 (11)	Fe1—C10—H10	125.9
C1—Fe1—C10	118.65 (11)	O1—C11—O2	122.9 (2)
C4—Fe1—C2	68.50 (9)	O1—C11—C4	126.9 (2)
C7—Fe1—C2	107.62 (11)	O2—C11—C4	110.1 (2)
C8—Fe1—C2	128.22 (13)	C13—C12—C17	121.7 (2)
C5—Fe1—C2	68.47 (10)	C13—C12—O2	120.5 (2)
C9—Fe1—C2	166.40 (13)	C17—C12—O2	117.7 (2)
C6—Fe1—C2	117.71 (10)	C12—C13—C14	117.9 (2)
C3—Fe1—C2	40.49 (9)	C12—C13—H13	121.0
C1—Fe1—C2	40.30 (9)	C14—C13—H13	121.0
C10—Fe1—C2	151.52 (11)	O3—C14—C13	116.2 (2)

C11—O2—C12	117.52 (18)	O3—C14—C15	121.31 (19)
C14—O3—C20	121.64 (19)	C13—C14—C15	122.5 (2)
C2—C1—C5	108.8 (2)	C14—C15—C16	117.2 (2)
C2—C1—Fe1	70.10 (13)	C14—C15—C18	118.70 (19)
C5—C1—Fe1	69.16 (13)	C16—C15—C18	124.1 (2)
C2—C1—H1	125.3	C17—C16—C15	121.4 (2)
C5—C1—H1	125.9	C17—C16—H16	119.3
Fe1—C1—H1	125.6	C15—C16—H16	119.3
C1—C2—C3	108.1 (2)	C16—C17—C12	119.2 (2)
C1—C2—Fe1	69.60 (13)	C16—C17—H17	120.4
C3—C2—Fe1	69.36 (12)	C12—C17—H17	120.4
C1—C2—H2	125.9	C19—C18—C15	118.3 (2)
C3—C2—H2	125.9	C19—C18—C21	122.1 (2)
Fe1—C2—H2	125.8	C15—C18—C21	119.6 (2)
C2—C3—C4	107.8 (2)	C18—C19—C20	123.4 (2)
C2—C3—Fe1	70.14 (13)	C18—C19—H19	118.3
C4—C3—Fe1	68.86 (13)	C20—C19—H19	118.3
C2—C3—H3	126.0	O4—C20—O3	116.0 (3)
C4—C3—H3	126.1	O4—C20—C19	127.3 (3)
Fe1—C3—H3	126.3	O3—C20—C19	116.6 (2)
C3—C4—C5	108.0 (2)	C18—C21—H21A	109.5
C3—C4—C11	123.9 (2)	C18—C21—H21B	109.5
C5—C4—C11	127.9 (2)	H21A—C21—H21B	109.5
C3—C4—Fe1	70.20 (12)	C18—C21—H21C	109.5
C5—C4—Fe1	69.60 (13)	H21A—C21—H21C	109.5
C11—C4—Fe1	121.63 (15)	H21B—C21—H21C	109.5
C1—C5—C4	107.3 (2)		
C4—Fe1—C1—C2	-81.64 (15)	C5—Fe1—C6—C10	73.7 (2)
C7—Fe1—C1—C2	72.90 (19)	C9—Fe1—C6—C10	-37.39 (17)
C8—Fe1—C1—C2	45.9 (5)	C3—Fe1—C6—C10	-169.9 (2)
C5—Fe1—C1—C2	-120.3 (2)	C1—Fe1—C6—C10	114.14 (18)
C9—Fe1—C1—C2	-173.15 (19)	C2—Fe1—C6—C10	156.54 (17)
C6—Fe1—C1—C2	112.64 (16)	C10—C6—C7—C8	-0.2 (3)
C3—Fe1—C1—C2	-37.43 (14)	Fe1—C6—C7—C8	-60.10 (18)
C10—Fe1—C1—C2	154.74 (15)	C10—C6—C7—Fe1	59.88 (19)
C4—Fe1—C1—C5	38.64 (14)	C4—Fe1—C7—C6	-170.58 (19)
C7—Fe1—C1—C5	-166.82 (16)	C8—Fe1—C7—C6	-118.6 (2)
C8—Fe1—C1—C5	166.2 (4)	C5—Fe1—C7—C6	39.2 (5)
C9—Fe1—C1—C5	-52.9 (3)	C9—Fe1—C7—C6	-81.18 (18)
C6—Fe1—C1—C5	-127.08 (17)	C3—Fe1—C7—C6	154.99 (15)
C3—Fe1—C1—C5	82.85 (16)	C1—Fe1—C7—C6	71.9 (2)
C10—Fe1—C1—C5	-84.98 (17)	C10—Fe1—C7—C6	-37.73 (16)
C2—Fe1—C1—C5	120.3 (2)	C2—Fe1—C7—C6	112.36 (17)
C5—C1—C2—C3	0.3 (3)	C4—Fe1—C7—C8	-51.9 (3)
Fe1—C1—C2—C3	58.83 (16)	C5—Fe1—C7—C8	157.9 (3)
C5—C1—C2—Fe1	-58.48 (16)	C9—Fe1—C7—C8	37.47 (18)
C4—Fe1—C2—C1	81.76 (15)	C6—Fe1—C7—C8	118.6 (2)



C7—Fe1—C2—C1	-126.44 (17)	C3—Fe1—C7—C8	-86.37 (19)
C8—Fe1—C2—C1	-166.21 (15)	C1—Fe1—C7—C8	-169.44 (17)
C5—Fe1—C2—C1	37.16 (14)	C10—Fe1—C7—C8	80.91 (19)
C9—Fe1—C2—C1	166.6 (4)	C2—Fe1—C7—C8	-128.99 (18)
C6—Fe1—C2—C1	-84.32 (18)	C6—C7—C8—C9	0.4 (3)
C3—Fe1—C2—C1	119.7 (2)	Fe1—C7—C8—C9	-60.04 (18)
C10—Fe1—C2—C1	-51.8 (3)	C6—C7—C8—Fe1	60.48 (18)
C4—Fe1—C2—C3	-37.91 (14)	C4—Fe1—C8—C9	-85.15 (18)
C7—Fe1—C2—C3	113.89 (17)	C7—Fe1—C8—C9	119.1 (2)
C8—Fe1—C2—C3	74.12 (19)	C5—Fe1—C8—C9	-47.3 (3)
C5—Fe1—C2—C3	-82.52 (15)	C6—Fe1—C8—C9	81.44 (18)
C9—Fe1—C2—C3	46.9 (5)	C3—Fe1—C8—C9	-128.94 (16)
C6—Fe1—C2—C3	156.00 (16)	C1—Fe1—C8—C9	153.4 (4)
C1—Fe1—C2—C3	-119.7 (2)	C10—Fe1—C8—C9	37.73 (16)
C10—Fe1—C2—C3	-171.4 (2)	C2—Fe1—C8—C9	-170.38 (15)
C1—C2—C3—C4	-0.2 (3)	C4—Fe1—C8—C7	155.77 (17)
Fe1—C2—C3—C4	58.74 (15)	C5—Fe1—C8—C7	-166.4 (2)
C1—C2—C3—Fe1	-58.98 (16)	C9—Fe1—C8—C7	-119.1 (2)
C4—Fe1—C3—C2	119.3 (2)	C6—Fe1—C8—C7	-37.64 (17)
C7—Fe1—C3—C2	-83.65 (19)	C3—Fe1—C8—C7	111.99 (18)
C8—Fe1—C3—C2	-126.80 (18)	C1—Fe1—C8—C7	34.3 (5)
C5—Fe1—C3—C2	81.00 (16)	C10—Fe1—C8—C7	-81.35 (18)
C9—Fe1—C3—C2	-167.16 (17)	C2—Fe1—C8—C7	70.5 (2)
C6—Fe1—C3—C2	-49.0 (3)	C7—C8—C9—C10	-0.5 (3)
C1—Fe1—C3—C2	37.26 (14)	Fe1—C8—C9—C10	-60.22 (18)
C10—Fe1—C3—C2	161.5 (4)	C7—C8—C9—Fe1	59.72 (18)
C7—Fe1—C3—C4	157.10 (15)	C4—Fe1—C9—C8	113.68 (17)
C8—Fe1—C3—C4	113.95 (16)	C7—Fe1—C9—C8	-37.70 (17)
C5—Fe1—C3—C4	-38.26 (13)	C5—Fe1—C9—C8	158.13 (16)
C9—Fe1—C3—C4	73.59 (19)	C6—Fe1—C9—C8	-81.09 (19)
C6—Fe1—C3—C4	-168.2 (2)	C3—Fe1—C9—C8	71.81 (19)
C1—Fe1—C3—C4	-81.99 (15)	C1—Fe1—C9—C8	-165.2 (2)
C10—Fe1—C3—C4	42.2 (5)	C10—Fe1—C9—C8	-118.6 (2)
C2—Fe1—C3—C4	-119.3 (2)	C2—Fe1—C9—C8	33.9 (5)
C2—C3—C4—C5	0.1 (3)	C4—Fe1—C9—C10	-127.68 (17)
Fe1—C3—C4—C5	59.59 (15)	C7—Fe1—C9—C10	80.94 (18)
C2—C3—C4—C11	-175.0 (2)	C8—Fe1—C9—C10	118.6 (2)
Fe1—C3—C4—C11	-115.4 (2)	C5—Fe1—C9—C10	-83.24 (18)
C2—C3—C4—Fe1	-59.54 (15)	C6—Fe1—C9—C10	37.54 (17)
C7—Fe1—C4—C3	-49.2 (3)	C3—Fe1—C9—C10	-169.56 (16)
C8—Fe1—C4—C3	-85.19 (17)	C1—Fe1—C9—C10	-46.6 (3)
C5—Fe1—C4—C3	118.92 (19)	C2—Fe1—C9—C10	152.6 (4)
C9—Fe1—C4—C3	-128.09 (16)	C8—C9—C10—C6	0.4 (3)
C6—Fe1—C4—C3	156.4 (4)	Fe1—C9—C10—C6	-59.46 (18)
C1—Fe1—C4—C3	80.97 (15)	C8—C9—C10—Fe1	59.82 (17)
C10—Fe1—C4—C3	-168.86 (15)	C7—C6—C10—C9	-0.1 (3)
C2—Fe1—C4—C3	37.51 (14)	Fe1—C6—C10—C9	59.44 (17)
C7—Fe1—C4—C5	-168.1 (2)	C7—C6—C10—Fe1	-59.53 (19)

C8—Fe1—C4—C5	155.88 (16)	C4—Fe1—C10—C9	73.2 (2)
C9—Fe1—C4—C5	112.99 (16)	C7—Fe1—C10—C9	-81.66 (19)
C6—Fe1—C4—C5	37.5 (4)	C8—Fe1—C10—C9	-37.79 (17)
C3—Fe1—C4—C5	-118.92 (19)	C5—Fe1—C10—C9	114.70 (18)
C1—Fe1—C4—C5	-37.95 (14)	C6—Fe1—C10—C9	-119.2 (2)
C10—Fe1—C4—C5	72.22 (18)	C3—Fe1—C10—C9	38.7 (5)
C2—Fe1—C4—C5	-81.41 (15)	C1—Fe1—C10—C9	157.75 (17)
C7—Fe1—C4—C11	69.1 (3)	C2—Fe1—C10—C9	-166.9 (2)
C8—Fe1—C4—C11	33.1 (3)	C4—Fe1—C10—C6	-167.59 (16)
C5—Fe1—C4—C11	-122.8 (3)	C7—Fe1—C10—C6	37.55 (17)
C9—Fe1—C4—C11	-9.8 (2)	C8—Fe1—C10—C6	81.42 (19)
C6—Fe1—C4—C11	-85.3 (4)	C5—Fe1—C10—C6	-126.09 (18)
C3—Fe1—C4—C11	118.3 (3)	C9—Fe1—C10—C6	119.2 (2)
C1—Fe1—C4—C11	-160.7 (2)	C3—Fe1—C10—C6	158.0 (4)
C10—Fe1—C4—C11	-50.6 (3)	C1—Fe1—C10—C6	-83.04 (19)
C2—Fe1—C4—C11	155.8 (2)	C2—Fe1—C10—C6	-47.7 (3)
C2—C1—C5—C4	-0.3 (3)	C12—O2—C11—O1	0.4 (3)
Fe1—C1—C5—C4	-59.37 (15)	C12—O2—C11—C4	-179.32 (19)
C2—C1—C5—Fe1	59.06 (16)	C3—C4—C11—O1	-3.3 (4)
C3—C4—C5—C1	0.2 (3)	C5—C4—C11—O1	-177.3 (2)
C11—C4—C5—C1	174.9 (2)	Fe1—C4—C11—O1	-89.5 (3)
Fe1—C4—C5—C1	60.13 (16)	C3—C4—C11—O2	176.36 (19)
C3—C4—C5—Fe1	-59.97 (15)	C5—C4—C11—O2	2.4 (3)
C11—C4—C5—Fe1	114.8 (2)	Fe1—C4—C11—O2	90.1 (2)
C4—Fe1—C5—C1	-118.4 (2)	C11—O2—C12—C13	72.1 (3)
C7—Fe1—C5—C1	41.8 (4)	C11—O2—C12—C17	-111.6 (3)
C8—Fe1—C5—C1	-171.8 (2)	C17—C12—C13—C14	0.5 (4)
C9—Fe1—C5—C1	155.16 (16)	O2—C12—C13—C14	176.7 (2)
C6—Fe1—C5—C1	72.29 (19)	C20—O3—C14—C13	-178.0 (2)
C3—Fe1—C5—C1	-80.49 (15)	C20—O3—C14—C15	1.4 (3)
C10—Fe1—C5—C1	112.95 (17)	C12—C13—C14—O3	178.7 (2)
C2—Fe1—C5—C1	-36.90 (14)	C12—C13—C14—C15	-0.8 (4)
C7—Fe1—C5—C4	160.2 (4)	O3—C14—C15—C16	-179.2 (2)
C8—Fe1—C5—C4	-53.5 (3)	C13—C14—C15—C16	0.2 (4)
C9—Fe1—C5—C4	-86.45 (17)	O3—C14—C15—C18	0.6 (3)
C6—Fe1—C5—C4	-169.33 (14)	C13—C14—C15—C18	180.0 (2)
C3—Fe1—C5—C4	37.89 (13)	C14—C15—C16—C17	0.8 (4)
C1—Fe1—C5—C4	118.4 (2)	C18—C15—C16—C17	-179.1 (2)
C10—Fe1—C5—C4	-128.67 (15)	C15—C16—C17—C12	-1.1 (4)
C2—Fe1—C5—C4	81.48 (15)	C13—C12—C17—C16	0.4 (4)
C4—Fe1—C6—C7	162.3 (3)	O2—C12—C17—C16	-175.9 (2)
C8—Fe1—C6—C7	37.96 (18)	C14—C15—C18—C19	-1.5 (3)
C5—Fe1—C6—C7	-167.46 (16)	C16—C15—C18—C19	178.3 (2)
C9—Fe1—C6—C7	81.50 (19)	C14—C15—C18—C21	178.6 (2)
C3—Fe1—C6—C7	-51.0 (3)	C16—C15—C18—C21	-1.6 (4)
C1—Fe1—C6—C7	-126.98 (17)	C15—C18—C19—C20	0.5 (4)
C10—Fe1—C6—C7	118.9 (2)	C21—C18—C19—C20	-179.6 (2)
C2—Fe1—C6—C7	-84.58 (19)	C14—O3—C20—O4	178.6 (2)

## supporting information

---

C4—Fe1—C6—C10	43.5 (5)	C14—O3—C20—C19	-2.4 (3)
C7—Fe1—C6—C10	-118.9 (2)	C18—C19—C20—O4	-179.7 (3)
C8—Fe1—C6—C10	-80.92 (19)	C18—C19—C20—O3	1.4 (4)

---