



# Crystal structure of (1*S*,3*R*,8*R*,9*S*,10*R*)-10-bromomethyl-2,2-dichloro-9,10-epoxy-3,7,7-trimethyltricyclo[6.4.0.0<sup>1,3</sup>]-dodecane

Ahmed Benharref,<sup>a</sup> Lahcen El Ammari,<sup>b</sup> Mohamed Saadi<sup>b</sup> and Moha Berraho<sup>a\*</sup>

<sup>a</sup>Laboratoire de Chimie des Substances Naturelles, Unité Associé au CNRST (URAC16), Faculté des Sciences Semlalia, BP 2390 Bd My Abdellah, 40000 Marrakech, Morocco, and <sup>b</sup>Laboratoire de Chimie du Solide Appliquée, Faculté des Sciences, Université Mohammed V, Avenue Ibn Battouta, BP 1014, Rabat, Morocco.  
\*Correspondence e-mail: berraho@uca.ma

Received 27 March 2015; accepted 31 March 2015

Edited by C. Rizzoli, Università degli Studi di Parma, Italy


The title compound, C<sub>16</sub>H<sub>23</sub>BrCl<sub>2</sub>O, was synthesized in three steps from β-himachalene (3,5,5,9-tetramethyl-2,4a,5,6,7,8-hexahydro-1*H*-benzocycloheptene), which was isolated from the essential oil of the Atlas cedar (*cedrus atlantica*). The molecule is built up from two fused six- and seven-membered rings, each linked to a three-membered ring. The six-membered ring has a screw-boat conformation, whereas the seven-membered ring displays a twist-boat conformation. The absolute structure was established unambiguously from anomalous dispersion effects.

**Keywords:** crystal structure; β-himachalene derivative; π–π interactions.

**CCDC reference:** 1057239

## 1. Related literature

For background to β-himachalene, see: El Haib *et al.* (2011). For the reactivity of this sesquiterpene and its derivatives, see: El Jamili *et al.* (2002); Benharref *et al.* (2013); Zaki *et al.* (2014). For the synthesis of the title compound, see: Bimoussa *et al.* (2013). For their potential antifungal activity against the phytopathogen *Botrytis cinerea*, see: Daoubi *et al.* (2004).



## 2. Experimental

### 2.1. Crystal data

C <sub>16</sub> H <sub>23</sub> BrCl <sub>2</sub> O	V = 1677.34 (15) Å <sup>3</sup>
M <sub>r</sub> = 382.15	Z = 4
Orthorhombic, P2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Mo Kα radiation
a = 8.8748 (5) Å	μ = 2.76 mm <sup>-1</sup>
b = 11.2102 (6) Å	T = 296 K
c = 16.8597 (8) Å	0.35 × 0.25 × 0.16 mm

### 2.2. Data collection

Bruker APEXII CCD diffractometer	23384 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 2003)	3419 independent reflections
T <sub>min</sub> = 0.648, T <sub>max</sub> = 0.746	3135 reflections with I > 2σ(I)
	R <sub>int</sub> = 0.034

### 2.3. Refinement

R[F <sup>2</sup> > 2σ(F <sup>2</sup> )] = 0.033	Δρ <sub>min</sub> = -0.53 e Å <sup>-3</sup>
wR(F <sup>2</sup> ) = 0.088	Absolute structure: Flack & Bernardinelli (2000), 1450 Friedel pairs
S = 1.04	Absolute structure parameter: 0.012 (9)
3419 reflections	
184 parameters	
H-atom parameters constrained	
Δρ <sub>max</sub> = 0.65 e Å <sup>-3</sup>	

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT-Plus (Bruker, 2009); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: WinGX (Farrugia, 2012).

## Acknowledgements

The authors thank the Unit of Support for Technical and Scientific Research (UATRS, CNRST) for the X-ray measurements.

Supporting information for this paper is available from the IUCr electronic archives (Reference: RZ5154).

## References

Benharref, A., Ourhiss, N., El Ammari, L., Saadi, M. & Berraho, M. (2013). *Acta Cryst.* **E69**, o933–o934.

- Bimoussa, A., Auhmani, A., Ait Itto, M. Y., Daran, J.-C. & Auhmani, A. (2013). *Acta Cryst.* **E69**, o1692–o1693.
- Bruker (2009). *APEX2* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Daoubi, M., Durán-Patrón, R., Hmamouchi, M., Hernández-Galán, R., Benharref, A. & Collado, I. G. (2004). *Pest. Manag. Sci.* **60**, 927–932.
- El Haib, A., Benharref, A., Parrès-Maynadié, S., Manoury, E., Urrutigoity, M. & Gouygou, M. (2011). *Tetrahedron Asymmetry*, **22**, 101–108.
- El Jamili, H., Auhmani, A., Dakir, M., Lassaba, E., Benharref, A., Pierrot, M., Chiaroni, A. & Riche, C. (2002). *Tetrahedron Lett.* **43**, 6645–6648.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Flack, H. D. & Bernardinelli, G. (2000). *J. Appl. Cryst.* **33**, 1143–1148.
- Sheldrick, G. M. (2003). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Zaki, M., Benharref, A., Daran, J.-C. & Berraho, M. (2014). *Acta Cryst.* **E70**, o526.

## supporting information

*Acta Cryst.* (2015). E71, o284–o285 [https://doi.org/10.1107/S205698901500657X]

## Crystal structure of (1*S*,3*R*,8*R*,9*S*,10*R*)-10-bromomethyl-2,2-dichloro-9,10-epoxy-3,7,7-trimethyltricyclo[6.4.0.0<sup>1,3</sup>]dodecane

Ahmed Benharref, Lahcen El Ammari, Mohamed Saadi and Moha Berraho

### S1. Comment

This work is a part of our ongoing program concerning the valorization of the most abundant essential oils in Morocco, such as *cedrus atlantica*. This oil is made up mainly (75%) of bicyclic sesquiterpene hydrocarbons, among which is found the compound  $\beta$ -himachalene (El Haib *et al.*, 2011). The reactivity of this sesquiterpene and its derivatives has been studied extensively by our team in order to prepare new products having biological properties (El jamili *et al.*, 2002; Benharref *et al.*, 2013, Zaki *et al.*, 2014). Indeed, these compounds were tested, using the food poisoning technique, for their potential antifungal activity against phytopathogen *Botrytis cinerea* (Daoubi *et al.*, 2004). In this work we present the crystal structure of the title compound.

The molecule contains fused six- and seven-membered rings, each linked to a three-membered ring as shown in Fig. 1. The six-membered ring has a screw boat conformation, as indicated by the total puckering amplitude  $Q_T = 0.476$  (3) Å and spherical polar angle  $\theta = 129.5$  (4)° with  $\varphi = 263.4$  (5)°, whereas the seven-membered ring displays a twist boat conformation with  $Q_T = 1.1465$  (34) Å,  $\theta = 88.54$  (18)°,  $\varphi_2 = -152.04$  (18)° and  $\varphi_3 = 72.86$  (6)°. The dihedral angle between the mean planes through the six- and seven-membered rings is 57.7 (2)°. The three-membered rings (C2/C3/O4) and (C6–C8) are nearly perpendicular to the six-membered ring (C1–C6) with a dihedral angles of 79.8 (3)° and 84.7 (3)°, respectively. Owing to the presence of Cl and Br atoms, the absolute configuration could be fully confirmed, by refining the Flack parameter (Flack & Bernardinelli, 2000) as *S*, *R*, *R*, *S* and *R* for C atoms at positions 1, 3, 8, 9 and 10, respectively.

### S2. Experimental

A stoichiometric quantity of *m*-chloroperbenzoic acid (*m*-CPBA) was added to a 100 ml flask containing a solution of (1*S*,3*R*,8*R*)-10-bromomethyl-2,2-dichloro-3,7,7-trimethyltricyclo[6.4.0.0<sup>1,3</sup>]dodec-9-ene (Bimoussa *et al.*, 2013) (750 mg, 2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (40 ml). The reaction mixture was stirred at ambient temperature for 2 h, then treated with a 10% solution of sodium hydrogencarbonate. The aqueous phase was extracted with ether and the organic phases were dried and concentrated. Chromatography of the residue on silica (hexane/ethyl acetate, 98:2 *v/v*) allowed the isolation of the title compound with a yield of 90% (700 mg, 1.8 mmol). The product was recrystallized from cyclohexane.

### S3. Refinement

All H atoms were fixed geometrically and treated as riding with C—H = 0.96–0.98 Å and with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$  or  $1.5 U_{\text{eq}}(\text{C})$  for methyl H atoms.




Figure 1

The molecular structure of the title compound with displacement ellipsoids drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii.

**(1*S*,3*R*,8*R*,9*S*,10*R*)-10-Bromomethyl-2,2-dichloro-9,10-epoxy-3,7,7-trimethyltricyclo[6.4.0.0<sup>1,3</sup>]dodecane**

*Crystal data*

C<sub>16</sub>H<sub>23</sub>BrCl<sub>2</sub>O

*M<sub>r</sub>* = 382.15

Orthorhombic, *P*2<sub>1</sub>2<sub>1</sub>2<sub>1</sub>

Hall symbol: P 2ac 2ab

*a* = 8.8748 (5) Å

*b* = 11.2102 (6) Å

*c* = 16.8597 (8) Å

*V* = 1677.34 (15) Å<sup>3</sup>

*Z* = 4

*F*(000) = 784

*D<sub>x</sub>* = 1.513 Mg m<sup>-3</sup>

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 3419 reflections

θ = 2.2–26.4°

μ = 2.76 mm<sup>-1</sup>

*T* = 296 K

Prism, colourless

0.35 × 0.25 × 0.16 mm

*Data collection*

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω and φ scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 2003)

*T<sub>min</sub>* = 0.648, *T<sub>max</sub>* = 0.746

23384 measured reflections

3419 independent reflections

3135 reflections with *I* > 2σ(*I*)

*R<sub>int</sub>* = 0.034

θ<sub>max</sub> = 26.4°, θ<sub>min</sub> = 2.2°

*h* = -11→11

*k* = -14→14

*l* = -21→14

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.033$  $wR(F^2) = 0.088$  $S = 1.04$ 

3419 reflections

184 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.035P)^2 + 1.0813P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} < 0.001$  $\Delta\rho_{\max} = 0.65 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\min} = -0.53 \text{ e } \text{\AA}^{-3}$ Absolute structure: Flack & Bernardinelli  
(2000), 1450 Friedel pairs

Absolute structure parameter: 0.012 (9)

*Special details*

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against all reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on all data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.8156 (3)	0.2763 (2)	0.75464 (16)	0.0311 (5)
H1	0.8658	0.3365	0.7875	0.037*
C2	0.7910 (3)	0.3332 (2)	0.67448 (17)	0.0349 (6)
H2	0.8806	0.3401	0.6408	0.042*
C3	0.6468 (3)	0.3276 (2)	0.63277 (17)	0.0382 (6)
C4	0.5134 (3)	0.2638 (3)	0.66859 (18)	0.0439 (7)
H4A	0.4679	0.2133	0.6285	0.053*
H4B	0.4389	0.3224	0.6845	0.053*
C5	0.5541 (3)	0.1874 (3)	0.74026 (17)	0.0362 (6)
H5A	0.4632	0.1688	0.7697	0.043*
H5B	0.5976	0.1128	0.7221	0.043*
C6	0.6652 (3)	0.2497 (2)	0.79482 (15)	0.0286 (5)
C7	0.6026 (3)	0.3367 (2)	0.85487 (16)	0.0347 (6)
C8	0.6680 (4)	0.2217 (2)	0.88460 (16)	0.0398 (6)
C9	0.8207 (4)	0.2258 (3)	0.9237 (2)	0.0570 (9)
H9A	0.8075	0.2281	0.9808	0.068*
H9B	0.8719	0.2985	0.9079	0.068*
C10	0.9190 (5)	0.1185 (4)	0.9020 (3)	0.0755 (13)
H10A	1.0238	0.1393	0.9107	0.091*
H10B	0.8946	0.0530	0.9373	0.091*
C11	0.9007 (5)	0.0760 (3)	0.8162 (3)	0.0603 (10)
H11A	0.7992	0.0451	0.8104	0.072*
H11B	0.9693	0.0098	0.8081	0.072*

C12	0.9274 (3)	0.1671 (3)	0.7491 (2)	0.0457 (7)
C13	0.5657 (5)	0.1268 (3)	0.9185 (2)	0.0592 (9)
H13A	0.5498	0.1418	0.9739	0.089*
H13B	0.6116	0.0499	0.9118	0.089*
H13C	0.4707	0.1284	0.8913	0.089*
C14	1.0895 (4)	0.2161 (4)	0.7539 (3)	0.0668 (11)
H14A	1.1026	0.2583	0.8029	0.100*
H14B	1.1071	0.2694	0.7103	0.100*
H14C	1.1597	0.1511	0.7514	0.100*
C15	0.9119 (4)	0.0994 (3)	0.6701 (3)	0.0586 (10)
H15A	0.9802	0.0329	0.6697	0.088*
H15B	0.9354	0.1520	0.6270	0.088*
H15C	0.8105	0.0709	0.6645	0.088*
C16	0.6457 (5)	0.3478 (3)	0.54456 (19)	0.0554 (9)
H16A	0.7275	0.4010	0.5303	0.067*
H16B	0.5517	0.3854	0.5293	0.067*
Br1	0.66817 (6)	0.19765 (4)	0.48772 (2)	0.07821 (17)
O4	0.6911 (3)	0.43512 (16)	0.67456 (12)	0.0418 (5)
Cl1	0.40574 (9)	0.36000 (8)	0.85958 (5)	0.0521 (2)
Cl2	0.69214 (10)	0.47409 (6)	0.87514 (5)	0.04980 (19)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0274 (12)	0.0251 (11)	0.0408 (14)	-0.0013 (10)	-0.0036 (11)	-0.0025 (10)
C2	0.0353 (13)	0.0311 (13)	0.0383 (14)	-0.0042 (10)	0.0061 (11)	-0.0017 (11)
C3	0.0477 (15)	0.0347 (14)	0.0323 (13)	0.0006 (11)	-0.0016 (12)	-0.0043 (11)
C4	0.0345 (14)	0.0591 (19)	0.0382 (15)	-0.0083 (13)	-0.0069 (12)	-0.0046 (13)
C5	0.0305 (13)	0.0362 (13)	0.0418 (15)	-0.0096 (11)	0.0009 (12)	-0.0039 (12)
C6	0.0271 (11)	0.0265 (11)	0.0323 (12)	-0.0027 (10)	-0.0045 (11)	0.0017 (9)
C7	0.0353 (13)	0.0355 (14)	0.0331 (14)	-0.0013 (11)	-0.0001 (11)	0.0001 (11)
C8	0.0485 (16)	0.0368 (14)	0.0341 (14)	-0.0056 (13)	-0.0056 (13)	0.0077 (11)
C9	0.064 (2)	0.061 (2)	0.0459 (18)	-0.0004 (18)	-0.0205 (18)	0.0138 (15)
C10	0.075 (3)	0.071 (3)	0.081 (3)	0.017 (2)	-0.029 (2)	0.027 (2)
C11	0.0554 (19)	0.0392 (16)	0.086 (3)	0.0153 (15)	-0.009 (2)	0.0085 (17)
C12	0.0330 (15)	0.0369 (15)	0.067 (2)	0.0061 (12)	-0.0024 (14)	-0.0027 (14)
C13	0.076 (3)	0.052 (2)	0.049 (2)	-0.0088 (18)	0.0071 (18)	0.0157 (16)
C14	0.0316 (16)	0.060 (2)	0.109 (3)	0.0024 (15)	-0.0107 (18)	-0.006 (2)
C15	0.0434 (17)	0.0463 (18)	0.086 (3)	0.0076 (14)	0.0107 (18)	-0.0195 (18)
C16	0.079 (2)	0.0550 (19)	0.0326 (15)	0.0012 (18)	-0.0015 (16)	-0.0008 (14)
Br1	0.0966 (3)	0.0883 (3)	0.0497 (2)	0.0133 (3)	-0.0049 (2)	-0.0260 (2)
O4	0.0549 (12)	0.0313 (9)	0.0394 (11)	0.0054 (9)	-0.0052 (10)	0.0000 (8)
Cl1	0.0415 (4)	0.0631 (5)	0.0516 (5)	0.0067 (4)	0.0126 (3)	0.0000 (4)
Cl2	0.0638 (5)	0.0353 (3)	0.0504 (4)	-0.0044 (3)	-0.0037 (4)	-0.0109 (3)

*Geometric parameters (Å, °)*

C1—C2	1.510 (4)	C9—C10	1.531 (6)
C1—C6	1.527 (4)	C9—H9A	0.9700
C1—C12	1.579 (4)	C9—H9B	0.9700
C1—H1	0.9800	C10—C11	1.530 (6)
C2—O4	1.446 (3)	C10—H10A	0.9700
C2—C3	1.462 (4)	C10—H10B	0.9700
C2—H2	0.9800	C11—C12	1.543 (5)
C3—O4	1.451 (3)	C11—H11A	0.9700
C3—C16	1.504 (4)	C11—H11B	0.9700
C3—C4	1.509 (4)	C12—C15	1.538 (5)
C4—C5	1.524 (4)	C12—C14	1.542 (4)
C4—H4A	0.9700	C13—H13A	0.9600
C4—H4B	0.9700	C13—H13B	0.9600
C5—C6	1.519 (4)	C13—H13C	0.9600
C5—H5A	0.9700	C14—H14A	0.9600
C5—H5B	0.9700	C14—H14B	0.9600
C6—C7	1.512 (4)	C14—H14C	0.9600
C6—C8	1.546 (4)	C15—H15A	0.9600
C7—C8	1.500 (4)	C15—H15B	0.9600
C7—C12	1.766 (3)	C15—H15C	0.9600
C7—C11	1.769 (3)	C16—Br1	1.947 (3)
C8—C9	1.507 (5)	C16—H16A	0.9700
C8—C13	1.511 (4)	C16—H16B	0.9700
C2—C1—C6	110.7 (2)	C8—C9—H9A	109.1
C2—C1—C12	111.4 (2)	C10—C9—H9A	109.1
C6—C1—C12	115.1 (2)	C8—C9—H9B	109.1
C2—C1—H1	106.3	C10—C9—H9B	109.1
C6—C1—H1	106.3	H9A—C9—H9B	107.8
C12—C1—H1	106.3	C11—C10—C9	114.2 (3)
O4—C2—C3	59.83 (17)	C11—C10—H10A	108.7
O4—C2—C1	114.9 (2)	C9—C10—H10A	108.7
C3—C2—C1	122.6 (2)	C11—C10—H10B	108.7
O4—C2—H2	115.8	C9—C10—H10B	108.7
C3—C2—H2	115.8	H10A—C10—H10B	107.6
C1—C2—H2	115.8	C10—C11—C12	118.1 (3)
O4—C3—C2	59.54 (17)	C10—C11—H11A	107.8
O4—C3—C16	110.9 (2)	C12—C11—H11A	107.8
C2—C3—C16	118.3 (3)	C10—C11—H11B	107.8
O4—C3—C4	114.4 (2)	C12—C11—H11B	107.8
C2—C3—C4	121.0 (2)	H11A—C11—H11B	107.1
C16—C3—C4	117.5 (3)	C15—C12—C14	107.7 (3)
C3—C4—C5	113.4 (2)	C15—C12—C11	107.1 (3)
C3—C4—H4A	108.9	C14—C12—C11	109.9 (3)
C5—C4—H4A	108.9	C15—C12—C1	112.2 (3)
C3—C4—H4B	108.9	C14—C12—C1	107.9 (2)

---

C5—C4—H4B	108.9	C11—C12—C1	111.9 (3)
H4A—C4—H4B	107.7	C8—C13—H13A	109.5
C6—C5—C4	112.1 (2)	C8—C13—H13B	109.5
C6—C5—H5A	109.2	H13A—C13—H13B	109.5
C4—C5—H5A	109.2	C8—C13—H13C	109.5
C6—C5—H5B	109.2	H13A—C13—H13C	109.5
C4—C5—H5B	109.2	H13B—C13—H13C	109.5
H5A—C5—H5B	107.9	C12—C14—H14A	109.5
C7—C6—C5	117.7 (2)	C12—C14—H14B	109.5
C7—C6—C1	119.5 (2)	H14A—C14—H14B	109.5
C5—C6—C1	112.9 (2)	C12—C14—H14C	109.5
C7—C6—C8	58.76 (18)	H14A—C14—H14C	109.5
C5—C6—C8	120.7 (2)	H14B—C14—H14C	109.5
C1—C6—C8	117.4 (2)	C12—C15—H15A	109.5
C8—C7—C6	61.77 (18)	C12—C15—H15B	109.5
C8—C7—C12	120.7 (2)	H15A—C15—H15B	109.5
C6—C7—C12	121.82 (19)	C12—C15—H15C	109.5
C8—C7—C11	119.6 (2)	H15A—C15—H15C	109.5
C6—C7—C11	119.2 (2)	H15B—C15—H15C	109.5
C12—C7—C11	107.87 (15)	C3—C16—Br1	110.8 (2)
C7—C8—C9	117.9 (2)	C3—C16—H16A	109.5
C7—C8—C13	119.9 (3)	Br1—C16—H16A	109.5
C9—C8—C13	113.3 (3)	C3—C16—H16B	109.5
C7—C8—C6	59.47 (17)	Br1—C16—H16B	109.5
C9—C8—C6	115.9 (3)	H16A—C16—H16B	108.1
C13—C8—C6	120.2 (3)	C2—O4—C3	60.63 (18)
C8—C9—C10	112.6 (3)		

---