



Crystal structure of orthorhombic {bis[(pyridin-2-yl)methyl](3,5,5,5-tetrachloropentyl)amine- κ^3N,N',N'' }chloridocopper(II) perchlorate

Katherine A. Bussey,^a Annie R. Cavalier,^a Jennifer R. Connell,^a Margaret E. Mraz,^a Kayode D. Oshin,^{a*} Tomislav Pintauer,^b Danielle L. Gray^c and Sean Parkin^d

Received 6 June 2015

Accepted 19 June 2015

Edited by A. J. Lough, University of Toronto, Canada

^aDepartment of Chemistry & Physics, Saint Mary's College, Notre Dame, IN 46556, USA, ^bDepartment of Chemistry & Biochemistry, Duquesne University, Pittsburgh, PA, 15282, USA, ^cSchool of Chemical Sciences, University of Illinois, Urbana-Champaign, IL 61801, USA, and ^dDepartment of Chemistry, University of Kentucky, Lexington, KY 40506, USA.

*Correspondence e-mail: koshin@saintmarys.edu

Keywords: crystal structure; four-coordinate copper(II); hetero-scorpionate complex; Atom Transfer Radical Addition (ATRA) reactions; disorder in cation and counter-ion

CCDC reference: 1407833

Supporting information: this article has supporting information at journals.iucr.org/e


In the title compound, [CuCl(C₁₇H₁₉Cl₄N₃)]ClO₄, the Cu^{II} ion adopts a distorted square-planar geometry defined by one chloride ligand and the three nitrogen atoms from the bis[(pyridin-2-yl)methyl](3,5,5,5-tetrachloropentyl)amine ligand. The perchlorate counter-ion is disordered over three sets of sites with refined occupancies 0.0634 (17), 0.221 (16) and 0.145 (7). In addition, the hetero-scorpionate arm of the bis[(pyridin-2-yl)methyl](3,5,5,5-tetrachloropentyl)amine ligand is disordered over two sets of sites with refined occupancies 0.839 (2) and 0.161 (2). In the crystal, weak Cu^{II}⋯Cl interactions between symmetry-related molecules create a dimerization with a chloride occupying the apical position of the square-pyramidal geometry typical of many copper(II) chloride hetero-scorpionate complexes.

1. Chemical context

The mechanistic and structural study of Atom Transfer Radical Addition (ATRA) reactions is a growing and promising field in organometallic chemistry. These reactions involve the formation of carbon–carbon bonds through addition of a poly-halogenated saturated hydrocarbon to alkenes (Eckenhoff & Pintauer, 2010). Also known as the Kharasch reaction, most proceed either in the presence of a free-radical precursor as the halogen transfer agent, or a transition metal complex as the halogen transfer agent (Muñoz-Molina *et al.*, 2011). What makes these types of reactions attractive is generation of halogen-group functionalities within the product; which can be used as starting reagents in further functionalization reactions (Kleij *et al.*, 2000). Of interest to this project is analysis of hetero-scorpionate complexes incorporating weakly coordinating olefinic moieties in ATRA reactions. Since their discovery in the 1960s by Swiatoslaw Trofimenko (Pettinari, 2004), scorpionate ligands are considered to be some of the most useful ligand structures available in modern coordination chemistry (Trofimenko, 1999). As such, we report the synthesis and crystal structure of the title compound [Cu(C₁₇H₁₉N₃Cl₄)(Cl)][ClO₄] (**1**).

2. Structural commentary

The title complex, (**1**) (Fig. 1), adopts a distorted square-planar geometry, as shown in the bond angles around the Cu^{II} ion. The Cu^{II} ion is coordinated by the binding of the two pyridine and amine nitrogen atoms and a chlorido ligand. A τ -



4 analysis of the distortions about the Cu^{II} ion yields a value of 0.15, slightly deviant from an ideal value of zero for perfect square-planar geometry [$\tau_4 = [360 - (\alpha + \beta)]/141$; Yang *et al.*, 2007] where α and β are the two greatest valence angles of the coordination center]. The Cu^{II} ion sits 0.0922 (4) Å out of the mean basal plane formed by Cl1 and the three coordinating N atoms, giving rise to the distortion from true square-planar geometry. The Cu–Cl1 [2.2519 (8) Å], Cu–N(amine) [2.027 (2) Å], and Cu–N(py) [1.982 (3) and 1.987 (3) Å] bond lengths are in the anticipated range for copper(II) complexes.





Figure 1
The molecular structure of (**1**), shown with 50% probability ellipsoids for non-H atoms and circles of arbitrary size for H atoms. Only the primary orientations of the disordered sites are shown.

weak Cu···Cl interaction becomes the apical position of a distorted square-pyramidal geometry for the Cu^{II} atoms. Further strengthening the dimer are weak electrostatic C—

3. Supramolecular features

Weak [2.8535 (9) Å] Cu···Cl interactions between adjacent molecules creates a dimerization with two Cl atoms bridging the Cu^{II} atoms (Fig. 2). The inter-copper distance between neighbouring cations is 3.4040 (7) Å. When considered, the

Table 1
Hydrogen-bond geometry (Å, °).

D–H···A	D–H	H···A	D···A	D–H···A
C11–H11A···Cl1 ⁱ	0.99	2.84	3.441 (3)	120
C12–H12B···Cl1 ⁱ	0.99	2.84	3.438 (3)	120
C2–H2···O1 ⁱⁱ	0.95	2.64	3.383 (8)	136
C11–H11A···O4 ⁱⁱⁱ	0.99	2.53	3.123 (7)	118
C12–H12A···O4	0.99	2.43	3.310 (9)	148
C13–H13A···O1 ^{iv}	0.99	2.64	3.578 (11)	157
C14–H14B···O3 ^{iv}	0.99	2.40	3.324 (13)	154
C7–H7···O2 ^{viii}	0.95	2.64	3.14 (3)	113
C12–H12A···O4 ^v	0.99	2.37	3.32 (2)	159
C14–H14C···O3 ^{vii}	0.99	2.17	3.13 (2)	163
C14–H14D···O4 ^v	0.99	2.41	3.26 (2)	143
C16 ^v –H16D···O2 ^{vii}	0.99	2.32	3.28 (3)	162
C7–H7···O2 ^{viii}	0.95	2.57	3.26 (2)	130
C12–H12A···O4 ⁱⁱ	0.99	2.53	3.33 (4)	138
C13–H13A···O1 ^{iv}	0.99	2.45	3.33 (2)	148
C2–H2···Cl4 ^{vii}	0.95	2.84	3.644 (19)	143
C11–H11B···Cl2 ^v	0.99	2.70	3.490 (7)	137
C13–H13B···Cl2 ^{vi}	0.99	2.90	3.863 (4)	165

Symmetry codes: (i) $-x, -y + 2, -z$; (ii) $x - \frac{1}{2}, -y + \frac{3}{2}, -z$; (iii) $-x + \frac{1}{2}, y + \frac{1}{2}, z$; (iv) $-x + \frac{1}{2}, y - \frac{1}{2}, z$; (v) $x, -y + 1, z - \frac{1}{2}$; (vi) $-x, y, -z + \frac{1}{2}$.




Figure 2
Dimer interactions between two [Cu(C₁₇H₁₉N₃Cl₄)(Cl)] (**1**) molecules, shown with 50% probability ellipsoids for the primary orientations of the disordered sites. H atoms are removed for clarity. The symmetry operation to generate the additional cation is $1 - x, 1 - y, 1 - z$.




Figure 3
Packing diagram viewed along the *b*-axis direction showing the electrostatic interactions for the primary orientations of the disordered sites.

H...Cl interactions between C11–H11A...Cl1ⁱ and C12–H12B...Cl1ⁱ (Cl1ⁱ is generated by the symmetry operation $-x, -y + 2, -z$; Table 1). The three-dimensional packing structure (Fig. 3) is comprised from many weak C–H...O interactions that occur between carbon donors on the scorpionate arm or the bis(pyridin-2-ylmethyl)amine and the oxygen atoms on varying orientations of the perchlorate counter-ion. Depending on the orientation of the chlorinated scorpionate arm, there are additional weak C–H...Cl interactions.

4. Database survey

There are 200 structures with the bis(pyridin-2-ylmethyl)amine ligand coordinating to copper with at least one bound chloride ligand (Groom & Allen 2014; CSD Version 5.36). Ignoring all the structures that have tethered pairs or tethered triplets of ligands, or have ligands whose amine group has substituents that additionally coordinate to the Cu^{II} atom, there are 58 remaining structures. Eighteen of these remaining structures have two bridging Cl ligands with one short axial Cu–Cl bond length (average 2.25 Å) and one long apical Cu–Cl bond length (average 2.72 Å).

5. Synthesis and crystallization

The synthetic procedure is outlined in Fig. 4. **Synthesis of 1-butene-bis(pyridin-2-ylmethyl)amine**, (**B**): bis(pyridin-2-ylmethyl)amine (BPMA) precursor (**A**) was synthesized and purified following literature procedures (Carvalho *et al.*, 2006).

BPMA (8.064 g, 40.5 mmol) was dissolved in 15 mL of acetonitrile followed by the addition of triethylamine (4.098 g, 40.5 mmol) and 4-bromobutene (5.468 g 40.5 mmol). The reaction was sealed and allowed to mix for 4 days to ensure complete deprotonation and coupling occurred. Generation of the triethylamine hydrogen bromide salt Et₃NH⁺·Br[−] was observed as white crystals in the brown-colored solution. The mixture was filtered and the desired product extracted from the filtrate using a hexane/water mixture. The hexane layer was separated and solvent removed to yield the ligand as a yellow-colored oil (8.516 g, 83%). The ligand was stored in a septum sealed round-bottom flask under argon gas in a refrigerator. ¹H NMR (CDCl₃, 400 MHz): δ 2.31 (*dd*, *J* = 8.0 and 21.6 Hz, 2H), δ 2.64 (*t*, *J* = 7.2 Hz, 2H), δ 3.83 (*s*, 4H), δ 4.97 (*d*, *J* = 10.4 Hz, 1H), δ 5.01 (*d*, *J* = 18.8 Hz, 1H), δ 5.75 (*m*, *J* = 10.4 Hz, 1H), δ 7.13 (*t*, *J* = 6.4 Hz, 2H), δ 7.53 (*d*, *J* = 8.0 Hz, 2H), δ 7.64 (*t*, *J* = 7.6 Hz, 2H), δ 8.51 (*d*, *J* = 4.4 Hz, 2H). ¹³C NMR (CDCl₃, 400 MHz): δ 159.75, 149.01, 136.38, 135.38, 122.80, 121.88, 117.93, 77.13, 59.90, 57.32. FT-IR (liquid): *ν* (cm^{−1}) = 3066 (*w*), 2922 (*w*), 2816 (*w*), 2158 (*s*), 1639 (*s*), 1588 (*s*), 1361 (*s*), 994 (*w*), 756 (*s*). FT-IR (solid): *ν* (cm^{−1}) = 3394 (*w*), 3067 (*w*), 3008 (*s*), 2923 (*w*), 2817 (*s*), 2359 (*s*), 1619 (*s*), 1589 (*s*), 1432 (*s*).

Synthesis of [Cu^I(butene-bis(pyridin-2-ylmethyl)amine)]-[ClO₄], (**C**): In the drybox, 1-butene-*bis*(pyridin-2-ylmethyl)amine (**A**) (1.00 g, 3.95 mmol) was dissolved in 5 mL acetonitrile in a 50 mL Schlenk flask. Cu(ClO₄) (1.292 g, 3.95 mmol) was added to the flask to give a yellow-colored solution. The reaction was allowed to mix for 6 h, then 15 mL of pentane was slowly added to the solution to generate a yellow precipi-




Figure 4
The synthetic scheme.

pitae. Solvent was removed from the flask through a vacuum line. The precipitate was washed twice by transferring 10 mL of pentane into the flask and stirring vigorously for thirty minutes. Solvent was removed and the precipitate dried under vacuum for 2 h to yield a yellow-colored solid (2.109 g, 92%). $^1\text{H NMR}$ (CD_3CN , 400 MHz): δ 2.45 (*dd*, $J = 8.8$ and 22.4 Hz, 2H), δ 2.77 (*t*, $J = 8.0$ Hz, 2H), δ 3.87 (*s*, 4H), δ 4.92 (*d*, $J = 10.0$ Hz, 1H), δ 4.98 (*d*, $J = 16.8$ Hz, 1H), δ 5.70 (*m*, $J = 10.4$ Hz, 1H), δ 7.33 (*d*, $J = 8.0$ Hz, 2H), δ 7.38 (*t*, $J = 6.0$ Hz, 2H), δ 7.80 (*t*, $J = 7.6$ Hz, 2H), δ 8.63 (*d*, $J = 4.8$ Hz, 2H). FT-IR (solid): ν (cm^{-1}) = 3271 (*w*), 3083 (*w*), 2923 (*w*), 2818 (*w*), 2325 (*s*), 2303 (*s*), 1602 (*s*), 1477 (*s*). TOF-ESI-MS: (m/z) [$M - (\text{ClO}_4)^+$], Calculated for $\text{C}_{16}\text{H}_{19}\text{N}_3\text{Cu}$ 316.0875, found 316.0897 (7 p.p.m.).

Synthesis of [Cu^{II}(1,1,1-trichloro-3-chloropentylbis(pyridin-2-ylmethyl)amine)][Cl][ClO₄] (1): In the drybox, [Cu^I(butene-*bis*(pyridin-2-ylmethyl)amine)][ClO₄] (C) (0.50 g, 1.20 mmol) was dissolved in 5 mL acetonitrile in a glass vial with a stir bar. Nitrogen gas purged CCl_4 (0.174 mL, 1.80 mmol) was added to the vial producing a bluish-green-colored mixture. The reaction vial was sealed with a plastic cap and allowed to mix for 4 h, then removed from the drybox.

Vapour diffusion crystallization at room temperature, incorporating 1 mL of the bluish-green solution with diethyl ether as the external diffusing solvent, produced blue-colored crystals suitable for X-ray analysis.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. The proposed structure model includes disorder of the hetero-scorpionate arm of the bis-[(pyridin-2-yl)methyl](3,5,5,5-tetrachloropentyl)amine ligand over two sets of sites and disorder of the perchlorate anion modelled over three sites. The geometries of the disordered [$\text{C}_5\text{H}_7\text{Cl}_4$] arm were restrained to be the same (s.u. 0.01 Å). The perchlorate anions were also restrained to have the same geometries (s.u. 0.01 Å). In addition, the sum of the occupancies of the three orientations for the perchlorate anions were restrained to add up to one (s.u. 0.001). All disordered sites were restrained to have similar displacement amplitudes (s.u. 0.01) for atoms overlapping by less than the sum of van der Waals radii. Displacement parameters for the perchlorate anion positions were also restrained to behave relatively

Table 2
Experimental details.

Crystal data	
Chemical formula	[CuCl(C ₁₇ H ₁₉ Cl ₄ N ₃)]ClO ₄
<i>M</i> _r	605.59
Crystal system, space group	Orthorhombic, <i>Pbcn</i>
Temperature (K)	150
<i>a</i> , <i>b</i> , <i>c</i> (Å)	17.4845 (13), 10.6593 (8), 25.1030 (18)
<i>V</i> (Å ³)	4678.5 (6)
<i>Z</i>	8
Radiation type	Mo <i>K</i> α
<i>μ</i> (mm ⁻¹)	1.65
Crystal size (mm)	0.72 × 0.31 × 0.04
Data collection	
Diffractometer	Bruker SMART APEXII CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2014)
<i>T</i> _{min} , <i>T</i> _{max}	0.638, 0.746
No. of measured, independent and observed [<i>I</i> > 2σ(<i>I</i>)] reflections	51373, 6305, 4147
<i>R</i> _{int}	0.081
(sin θ/λ) _{max} (Å ⁻¹)	0.686
Refinement	
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.045, 0.108, 1.03
No. of reflections	6305
No. of parameters	437
No. of restraints	647
H-atom treatment	H-atom parameters constrained
Δρ _{max} , Δρ _{min} (e Å ⁻³)	0.56, -0.39

Computer programs: *APEX2*, *SAINT*, *XPREP* and *SADABS* (Bruker, 2014), *SHELXS2014/7* (Sheldrick, 2008), *SHELXL2014/7* (Sheldrick, 2015), *CrystalMaker* (CrystalMaker, 1994) and *pubCIF* (Westrip, 2010).

isotropic. All non-H atoms were refined with anisotropic displacement parameters. H atoms were included as riding

idealized contributors, with C–H = 0.95 (aromatic), 0.99 (*sp*³ C–R2H2), and 1.00 Å (*sp*³ C–R3H). The *U*_{iso}(H) values were set to 1.2*U*_{eq}(C) of the carrier atom. The (002) reflection was omitted from the final refinement because it was partially obscured by the beam stop.

Acknowledgements

The authors would like to thank Duquesne University for instrument support.

References

- Bruker (2014). *APEX2*, *SAINT*, *XPREP*, *SADABS*. Bruker AXS, Inc., Madison, Wisconsin, USA.
- Carvalho, N. M. F., Horn, A. Jr, Bortoluzzi, A. J., Drago, V. & Antunes, O. A. C. (2006). *Inorg. Chim. Acta*, **359**, 90–98.
- CrystalMaker (1994). *CrystalMaker*. CrystalMaker Software Ltd, Oxford, England (www.CrystalMaker.com).
- Eckenhoff, W. T. & Pintauer, T. (2010). *Catal. Rev.* **52**, 1–59.
- Groom, C. R. & Allen, F. H. (2014). *Angew. Chem. Int. Ed.* **53**, 662–671.
- Kleij, A. W., Gossage, R. A., Jastrzebski, J. T. B. H., Boersma, J. & van Koten, G. (2000). *Angew. Chem. Int. Ed.* **39**, 176–178.
- Muñoz-Molina, J. M., Sameera, W. M. C., Álvarez, E., Maseras, F., Belderrain, T. R. & Pérez, P. J. (2011). *Inorg. Chem.* **50**, 2458–2467.
- Pettinari, C. (2004). *Chim. Ind.* **10**, 94–100.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **A71**, 3–8.
- Trofimenko, S. (1999). In *Scorpionates: The Coordination Chemistry of Poly(pyrazolyl)borate Ligands*. London: Imperial College Press.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Yang, L., Powell, D. R. & Houser, R. P. (2007). *Dalton Trans.* pp. 955–964.

supporting information

Acta Cryst. (2015). E71, 847-851 [doi:10.1107/S2056989015011792]

Crystal structure of orthorhombic {bis[(pyridin-2-yl)methyl](3,5,5,5-tetrachloropentyl)amine- κ^3N,N',N'' }chloridocopper(II) perchlorate

Katherine A. Bussey, Annie R. Cavalier, Jennifer R. Connell, Margaret E. Mraz, Kayode D. Oshin, Tomislav Pintauer, Danielle L. Gray and Sean Parkin

Computing details

Data collection: *APEX2* (Bruker, 2014); cell refinement: *SAINT* (Bruker, 2014); data reduction: *SAINT*, *XPREP* and *SADABS* (Bruker, 2014); program(s) used to solve structure: *SHELXS2014/7* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014/7* (Sheldrick, 2015); molecular graphics: *CrystalMaker* (*CrystalMaker*, 1994); software used to prepare material for publication: *publCIF* (Westrip, 2010).

{Bis[(pyridin-2-yl)methyl](1,1,1,3-tetrachloropentyl)amine- κ^3N,N',N'' }chloridocopper(II) perchlorate

Crystal data

[CuCl(C₁₇H₁₉Cl₄N₃)]ClO₄

M_r = 605.59

Orthorhombic, *Pbcn*

a = 17.4845 (13) Å

b = 10.6593 (8) Å

c = 25.1030 (18) Å

V = 4678.5 (6) Å³

Z = 8

F(000) = 2440

D_x = 1.720 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 5135 reflections

θ = 2.2–23.3°

μ = 1.65 mm⁻¹

T = 150 K

Prism, blue

0.72 × 0.31 × 0.04 mm

Data collection

Bruker SMART APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

profile data from φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2014)

T_{min} = 0.638, *T_{max}* = 0.746

51373 measured reflections

6305 independent reflections

4147 reflections with *I* > 2σ(*I*)

R_{int} = 0.081

θ_{max} = 29.2°, θ_{min} = 2.0°

h = -23→23

k = -14→14

l = -34→34

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.045

wR(*F*²) = 0.108

S = 1.03

6305 reflections

437 parameters

647 restraints

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

w = 1/[σ²(*F_o*²) + (0.0376*P*)² + 5.7335*P*]

where *P* = (*F_o*² + 2*F_c*²)/3

(Δ/σ)_{max} = 0.001

Δρ_{max} = 0.56 e Å⁻³

Δρ_{min} = -0.39 e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cu1	0.01473 (2)	0.91372 (4)	0.05612 (2)	0.02853 (11)	
N1	-0.01787 (15)	1.0476 (2)	0.10598 (10)	0.0314 (6)	
N2	0.06695 (16)	0.7705 (2)	0.02079 (10)	0.0307 (6)	
N3	0.10728 (14)	0.9061 (2)	0.10463 (10)	0.0255 (5)	
Cl1	-0.09392 (4)	0.90746 (8)	0.00802 (3)	0.03601 (19)	
C1	0.0368 (2)	0.6854 (3)	-0.01322 (13)	0.0409 (9)	
H1	-0.0161	0.6897	-0.0218	0.049*	
C2	0.0808 (3)	0.5933 (4)	-0.03554 (15)	0.0499 (10)	
H2	0.0586	0.5343	-0.0593	0.060*	
C3	0.1574 (3)	0.5865 (4)	-0.02351 (16)	0.0535 (11)	
H3	0.1884	0.5230	-0.0390	0.064*	
C4	0.1888 (2)	0.6722 (3)	0.01107 (15)	0.0439 (9)	
H4	0.2418	0.6693	0.0195	0.053*	
C5	0.14184 (18)	0.7632 (3)	0.03345 (12)	0.0301 (7)	
C6	0.03905 (18)	1.0893 (3)	0.13719 (12)	0.0299 (7)	
C7	0.0263 (2)	1.1791 (3)	0.17579 (14)	0.0394 (8)	
H7	0.0667	1.2054	0.1984	0.047*	
C8	-0.0459 (2)	1.2302 (3)	0.18115 (16)	0.0481 (10)	
H8	-0.0554	1.2931	0.2071	0.058*	
C9	-0.1039 (2)	1.1889 (4)	0.14842 (17)	0.0495 (10)	
H9	-0.1537	1.2238	0.1511	0.059*	
C10	-0.0885 (2)	1.0960 (4)	0.11176 (15)	0.0411 (9)	
H10	-0.1288	1.0654	0.0900	0.049*	
C11	0.11701 (18)	1.0358 (3)	0.12597 (13)	0.0286 (7)	
H11A	0.1441	1.0887	0.0996	0.034*	
H11B	0.1477	1.0337	0.1591	0.034*	
C12	0.17104 (17)	0.8613 (3)	0.07094 (12)	0.0283 (7)	
H12A	0.2119	0.8256	0.0936	0.034*	
H12B	0.1928	0.9323	0.0505	0.034*	
C13	0.08923 (19)	0.8171 (3)	0.14857 (12)	0.0313 (7)	
H13A	0.0711	0.7373	0.1328	0.038*	
H13B	0.0468	0.8522	0.1700	0.038*	
C14	0.1563 (2)	0.7886 (3)	0.18580 (13)	0.0380 (8)	
H14A	0.1893	0.8635	0.1894	0.046*	0.839 (2)
H14B	0.1875	0.7192	0.1711	0.046*	0.839 (2)
H14C	0.1674	0.6985	0.1800	0.046*	0.161 (2)
H14D	0.2001	0.8345	0.1702	0.046*	0.161 (2)
C15	0.1234 (2)	0.7506 (4)	0.24114 (14)	0.0278 (8)	0.839 (2)
H15	0.0768	0.6978	0.2358	0.033*	0.839 (2)

C12	0.09677 (7)	0.89437 (12)	0.27515 (4)	0.0431 (3)	0.839 (2)
C16	0.1823 (2)	0.6764 (4)	0.27201 (14)	0.0298 (8)	0.839 (2)
H16A	0.2273	0.7311	0.2777	0.036*	0.839 (2)
H16B	0.1991	0.6055	0.2494	0.036*	0.839 (2)
C17	0.1586 (2)	0.6238 (4)	0.32558 (16)	0.0352 (9)	0.839 (2)
C13	0.16533 (13)	0.73299 (18)	0.37901 (7)	0.0524 (5)	0.839 (2)
C14	0.06429 (15)	0.5634 (3)	0.32483 (14)	0.0415 (5)	0.839 (2)
C15	0.22215 (6)	0.49641 (11)	0.34156 (5)	0.0460 (3)	0.839 (2)
C15'	0.1645 (9)	0.8073 (10)	0.2458 (4)	0.032 (3)	0.161 (2)
H15'	0.2176	0.7933	0.2592	0.038*	0.161 (2)
C12'	0.1272 (4)	0.9613 (6)	0.2612 (2)	0.0464 (17)	0.161 (2)
C16'	0.1082 (9)	0.7056 (12)	0.2629 (5)	0.033 (3)	0.161 (2)
H16C	0.0566	0.7432	0.2661	0.040*	0.161 (2)
H16D	0.1058	0.6407	0.2347	0.040*	0.161 (2)
C17'	0.1283 (7)	0.6432 (11)	0.3150 (5)	0.040 (3)	0.161 (2)
C13'	0.1395 (7)	0.7557 (10)	0.3665 (4)	0.062 (3)	0.161 (2)
C14'	0.0539 (8)	0.5400 (16)	0.3321 (8)	0.049 (3)	0.161 (2)
C15'	0.2150 (4)	0.5561 (6)	0.3065 (3)	0.0541 (19)	0.161 (2)
C16	0.3626 (3)	0.9480 (5)	0.1331 (2)	0.0408 (9)	0.634 (17)
O1	0.4211 (4)	1.0019 (10)	0.1000 (3)	0.065 (2)	0.634 (17)
O2	0.3883 (6)	0.9535 (10)	0.1871 (3)	0.061 (2)	0.634 (17)
O3	0.2938 (5)	1.0211 (10)	0.1273 (5)	0.058 (2)	0.634 (17)
O4	0.3463 (6)	0.8219 (6)	0.1179 (4)	0.052 (2)	0.634 (17)
C16'	0.3611 (9)	0.9340 (15)	0.1444 (7)	0.051 (2)	0.221 (16)
O1'	0.4122 (12)	0.951 (2)	0.0995 (8)	0.059 (4)	0.221 (16)
O2'	0.4055 (14)	0.949 (3)	0.1915 (8)	0.056 (4)	0.221 (16)
O3'	0.2990 (12)	1.022 (2)	0.1436 (9)	0.051 (4)	0.221 (16)
O4'	0.3301 (11)	0.8105 (15)	0.1398 (10)	0.047 (4)	0.221 (16)
C16''	0.3552 (10)	0.9434 (19)	0.1397 (7)	0.052 (3)	0.145 (7)
O1''	0.4277 (11)	1.007 (2)	0.1349 (11)	0.061 (4)	0.145 (7)
O2''	0.3490 (14)	0.902 (2)	0.1936 (7)	0.059 (4)	0.145 (7)
O3''	0.2941 (15)	1.027 (4)	0.1259 (16)	0.053 (5)	0.145 (7)
O4''	0.356 (2)	0.843 (3)	0.1021 (10)	0.051 (4)	0.145 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.02551 (19)	0.0343 (2)	0.02575 (19)	-0.00116 (17)	-0.00341 (16)	0.00344 (17)
N1	0.0324 (14)	0.0322 (14)	0.0296 (14)	0.0024 (12)	0.0030 (12)	0.0078 (11)
N2	0.0373 (15)	0.0336 (15)	0.0212 (13)	-0.0058 (12)	-0.0016 (11)	-0.0007 (11)
N3	0.0282 (13)	0.0250 (13)	0.0234 (12)	0.0014 (11)	-0.0012 (10)	-0.0007 (10)
C11	0.0270 (4)	0.0476 (5)	0.0334 (4)	-0.0072 (4)	-0.0093 (3)	0.0132 (4)
C1	0.053 (2)	0.042 (2)	0.0279 (17)	-0.0150 (17)	-0.0050 (16)	-0.0012 (15)
C2	0.076 (3)	0.043 (2)	0.0310 (19)	-0.006 (2)	-0.0020 (19)	-0.0086 (17)
C3	0.075 (3)	0.046 (2)	0.040 (2)	0.007 (2)	0.006 (2)	-0.0131 (18)
C4	0.047 (2)	0.046 (2)	0.039 (2)	0.0101 (18)	0.0055 (17)	-0.0053 (17)
C5	0.0344 (17)	0.0332 (17)	0.0227 (15)	-0.0004 (14)	-0.0009 (13)	-0.0001 (13)
C6	0.0364 (17)	0.0248 (15)	0.0284 (16)	0.0010 (14)	0.0046 (13)	0.0050 (13)

C7	0.053 (2)	0.0288 (17)	0.0361 (19)	-0.0016 (16)	0.0088 (17)	0.0013 (15)
C8	0.068 (3)	0.0301 (19)	0.046 (2)	0.0114 (18)	0.027 (2)	0.0071 (17)
C9	0.043 (2)	0.047 (2)	0.059 (3)	0.0165 (18)	0.023 (2)	0.017 (2)
C10	0.0326 (18)	0.047 (2)	0.044 (2)	0.0082 (16)	0.0095 (15)	0.0172 (18)
C11	0.0317 (16)	0.0243 (15)	0.0297 (16)	-0.0005 (13)	-0.0034 (13)	-0.0013 (13)
C12	0.0277 (16)	0.0302 (16)	0.0268 (16)	0.0015 (13)	-0.0011 (13)	-0.0007 (13)
C13	0.0423 (19)	0.0257 (16)	0.0258 (16)	0.0009 (14)	-0.0019 (14)	0.0022 (13)
C14	0.051 (2)	0.0356 (19)	0.0277 (17)	0.0136 (16)	-0.0045 (15)	-0.0004 (15)
C15	0.030 (2)	0.029 (2)	0.0240 (19)	0.0028 (16)	-0.0020 (16)	-0.0036 (15)
Cl2	0.0539 (7)	0.0435 (7)	0.0319 (6)	0.0184 (6)	0.0005 (5)	-0.0079 (5)
C16	0.0327 (19)	0.0306 (18)	0.0262 (17)	0.0009 (16)	-0.0005 (15)	0.0009 (15)
C17	0.038 (2)	0.033 (2)	0.034 (2)	0.0022 (18)	0.0023 (18)	0.0034 (17)
Cl3	0.0836 (14)	0.0450 (9)	0.0286 (8)	-0.0038 (8)	-0.0016 (7)	-0.0030 (6)
Cl4	0.0385 (8)	0.0382 (11)	0.0478 (11)	-0.0028 (7)	0.0107 (7)	0.0139 (8)
Cl5	0.0458 (6)	0.0461 (6)	0.0462 (7)	0.0073 (5)	-0.0033 (5)	0.0156 (5)
C15'	0.035 (5)	0.033 (5)	0.027 (5)	0.004 (5)	-0.004 (5)	-0.006 (5)
Cl2'	0.069 (4)	0.032 (3)	0.038 (3)	-0.006 (3)	-0.001 (3)	-0.002 (2)
C16'	0.039 (5)	0.033 (5)	0.027 (5)	0.000 (5)	0.001 (5)	-0.006 (5)
C17'	0.046 (5)	0.038 (5)	0.036 (5)	0.002 (5)	0.002 (5)	-0.003 (5)
Cl3'	0.082 (7)	0.061 (5)	0.042 (5)	-0.008 (5)	-0.003 (4)	-0.003 (4)
Cl4'	0.054 (6)	0.043 (6)	0.050 (6)	-0.009 (4)	0.012 (4)	0.019 (4)
Cl5'	0.052 (3)	0.054 (4)	0.057 (4)	0.011 (3)	-0.010 (3)	0.011 (3)
Cl6	0.0335 (13)	0.0350 (13)	0.0539 (17)	0.0105 (10)	-0.0121 (11)	-0.0163 (11)
O1	0.047 (3)	0.068 (5)	0.079 (4)	-0.007 (3)	-0.008 (3)	-0.002 (4)
O2	0.052 (5)	0.073 (4)	0.058 (3)	0.013 (3)	-0.031 (3)	-0.026 (3)
O3	0.039 (3)	0.039 (3)	0.095 (5)	0.013 (3)	-0.028 (3)	-0.029 (3)
O4	0.054 (4)	0.031 (3)	0.070 (5)	0.010 (3)	-0.020 (4)	-0.018 (3)
Cl6'	0.037 (3)	0.040 (3)	0.075 (4)	0.009 (3)	-0.021 (3)	-0.015 (3)
O1'	0.047 (6)	0.058 (7)	0.073 (6)	-0.001 (6)	0.007 (6)	-0.002 (6)
O2'	0.042 (7)	0.063 (7)	0.064 (7)	0.007 (6)	-0.023 (6)	-0.018 (6)
O3'	0.040 (6)	0.042 (6)	0.069 (8)	0.013 (5)	-0.032 (6)	-0.025 (6)
O4'	0.036 (6)	0.034 (6)	0.071 (7)	0.002 (5)	-0.015 (6)	-0.010 (6)
Cl6''	0.043 (4)	0.045 (4)	0.070 (4)	0.001 (4)	-0.018 (4)	-0.017 (4)
O1''	0.047 (6)	0.063 (7)	0.072 (7)	0.004 (6)	-0.016 (6)	-0.013 (7)
O2''	0.047 (7)	0.057 (7)	0.071 (7)	0.008 (6)	-0.012 (6)	-0.021 (6)
O3''	0.041 (8)	0.042 (8)	0.076 (9)	0.010 (8)	-0.021 (8)	-0.020 (8)
O4''	0.049 (7)	0.038 (7)	0.067 (8)	0.008 (7)	-0.013 (7)	-0.024 (7)

Geometric parameters (Å, °)

Cu1—N1	1.982 (3)	C14—C15'	1.527 (10)
Cu1—N2	1.987 (3)	C14—C15	1.557 (5)
Cu1—N3	2.027 (2)	C14—H14A	0.9900
Cu1—C11	2.2519 (8)	C14—H14B	0.9900
N1—C6	1.342 (4)	C14—H14C	0.9900
N1—C10	1.346 (4)	C14—H14D	0.9900
N2—C5	1.350 (4)	C15—C16	1.512 (5)
N2—C1	1.353 (4)	C15—Cl2	1.815 (4)

N3—C12	1.479 (4)	C15—H15	1.0000
N3—C13	1.489 (4)	C16—C17	1.515 (5)
N3—C11	1.492 (4)	C16—H16A	0.9900
C1—C2	1.367 (5)	C16—H16B	0.9900
C1—H1	0.9500	C17—C14	1.771 (5)
C2—C3	1.375 (6)	C17—C13	1.780 (4)
C2—H2	0.9500	C17—C15	1.800 (4)
C3—C4	1.375 (5)	C15'—C16'	1.526 (10)
C3—H3	0.9500	C15'—C12'	1.807 (10)
C4—C5	1.390 (5)	C15'—H15'	1.0000
C4—H4	0.9500	C16'—C17'	1.509 (10)
C5—C12	1.496 (4)	C16'—H16C	0.9900
C6—C7	1.380 (4)	C16'—H16D	0.9900
C6—C11	1.504 (4)	C17'—C14'	1.758 (9)
C7—C8	1.381 (5)	C17'—C13'	1.774 (9)
C7—H7	0.9500	C17'—C15'	1.790 (10)
C8—C9	1.377 (6)	C16—O4	1.427 (4)
C8—H8	0.9500	C16—O2	1.429 (5)
C9—C10	1.378 (5)	C16—O1	1.438 (6)
C9—H9	0.9500	C16—O3	1.442 (4)
C10—H10	0.9500	C16'—O2'	1.424 (8)
C11—H11A	0.9900	C16'—O4'	1.429 (8)
C11—H11B	0.9900	C16'—O3'	1.434 (8)
C12—H12A	0.9900	C16'—O1'	1.449 (9)
C12—H12B	0.9900	C16"—O2"	1.427 (9)
C13—C14	1.530 (4)	C16"—O4"	1.429 (8)
C13—H13A	0.9900	C16"—O3"	1.436 (8)
C13—H13B	0.9900	C16"—O1"	1.444 (9)
N1—Cu1—N2	165.22 (11)	C15'—C14—C13	130.4 (6)
N1—Cu1—N3	83.05 (10)	C13—C14—C15	108.2 (3)
N2—Cu1—N3	82.56 (10)	C13—C14—H14A	110.0
N1—Cu1—C11	96.74 (8)	C15—C14—H14A	110.0
N2—Cu1—C11	97.22 (8)	C13—C14—H14B	110.0
N3—Cu1—C11	173.97 (8)	C15—C14—H14B	110.0
C6—N1—C10	119.3 (3)	H14A—C14—H14B	108.4
C6—N1—Cu1	113.2 (2)	C15'—C14—H14C	104.7
C10—N1—Cu1	127.5 (3)	C13—C14—H14C	104.7
C5—N2—C1	119.2 (3)	C15'—C14—H14D	104.7
C5—N2—Cu1	112.6 (2)	C13—C14—H14D	104.7
C1—N2—Cu1	128.2 (2)	H14C—C14—H14D	105.7
C12—N3—C13	112.2 (2)	C16—C15—C14	110.0 (3)
C12—N3—C11	114.7 (2)	C16—C15—C12	112.1 (3)
C13—N3—C11	110.4 (2)	C14—C15—C12	107.2 (3)
C12—N3—Cu1	105.74 (18)	C16—C15—H15	109.2
C13—N3—Cu1	107.54 (18)	C14—C15—H15	109.2
C11—N3—Cu1	105.65 (18)	C12—C15—H15	109.2
N2—C1—C2	121.4 (4)	C15—C16—C17	117.6 (3)

N2—C1—H1	119.3	C15—C16—H16A	107.9
C2—C1—H1	119.3	C17—C16—H16A	107.9
C1—C2—C3	119.8 (4)	C15—C16—H16B	107.9
C1—C2—H2	120.1	C17—C16—H16B	107.9
C3—C2—H2	120.1	H16A—C16—H16B	107.2
C2—C3—C4	119.5 (4)	C16—C17—C14	112.3 (3)
C2—C3—H3	120.2	C16—C17—C13	114.1 (3)
C4—C3—H3	120.2	C14—C17—C13	107.9 (2)
C3—C4—C5	118.9 (4)	C16—C17—C15	108.0 (3)
C3—C4—H4	120.6	C14—C17—C15	107.6 (2)
C5—C4—H4	120.6	C13—C17—C15	106.6 (2)
N2—C5—C4	121.2 (3)	C16'—C15'—C14	97.1 (8)
N2—C5—C12	116.0 (3)	C16'—C15'—C12'	110.7 (9)
C4—C5—C12	122.7 (3)	C14—C15'—C12'	107.2 (7)
N1—C6—C7	121.3 (3)	C16'—C15'—H15'	113.5
N1—C6—C11	115.9 (3)	C14—C15'—H15'	113.5
C7—C6—C11	122.8 (3)	C12'—C15'—H15'	113.5
C6—C7—C8	119.3 (4)	C17'—C16'—C15'	113.9 (9)
C6—C7—H7	120.3	C17'—C16'—H16C	108.8
C8—C7—H7	120.3	C15'—C16'—H16C	108.8
C9—C8—C7	119.2 (4)	C17'—C16'—H16D	108.8
C9—C8—H8	120.4	C15'—C16'—H16D	108.8
C7—C8—H8	120.4	H16C—C16'—H16D	107.7
C8—C9—C10	118.9 (3)	C16'—C17'—C14'	108.3 (9)
C8—C9—H9	120.5	C16'—C17'—C13'	111.0 (8)
C10—C9—H9	120.5	C14'—C17'—C13'	109.0 (9)
N1—C10—C9	121.8 (4)	C16'—C17'—C15'	108.9 (8)
N1—C10—H10	119.1	C14'—C17'—C15'	109.4 (8)
C9—C10—H10	119.1	C13'—C17'—C15'	110.2 (8)
N3—C11—C6	108.4 (2)	O4—C16—O2	110.8 (5)
N3—C11—H11A	110.0	O4—C16—O1	111.4 (4)
C6—C11—H11A	110.0	O2—C16—O1	108.0 (5)
N3—C11—H11B	110.0	O4—C16—O3	108.3 (4)
C6—C11—H11B	110.0	O2—C16—O3	109.7 (5)
H11A—C11—H11B	108.4	O1—C16—O3	108.6 (5)
N3—C12—C5	109.2 (2)	O2'—C16'—O4'	112.0 (10)
N3—C12—H12A	109.8	O2'—C16'—O3'	110.6 (10)
C5—C12—H12A	109.8	O4'—C16'—O3'	108.3 (10)
N3—C12—H12B	109.8	O2'—C16'—O1'	107.2 (10)
C5—C12—H12B	109.8	O4'—C16'—O1'	106.7 (10)
H12A—C12—H12B	108.3	O3'—C16'—O1'	111.9 (10)
N3—C13—C14	114.6 (3)	O2''—C16''—O4''	113.4 (12)
N3—C13—H13A	108.6	O2''—C16''—O3''	111.3 (12)
C14—C13—H13A	108.6	O4''—C16''—O3''	108.4 (11)
N3—C13—H13B	108.6	O2''—C16''—O1''	106.9 (11)
C14—C13—H13B	108.6	O4''—C16''—O1''	106.9 (11)
H13A—C13—H13B	107.6	O3''—C16''—O1''	109.8 (12)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C11—H11 <i>A</i> \cdots C11 ⁱ	0.99	2.84	3.441 (3)	120
C12—H12 <i>B</i> \cdots C11 ⁱ	0.99	2.84	3.438 (3)	120
C2—H2 \cdots O1 ⁱⁱ	0.95	2.64	3.383 (8)	136
C11—H11 <i>A</i> \cdots O4 ⁱⁱⁱ	0.99	2.53	3.123 (7)	118
C12—H12 <i>A</i> \cdots O4	0.99	2.43	3.310 (9)	148
C13—H13 <i>A</i> \cdots O1 ^{iv}	0.99	2.64	3.578 (11)	157
C14—H14 <i>B</i> \cdots O3 ^{iv}	0.99	2.40	3.324 (13)	154
C7—H7 \cdots O2 ^{viii}	0.95	2.64	3.14 (3)	113
C12—H12 <i>A</i> \cdots O4'	0.99	2.37	3.32 (2)	159
C14—H14 <i>C</i> \cdots O3 ^{iv}	0.99	2.17	3.13 (2)	163
C14—H14 <i>D</i> \cdots O4'	0.99	2.41	3.26 (2)	143
C16'—H16 <i>D</i> \cdots O2 ^{iv}	0.99	2.32	3.28 (3)	162
C7—H7 \cdots O2 ⁱⁱⁱ	0.95	2.57	3.26 (2)	130
C12—H12 <i>A</i> \cdots O4''	0.99	2.53	3.33 (4)	138
C13—H13 <i>A</i> \cdots O1 ^{iv}	0.99	2.45	3.33 (2)	148
C2—H2 \cdots C14 ^v	0.95	2.84	3.644 (19)	143
C11—H11 <i>B</i> \cdots C12'	0.99	2.70	3.490 (7)	137
C13—H13 <i>B</i> \cdots C12 ^{vi}	0.99	2.90	3.863 (4)	165

Symmetry codes: (i) $-x, -y+2, -z$; (ii) $x-1/2, -y+3/2, -z$; (iii) $-x+1/2, y+1/2, z$; (iv) $-x+1/2, y-1/2, z$; (v) $x, -y+1, z-1/2$; (vi) $-x, y, -z+1/2$.