

Received 12 August 2015

Accepted 27 August 2015

Edited by A. J. Lough, University of Toronto,
Canada

Keywords: crystal structure; 2-oxazolines; bromoaryl; iodoaryl; Phebox ligands; hydrogen bonding; parallel-displaced π - π interaction; N \cdots I contacts; isostructural compounds

CCDC references: 1420839; 1420838

Supporting information: this article has supporting information at journals.iucr.org/e

Crystal structures of 1-bromo-3,5-bis(4,4-dimethyl-1,3-oxazolin-2-yl)benzene 0.15-hydrate and 3,5-bis(4,4-dimethyl-1,3-oxazolin-2-yl)-1-iodobenzene

Timo Stein, Frank Hoffmann and Michael Fröba*

Institute of Inorganic and Applied Chemistry, Department of Chemistry, University of Hamburg, Martin-Luther-King-Platz 6, D-20146 Hamburg, Germany. *Correspondence e-mail: michael.froeba@chemie.uni-hamburg.de



The bromo and iodo derivatives of a *meta*-bis(1,3-oxazolin-2-yl)-substituted benzene, C₁₆H₁₉BrN₂O₂·0.15H₂O (**1**) and C₁₆H₁₉IN₂O₂ (**2**), have been prepared and studied in terms of their molecular and crystal structures. While the former crystallizes as a sub-hydrate, with 0.15 formula units of water and shows an almost all-planar arrangement of the three ring systems, the latter crystallizes solvate-free with the flanking heterocycles twisted considerably with respect to the central arene. Non-covalent contacts include parallel-displaced π - π interactions and (non-classical) hydrogen bonding for both (**1**) and (**2**), as well as relatively short I \cdots N contacts for (**2**).

1. Chemical context

The 2-oxazolinyll functional group has been employed as a protective group for carboxylic acids rendering them stable against organometallic reagents (Wuts & Greene, 2007). Aromatic 1,3-substituted bis(1,3-oxazolin-2-yl) compounds have shown to be efficient for directed *ortho* metallation (DoM) reactions (Harris *et al.*, 1978), while competitive halogen-metal exchange reactions should be considered for the halide-substituted title compounds reported herein. The substitutional pattern also gives access to bis(1,3-oxazolin-2-yl) systems suitable for the preparation of *N,C,N*-tridentate pincer ligands which have come to general attention as Phebox ligands [Phebox: 2,6-bis(1,3-oxazolin-2-yl)phenyl].

There are many examples for respective organometallic complexes of transition and rare-earth metals known in the literature with many of them bearing chiral 2-oxazolinyll substituents within the Phebox ligand. Exemplary compounds include those of some early transition metals (Chuchuryukin *et al.*, 2011), iridium (Allen *et al.*, 2014), or palladium (Lu *et al.*, 2010), to name but a few.

The substitution with bromine and iodine renders the title compounds capable of being utilized as potential precursors for C-C cross-coupling building blocks bearing the Phebox motif.






Figure 1
The asymmetric unit of **(1)**, shown with 50% probability displacement ellipsoids. H atoms are drawn as green spheres of an arbitrary radius.

2. Structural commentary

The structural considerations below take least-squares mean planes for rings *A* (O1/C7/N1/C9/C8) and *C* (O2/C12/N2/C14/C13) of the heterocyclic 2-oxazolynyl groups and *B* (C1/C2/C3/C4/C5/C6) for the central arene into account. For both **(1)** and **(2)**, the 2-oxazoline moiety with an approximately accordant orientation of the C=N bond with respect to the C5–X1 (*X* = Br, I) bond was chosen to be denoted as ring *A*.

Compound **(1)** crystallized from water-containing acetonitrile as an adduct with 0.15 H₂O in the monoclinic space group *P*2₁/*n*. Within the molecular structure (Fig. 1) the 2-oxazolynyl functional groups are oriented antiperiplanar to each other. The elongated displacement ellipsoid of the methylene carbon atom C13 indicates a noteworthy vibrational freedom orthogonal to plane *C* which was not observed for atoms within ring *A*. The absence of a comparable displacement component of C14 perpendicular to plane *C* disagrees with pseudorotational disorder around the atomic positions mentioned.

The five-membered heterocyclic moieties possess different conformational characteristics. Within ring *A*, significant puckering with parameters $\tau_m = 13.2$ (1)°, $q_2 = 0.1284$ (19) Å, and $\varphi_2 = 134.2$ (9)° indicate a ^{C8}*T*_{C9} conformation distorted towards ^{C8}*E* (Altona & Sundaralingam, 1972; Cremer & Pople, 1975). Ring *C* shows only slight deviation from ideal planarity with $\tau_m = 3.5$ (1)°.

Interplanar angles of 2.15 (12)° and 3.67 (16)° of the planes N1/C7/O1 and N2/C12/O2 with plane *B*, respectively, have been found for the almost all-planar overall structure. Considerable angular deviations from ideal (120°) angles within ring *B* were found for C4–C5–C6 = 122.04 (15)°, C3–C4–C5 = 118.75 (15), and C1–C6–C5 = 118.59 (14)°.

The water molecule is involved in hydrogen bonds with N1···H3A = 1.96 (2) Å and O3···H6 = 2.548 (9) Å, where the corresponding angles are O3–H3A···N1 = 178 (15)° and C6–H6···O3 = 148.6 (3)°, respectively. Hydrogen-bond geometries for **(1)** are summarized in Table 1.

Table 1
Hydrogen-bond geometry (Å, °) for **(1)**.

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
C6–H6···O3	0.95	2.55 (1)	3.395 (10)	149 (1)
O3–H3A···N1	0.84 (2)	1.96 (2)	2.796 (10)	178 (15)
C16–H16B···O3 ⁱ	0.98	2.18 (1)	3.045 (11)	146 (1)
C13–H13B···O3 ⁱ	0.99	2.94 (1)	3.656 (12)	130 (1)
C8–H8B···Br1 ⁱⁱ	0.99	3.09 (1)	4.054 (2)	165 (1)
C11–H11C···O3 ⁱⁱⁱ	0.98	2.56 (1)	3.391 (12)	143 (1)
O3–H3B···N1 ⁱⁱⁱ	0.84 (2)	2.37 (9)	3.107 (11)	148 (15)
O3–H3B···O3 ⁱⁱⁱ	0.84 (2)	2.32 (13)	2.90 (2)	126 (13)

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x - 1, y, z$; (iii) $-x, -y + 1, -z$.

The isostructural iodo derivative **(2)** was crystallized from dichloromethane with no evidence of co-crystallized solvent. Again, an antiperiplanar orientation of the 2-oxazolynyl moieties within the molecular structure (Fig. 2) was found. In contrast to **(1)**, a more distinct twisting of the planes N1/C7/O1 and N2/C12/O2 compared to the plane *B* with interplanar angles of 16.16 (4) and 15.14 (4)°, respectively, was found.

For **(2)**, similar conformational considerations as for **(1)** apply. Puckering parameters $\tau_m = 19.6$ (1)°, $q_2 = 0.1902$ (11) Å, and $\varphi_2 = 135.5$ (3)° indicate a conformation between ^{C8}*T*_{C9} and ^{C8}*E* for ring *C*. A higher degree of planarity was found for ring *A* with $\tau_m = 4.6$ (1)°. The bond length C5–I1 is 2.0968 (9) Å with I1 located 0.1183 (14) Å above ring *B* and thus considerably more distant than the bromine atom in **(1)**, where Br1 lies only 0.005 (2) Å above plane *B*. The deviation occurs away from a π – π stacked (see also below) inversion-equivalent formula unit and might be the consequence of steric repulsion between the bulky iodine substituent and the 2-oxazolynyl groups of the neighboring formula unit. Noteworthy deviations from ideal angles within ring *B* were found with C4–C5–C6 = 121.35 (9)° and C3–C4–C5 = 118.60 (9)°. I1 shows an angular adjustment to C6–C5–I1 = 117.08 (7)°, thus improving intermolecular N···I contacts (see below).




Figure 2
The asymmetric unit of **(2)**, shown with displacement ellipsoids drawn at the 50% probability level. H atoms are drawn as green spheres of an arbitrary radius.




Figure 3
View along the *a* axis of the crystal packing of compound (1).

All bond lengths for (1) and (2) fall within expected ranges (Allen *et al.*, 1987).

3. Supramolecular features

Within the crystal structure of (1) (Fig. 3), intermolecular bonding is established by classical and non-classical hydrogen




Figure 4
Intermolecular contacts within the crystal structure of (1) are established by means of parallel-displaced π - π interactions and hydrogen bonding. Displacement ellipsoids are at the 50% probability level and intermolecular contacts are depicted as dashed lines. Only H atoms involved in hydrogen bonding or van der Waals contacts (black dashed lines) are shown as green spheres at an arbitrary radius. Purple dashed lines indicate centroid-centroid connecting lines. [Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x - \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$]




Figure 5
Supramolecular features within the crystal structure of (2) comprise π - π contacts and mutual $N1 \cdots I1^{ii}$ and $N1^{ii} \cdots I1$ interactions. Moreover, there are non-conventional hydrogen bonds $I1^{ii} \cdots H6$, $I1 \cdots H6^{ii}$, and $H10B \cdots N2^{iii}$. Intermolecular contacts are shown as dashed lines. Only atoms H6, $H6^{ii}$, and H10B which are involved in hydrogen bonding (black dashed lines) are shown as green spheres of an arbitrary radius. Purple dashed lines connect centroids of π - π associated dimers. [Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 1, -y + 2, -z + 1$; (iii) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$]

bonding (see Table 1) as well as π - π interactions. Close contacts $O3 \cdots H16B^i = 2.181(10)$ Å with $C16^i - H16B^i \cdots O3 = 146.2(3)^\circ$ and $H13B \cdots O3^i = 2.936(11)$ Å with $C13 - H13B \cdots O3^i = 130.4(3)^\circ$ were found. Further hydrogen-bonding interactions arise from mutual interactions of two inversion-related water molecules with $H3B \cdots O3^{iii} = 2.32(13)$ Å and $O3 - H3B \cdots O3^{iii} = 126(13)^\circ$.

Additionally, parallel-displaced π - π stacking with $Cg1 \cdots Cg1^i = 3.5064(12)$ Å (*Cg1* denotes the centroid calculated for the arene atoms) and a horizontal displacement of $I = 0.996(2)$ Å (for a description of geometric parameters of π - π associated arenes, see: Snyder *et al.*, 2012) between the centroids was found. The interplanar distance is $R = 3.3620(13)$ Å. The π - π interaction gives rise to antiperiplanar dimers linked to each other by an inversion. Features of intermolecular bonding for (1) are illustrated in Fig. 4.

The supramolecular contacts found for (2) are depicted in Fig. 5. A comparable π - π stacking motif as for compound (1) was found within the crystal structure of (2) (Fig. 6). The centroid-centroid distance $Cg1 \cdots Cg1^i = 3.6142(8)$ Å is slightly increased compared to (1). The interplanar distance is $R = 3.2862(9)$ Å with a remarkably increased horizontal centroid-centroid displacement of $I = 1.5044(15)$ Å.

Mutual hydrogen bonds $I1 \cdots H6^{ii} = I1^{ii} \cdots H6 = 3.11$ Å with $C6 - H6 \cdots I1^{ii} = C6^{ii} - H6^i \cdots I1 = 150^\circ$ and $H10B \cdots N2^{ii} =$




Figure 6
View along the *a* axis of the crystal packing of compound **(2)**.

2.745 Å with an angle of $\text{C10-H10B}\cdots\text{N2}^{\text{ii}} = 172^\circ$ establish intermolecular bonding (Table 2). Additionally, particularly short mutual $\text{N1}\cdots\text{I1}^{\text{ii}}$ and $\text{N1}^{\text{ii}}\cdots\text{I1}$ contacts at 3.2779 (9) Å were found. The $\text{N}\cdots\text{I}$ distance corresponds to 89% of the sum of the van der Waals radii (3.70 Å; Alvarez, 2013) of the atoms involved. To the best of our knowledge, the shortest $\text{N}\cdots\text{I}$ contact found in a crystal structure was recently reported to be 2.622 Å (Bosch, 2014), corresponding to 71% of the sum of the van der Waals radii. With respect to $\text{I1}\cdots\text{N1}^{\text{ii}}$ contacts, the angle $\text{C5-I1}\cdots\text{N1}^{\text{ii}} = 155.61 (3)^\circ$ is found to be slightly linearized by an angular adaption C6-C5-I1 (see above).

The halogen-halogen distances for **(1)** and **(2)** exceed the sum of the van der Waals radii and lack appropriate angles for halogen-halogen bonding.

4. Database survey

The crystal structure of the reagent used to prepare compounds **(1)** and **(2)** (see also below), 5-bromo-1,3-dicyanobenzene, has been published recently (CCDC reference number DIRLEX; Seidel *et al.*, 2013).

A WebCSD search (Version 1.1.1, updated July 2015; Groom & Allen, 2014) for the 1,3-bis(1,3-oxazolin-2-yl)benzene substructure gave 143 hits. Very few of the crystal structures show the parent motif without any metal ion bound to it. The following considerations take into account only purely organic structures with a 1,3-substitutional pattern, although a significant number of (metal coordinated) 1,3,5-tri(1,3-oxazolin-2-yl)benzenes has been reported.

1-Iodo-2,6-bis(4'-isopropyl-1,3-oxazolin-2-yl)-4-*tert*-butylbenzene (ROHMIL; Bugarin & Connell, 2008) and 1,3-bis(4,4-dimethyl-1,3-oxazolin-2-yl)-2-(trimethylstannyl)benzene (FILNAQ; Stol *et al.*, 2005) show substitutional variation from compounds **(1)** and **(2)**. By introducing sterically demanding substituents *ortho* to the 1,3-oxazolin-2-yl groups, these are considerably more twisted against the parent arene ring plane with twisting angles ranging from 47.3 (2)–63.6 (2)°

Table 2
Hydrogen-bond geometry (Å, °) for **(2)**.

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
C6–H6···I1 ⁱ	0.95	3.11	3.9679 (9)	150
C10–H10B···N2 ⁱⁱ	0.98	2.75	3.7228 (15)	172

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

and 22.35 (8)–22.75 (8)° for the iodo and tin derivatives, respectively. The more distinct twisting of the respective groups in **(2)** with 16.16 (4) and 15.14 (4)° compared to **(1)** [2.15 (12) and 3.67 (16)°] might be attributed to geometrical requirements for the formation of supramolecular bonding observed for **(2)**. 1-Iodo-2,6-bis(4'-isopropyl-1,3-oxazolin-2-yl)-4-*tert*-butylbenzene shows short $\text{N}\cdots\text{I}$ distances at 3.041 (6) Å corresponding to 82% of the sum of van der Waals radii.


Other structures generally lack halide substitution at the central arene, although substitutional variety at the 4'-positions was found. Instead of 4,4-dimethyl substitution, derivatives bearing hydrogen atoms (LAFNEM; Chen *et al.*, 2004), hydroxymethyl (MODQIH; Javadi *et al.*, 2014) or isopropyl groups (DOWGOM; Mei *et al.*, 2009) have been found. For those structures, no features of intermolecular π – π interactions could be observed. Instead, $\text{C-H}\cdots\pi$ interaction becomes obvious for LAFNEM and DOWGOM. It is supposed that halide substitution increases dispersion interaction at modest horizontal centroid-centroid separations (Arnstein & Sherrill, 2008), thus promoting π – π stacking for **(1)** and **(2)**. As a result of the presence of electronegative atoms, hydrogen bonding and van der Waals contacts can also be found for the structures mentioned.

Structures DOWGOM, FILNAQ, MODQIH, and ROHMIL show a synperiplanar orientation of the 2-oxazolinyl groups. The antiperiplanar arrangement as in **(1)** and **(2)** was only found for LAFNEM.

Langer *et al.* found the same ${}^{\text{C8}}T_{\text{C9}}$ conformation (with respect to the numbering scheme used herein) within the 2-oxazoline ring of 2-(4-hydroxyphenyl)-4,4-dimethyl-2-oxazoline (CELMAI; Langer *et al.*, 2006) as for the 2-oxazoline moieties in **(1)** and **(2)**. Conformational analysis of the 2-oxazolines within the structures discussed herein showed that all possible conformations twisted around C8–C9 can be found, that are ${}^{\text{C8}}T_{\text{C9}}$ (two times within DOWGOM, CELMAI) and ${}^{\text{C9}}T_{\text{C8}}$ (LAWCAO, ROHMIL). Folding into an envelope conformation was only observed at C8 with examples for ${}^{\text{C8}}E$ (two times within DOWGOM, ROHMIL) and E_{C8} (FILNAQ, two times within ROHMIL) conformations. This might suggest that folding happens preferentially at the less-substituted methylene position. Planar conformations have been found for MODQIH, LAFNEM, and FILNAQ.

5. Synthesis and crystallization

(1) and **(2)** were synthesized starting from 5-bromo-1,3-dicyanobenzene (Fig. 7). The multi-step preparation of


Figure 7

The synthesis scheme of **(1)** and **(2)**, starting from 1-bromo-3,5-dicyanobenzene. The numbering scheme for the title compounds is given.

5-bromo-1,3-dicyanobenzene starting from isophthalic acid has been described in a patent (Dillard *et al.*, 2010). For the preparation of compound **(1)**, 5-bromo-1,3-dicyanobenzene was subjected to cyclization at the cyano positions with 2-amino-2-methylpropan-1-ol under zinc(II) catalysis (Button *et al.*, 2002) to give the *meta*-bis(1,3-oxazolin-2-yl) arene. The iodo derivative **(2)** was synthesized from **(1)** by an aromatic Finkelstein reaction (Klapars & Buchwald, 2002). All reactions were carried out under an atmosphere of dry nitrogen using standard Schlenk techniques.

5.1. 1-Bromo-3,5-bis(4,4-dimethyl-1,3-oxazolin-2-yl)benzene sub-hydrate (**1**)

Zinc(II) chloride (73 mg, 0.54 mmol, 0.1 eq.) was melted three times *in vacuo* with help of a heat gun. 5-Bromo-1,3-dicyanobenzene (1.13 g, 4.65 mmol, 1.0 eq.), 2-amino-2-methylpropan-1-ol (870 mg, 9.77 mmol, 2.1 eq.) and chlorobenzene (15 mL) were added, the colorless suspension was magnetically stirred and heated to reflux for 48 h where it became a pink-colored solution. After quantitative conversion of the dicyano compound was confirmed *via* TLC, the reaction mixture was cooled to 353 K. Volatiles were removed *in vacuo* at 353–433 K. After cooling to room temperature, dichloromethane (50 mL) and distilled water (50 mL) were added. The organic layer was separated, the aqueous layer extracted with dichloromethane (three times with 20 mL each). The combined organic layers were dried over sodium sulfate, filtered, and the solvent was removed from the filtrate under reduced pressure. Column chromatography of the crude product (SiO₂; petrol ether/ethyl acetate = 4:1) gave **(1)** (1.52 g, 4.33 mmol, 93%) as a colorless solid. Crystallization of **(1)** from a concentrated solution in acetonitrile at 277 K gave single crystals suitable for X-ray structural analysis.

¹H-NMR (DMSO-*d*₆, 300.21 MHz): δ (p.p.m.) = 8.28 (*t*, 1H; ⁴*J*(H,H) = 1.5 Hz, H2), 8.06 (*d*, 2H; ⁴*J*(H,H) = 1.5 Hz, H4, H6), 4.16 (*s*, 4H; H8, H13), 1.30 (*s*, 12H; H10, H11, H15, H16). ¹³C{¹H}-NMR (DMSO-*d*₆, 75.50 MHz): δ (p.p.m.) = 158.7 (C7, C12), 132.5 (C4, C6), 130.1 (C1, C3), 126.0 (C2), 121.9 (C5), 78.9 (C8, C13), 67.7 (C9, C14), 28.1 (C10, C11, C15, C16). ESI–

HRMS(+) *m/z*: calculated for [M+H]⁺ 351.0703/353.0688, found 351.0707/353.0689 (⁷⁹Br/⁸¹Br).

5.2. 1-Iodo-3,5-bis(4,4-dimethyl-1,3-oxazolin-2-yl)benzene (**2**)

A *J. Young* ampoule was charged with **(1)** (500 mg, 1.42 mmol, 1.0 eq.), copper(I) iodide (14 mg, 74 μ mol, 5 mol%), and sodium iodide (426 mg, 2.84 mmol, 2.0 eq.). The ampoule was put to vacuum and flushed with nitrogen three times. 1,4-Dioxane (3 mL) and *N,N'*-dimethylethylenediamine (15 μ L, 0.14 mmol, 10 mol%) were added. The ampoule was sealed with a Teflon screw valve, the colorless suspension was magnetically stirred and heated to 373 K for 24 h. An intense blue coloring was observed during the reaction course. After cooling to room temperature, ammonia solution (10 mL, 25%), distilled water (20 mL), and dichloromethane (30 mL) were added. The organic layer was separated, the aqueous layer extracted with dichloromethane (three times with 10 mL each) and the combined organic phases were dried over sodium sulfate. After filtration, dichloromethane was removed from the filtrate under reduced pressure. The crude product was purified by means of column chromatography (SiO₂; petrol ether/ethyl acetate = 4:1) to give a colorless solid **(2)** (505 mg, 1.27 mmol, 89%). Crystallization from a concentrated solution of **(2)** in dichloromethane by slow evaporation of the solvent yielded single crystals suitable for X-ray single crystal diffraction.

¹H-NMR (DMSO-*d*₆, 300.21 MHz): δ (p.p.m.) = 8.28 (*t*, 1H; ⁴*J*(H,H) = 1.5 Hz, H2), 8.24 (*d*, 2H; ⁴*J*(H,H) = 1.5 Hz, H4, H6), 4.14 (*s*, 4H; H8, H13), 1.29 (*s*, 12H; H10, H11, H15, H16). ¹³C{¹H}-NMR (DMSO-*d*₆, 75.50 MHz): δ (p.p.m.) = 158.6 (C7, C12), 138.2 (C1, C3), 132.5 (C4, C6), 129.8 (C2), 126.2 (C1), 78.8 (C8, C13), 67.7 (C9, C14), 28.1 (C10, C11, C15, C16). EI–MS *m/z*: calcd. for *M*⁺ 398.08, found 398.08 (*M*⁺), 383.06 (*M*⁺ – CH₃), 368.07 (*M*⁺ – 2CH₃).

6. Refinement

Crystal data, data collection, and structure refinement details are summarized in Table 3.

Primary atom site locations were assigned with *EDMA* (Palatinus *et al.*, 2012) from electron densities obtained by *SUPERFLIP* (Palatinus & Chapuis, 2007). The remaining secondary non-carbon atom sites were located from the difference Fourier map. All non-hydrogen atoms were refined with anisotropic displacement parameters.

Full-matrix least-squares refinement on *F*² was done with *SHELXL2014/7* (Sheldrick, 2008). Carbon-bound hydrogen atoms were positioned geometrically and refined riding on their respective carbon atom. Bond lengths were fixed at 0.95 Å (aromatic H), 0.98 Å (methyl H), and 0.99 Å (methylene H). *U*_{iso}(H) was fixed at 1.5 *U*_{eq}(O) and *U*_{eq}(C) for hydroxyl and methyl hydrogens or 1.2 *U*_{eq}(C) for the remaining hydrogens. Methyl hydrogens were fitted to the experimental electron density by allowing them to rotate around the C–C bond with a fixed angle (HFIX 137).

Table 3
Experimental details.

	(1)	(2)
Crystal data		
Chemical formula	C ₁₆ H ₁₉ BrN ₂ O ₂ ·0.15H ₂ O	C ₁₆ H ₁₉ IN ₂ O ₂
<i>M_r</i>	353.94	398.23
Crystal system, space group	Monoclinic, <i>P</i> ₂ ₁ / <i>n</i>	Monoclinic, <i>P</i> ₂ ₁ / <i>n</i>
Temperature (K)	100	100
<i>a</i> , <i>b</i> , <i>c</i> (Å)	10.0661 (1), 16.2960 (2), 11.0400 (1)	9.6195 (2), 9.9759 (2), 17.2951 (4)
β (°)	114.496 (2)	94.648 (1)
<i>V</i> (Å ³)	1647.96 (4)	1654.23 (6)
<i>Z</i>	4	4
Radiation type	Mo <i>K</i> α	Mo <i>K</i> α
μ (mm ⁻¹)	2.50	1.94
Crystal size (mm)	0.34 × 0.26 × 0.08	0.22 × 0.12 × 0.08
Data collection		
Diffractometer	Agilent SuperNova Dual Source diffractometer with an Atlas detector	Bruker SMART APEX CCD area-detector diffractometer
Absorption correction	Multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2013)	Numerical (<i>SADABS</i> ; Bruker, 2014)
<i>T</i> _{min} , <i>T</i> _{max}	0.670, 1.000	0.649, 0.747
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	29541, 5438, 4598	45346, 6463, 6035
<i>R</i> _{int}	0.035	0.019
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.735	0.776
Refinement		
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.035, 0.080, 1.05	0.017, 0.043, 1.07
No. of reflections	5438	6463
No. of parameters	209	194
No. of restraints	3	0
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement	H-atom parameters constrained
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	0.85, -0.74	0.82, -0.62

Computer programs: *CrysAlis PRO* (Agilent, 2013), *APEX2* and *SAINTE* (Bruker, 2014), *SUPERFLIP* (Palatinus & Chapuis, 2007), *SHELXL2014* (Sheldrick, 2015), *XP* in *SHELXTL* (Sheldrick, 2008), *WinGX* (Farrugia, 2012), *OLEX2* (Dolomanov *et al.*, 2009), *pubCIF* (Westrip, 2010) and *PLATON* (Spek, 2009).

For (1), occupancy refinement of O3 gave an occupancy of 0.15543 (611) which was subsequently fixed at 0.15. The hydrogen atoms H3A and H3B were located with the help of CALC-OH, which is the *WinGX* (Farrugia, 2012) implementation of Nardelli's method (Nardelli, 1999) of OH atom positioning. The coordinates of those hydrogen atoms were refined freely while applying restraints to the overall water geometry (O–H = 0.84 Å and H...H = 1.328 Å).

Acknowledgements

The authors gratefully acknowledge the continuous support of Professor Ulrich Behrens, University of Hamburg, for helpful discussions concerning X-ray structural analysis.

References

Agilent Technologies (2013). *CrysAlis PRO*. Agilent Technologies, Santa Clara, California, USA.
 Allen, K. E., Heinekey, D. M., Goldman, A. S. & Goldberg, K. I. (2014). *Organometallics*, **33**, 1337–1340.
 Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
 Altona, C. & Sundaralingam, M. (1972). *J. Am. Chem. Soc.* **94**, 8205–8212.
 Alvarez, S. (2013). *Dalton Trans.* **42**, 8617–8636.
 Arnstein, S. A. & Sherrill, C. D. (2008). *Phys. Chem. Chem. Phys.* **10**, 2581–2583.

Bosch, E. (2014). *Cryst. Growth Des.* **14**, 126–130.
 Bruker (2014). *APEX2*, *SAINTE* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bugarin, A. & Connell, B. T. (2008). *Organometallics*, **27**, 4357–4369.
 Button, K. M., Gossage, R. A. & Phillips, R. K. R. (2002). *Synth. Commun.* **32**, 363–368.
 Chen, S.-H., Li, S.-F., Zou, Y., Yang, L., Chen, B. & Zhu, H.-L. (2004). *Z. Kristallogr. New Cryst. Struct.* **219**, 153–154.
 Chuchuryukin, A. V., Huang, R., Lutz, M., Chadwick, J. C., Spek, A. L. & van Koten, G. (2011). *Organometallics*, **30**, 2819–2830.
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
 Dillard, L. W., Yuan, J., Jia, L. & Zheng, Y. (2010). WO Patent No. 021680 A2.
 Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
 Groom, C. R. & Allen, F. H. (2014). *Angew. Chem. Int. Ed.* **53**, 662–671.
 Harris, T. D., Neuschwander, B. & Boekelheide, V. (1978). *J. Org. Chem.* **43**, 727–730.
 Javadi, M. M., Moghadam, M., Mohammadpoor-Baltork, I., Tangestaninejad, S., Mirkhani, V., Kargar, H. & Tahir, M. N. (2014). *Polyhedron*, **72**, 19–26.
 Klapsars, A. & Buchwald, S. L. (2002). *J. Am. Chem. Soc.* **124**, 14844–14845.
 Langer, V., Gyepesová, D., Scholtzová, E., Lustoň, J., Kronek, J. & Koš, M. (2006). *Acta Cryst.* **C62**, o416–o418.
 Lu, Z.-L., Yang, X.-S., Guo, Z.-F. & Wang, R.-Y. (2010). *J. Coord. Chem.* **63**, 2659–2672.
 Mei, L., Hai, Z. J., Jie, S., Ming, Z. S., Hao, Y. & Liang, H. K. (2009). *J. Comb. Chem.* **11**, 220–227.

- Nardelli, M. (1999). *J. Appl. Cryst.* **32**, 563–571.
- Palatinus, L. & Chapuis, G. (2007). *J. Appl. Cryst.* **40**, 786–790.
- Palatinus, L., Prathapa, S. J. & van Smaalen, S. (2012). *J. Appl. Cryst.* **45**, 575–580.
- Seidel, N., Seichter, W. & Weber, E. (2013). *Acta Cryst.* **E69**, o1732–o1733.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Snyder, S. E., Huang, B.-S., Chu, Y. W., Lin, H.-S. & Carey, J. R. (2012). *Chem. Eur. J.* **18**, 12663–12671.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Stol, M., Snelders, D. J. M., de Pater, J. J. M., van Klink, G. P. M., Kooijman, H., Spek, A. L. & van Koten, G. (2005). *Organometallics*, **24**, 743–749.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Wuts, P. G. W. & Greene, T. W. (2007). *Organic Synthesis*, 4th ed. New Jersey: Wiley.

supporting information

Acta Cryst. (2015). E71, 1125-1131 [doi:10.1107/S2056989015016059]

Crystal structures of 1-bromo-3,5-bis(4,4-dimethyl-1,3-oxazolin-2-yl)benzene 0.15-hydrate and 3,5-bis(4,4-dimethyl-1,3-oxazolin-2-yl)-1-iodobenzene

Timo Stein, Frank Hoffmann and Michael Fröba

Computing details

Data collection: *CrysAlis PRO* (Agilent, 2013) for (1); *APEX2* (Bruker, 2014) for (2). Cell refinement: *CrysAlis PRO* (Agilent, 2013) for (1); *SAINTE* (Bruker, 2014) for (2). Data reduction: *CrysAlis PRO* (Agilent, 2013) for (1); *SAINTE* (Bruker, 2014) for (2). For both compounds, program(s) used to solve structure: *SUPERFLIP* (Palatinus & Chapuis, 2007); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *XP in SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *WinGX* (Farrugia, 2012), *OLEX2* (Dolomanov *et al.*, 2009), *publCIF* (Westrip, 2010) and *PLATON* (Spek, 2009).

(1) 1-Bromo-3,5-bis(4,4-dimethyl-1,3-oxazolin-2-yl)benzene 0.15-hydrate

Crystal data

$C_{16}H_{19}BrN_2O_2 \cdot 0.15H_2O$

$M_r = 353.94$

Monoclinic, $P2_1/n$

$a = 10.0661$ (1) Å

$b = 16.2960$ (2) Å

$c = 11.0400$ (1) Å

$\beta = 114.496$ (2)°

$V = 1647.96$ (4) Å³

$Z = 4$

$F(000) = 726$

$D_x = 1.427$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 11733 reflections

$\theta = 3.2\text{--}31.8^\circ$

$\mu = 2.50$ mm⁻¹

$T = 100$ K

Plate, colorless

$0.34 \times 0.26 \times 0.08$ mm

Data collection

Agilent SuperNova Dual Source
diffractometer with an Atlas detector
Radiation source: SuperNova (Mo) X-ray
Source
Mirror monochromator
Detector resolution: 10.4127 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2013)

$T_{\min} = 0.670$, $T_{\max} = 1.000$

29541 measured reflections

5438 independent reflections

4598 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

$\theta_{\max} = 31.5^\circ$, $\theta_{\min} = 3.2^\circ$

$h = -14 \rightarrow 14$

$k = -23 \rightarrow 23$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.035$

$wR(F^2) = 0.080$

$S = 1.05$

5438 reflections

209 parameters

3 restraints

Primary atom site location: iterative

Secondary atom site location: difference Fourier
map

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0278P)^2 + 1.5569P]$
where $P = (F_o^2 + 2F_c^2)/3$

$$\begin{aligned}(\Delta/\sigma)_{\max} &= 0.001 \\ \Delta\rho_{\max} &= 0.85 \text{ e } \text{\AA}^{-3} \\ \Delta\rho_{\min} &= -0.74 \text{ e } \text{\AA}^{-3}\end{aligned}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. All non-hydrogen atoms were refined with anisotropic displacement parameters. Carbon-bound hydrogen atoms were positioned geometrically and refined riding on their respective carbon atom. Bond lengths were fixed at 0.95 Å (aromatic H), 0.98 Å (methyl H), and 0.99 Å (methylene H). $U_{\text{iso}}(\text{H})$ was fixed at 1.5 $U_{\text{eq}}(\text{O})$ and $U_{\text{eq}}(\text{C})$ for hydroxyl and methyl H atoms or 1.2 $U_{\text{eq}}(\text{C})$ for the remaining H atoms. Methyl H atoms were fitted to the experimental electron density by allowing them to rotate around the C–C bond with a fixed angle (HFIX 137). Occupancy refinement of O3 gave an occupancy of 0.15543 (611) which was subsequently fixed at 0.15. The H atoms H3A and H3B were located with help of *CALC-OH* which is the *WinGX* (Farrugia, 2012) implementation of Nardelli's method (Nardelli, 1999) of OH atom positioning. The coordinates of those H atoms were refined freely while applying restraints to the overall water geometry (O–H = 0.84 Å and H⋯H = 1.328 Å).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.26376 (17)	0.48374 (9)	0.34172 (15)	0.0162 (3)	
C3	0.44468 (17)	0.38604 (9)	0.47611 (15)	0.0161 (3)	
C5	0.45798 (18)	0.44445 (10)	0.28229 (15)	0.0180 (3)	
C4	0.51639 (17)	0.39301 (10)	0.39218 (15)	0.0176 (3)	
H4	0.6035	0.3631	0.4099	0.021*	
C12	0.49957 (17)	0.32940 (10)	0.59022 (16)	0.0184 (3)	
C7	0.13236 (18)	0.53354 (9)	0.31540 (15)	0.0172 (3)	
C6	0.33230 (18)	0.48959 (10)	0.25469 (15)	0.0179 (3)	
H6	0.2934	0.5238	0.1783	0.022*	
C9	−0.0526 (2)	0.62058 (12)	0.23624 (19)	0.0268 (4)	
C14	0.53332 (19)	0.25360 (11)	0.76730 (17)	0.0226 (3)	
C16	0.4386 (3)	0.18050 (16)	0.7561 (4)	0.0682 (10)	
H16A	0.4962	0.1380	0.8186	0.102*	
H16B	0.4009	0.1589	0.6651	0.102*	
H16C	0.3568	0.1968	0.7771	0.102*	
C15	0.5965 (4)	0.28651 (17)	0.9079 (2)	0.0632 (9)	
H15A	0.6590	0.2447	0.9687	0.095*	
H15B	0.5170	0.3004	0.9336	0.095*	
H15C	0.6544	0.3357	0.9127	0.095*	
C2	0.31925 (17)	0.43183 (9)	0.45143 (15)	0.0159 (3)	
H2	0.2716	0.4276	0.5096	0.019*	
C13	0.6535 (3)	0.2366 (2)	0.7218 (3)	0.0757 (12)	
H13A	0.7495	0.2531	0.7914	0.091*	
H13B	0.6568	0.1775	0.7028	0.091*	

C8	-0.0606 (2)	0.57193 (12)	0.35229 (19)	0.0257 (4)	
H8A	-0.0676	0.6094	0.4201	0.031*	
H8B	-0.1460	0.5347	0.3203	0.031*	
C10	-0.0128 (3)	0.70964 (14)	0.2754 (3)	0.0525 (7)	
H10A	-0.0925	0.7364	0.2894	0.079*	
H10B	0.0031	0.7380	0.2043	0.079*	
H10C	0.0766	0.7120	0.3578	0.079*	
C11	-0.1925 (3)	0.6130 (2)	0.1101 (2)	0.0564 (8)	
H11A	-0.2755	0.6304	0.1283	0.085*	
H11C	-0.2061	0.5558	0.0801	0.085*	
H11B	-0.1859	0.6479	0.0405	0.085*	
N2	0.44454 (16)	0.31821 (9)	0.67357 (14)	0.0217 (3)	
N1	0.07115 (17)	0.58157 (10)	0.21786 (15)	0.0248 (3)	
O2	0.61878 (15)	0.28455 (9)	0.60266 (13)	0.0309 (3)	
O1	0.07500 (14)	0.52563 (8)	0.40654 (13)	0.0257 (3)	
Br1	0.55330 (2)	0.45321 (2)	0.16706 (2)	0.02642 (6)	
O3	0.0751 (12)	0.5663 (7)	-0.0328 (10)	0.035 (2)	0.15
H3A	0.076 (18)	0.572 (11)	0.043 (8)	0.050*	0.15
H3B	0.010 (15)	0.532 (9)	-0.071 (14)	0.050*	0.15

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0199 (7)	0.0131 (6)	0.0129 (6)	-0.0033 (5)	0.0042 (5)	-0.0015 (5)
C3	0.0180 (7)	0.0150 (7)	0.0121 (6)	-0.0028 (5)	0.0030 (5)	-0.0007 (5)
C5	0.0211 (7)	0.0192 (7)	0.0136 (6)	-0.0068 (6)	0.0072 (6)	-0.0031 (6)
C4	0.0180 (7)	0.0174 (7)	0.0148 (7)	-0.0036 (5)	0.0042 (6)	-0.0034 (5)
C12	0.0174 (7)	0.0176 (7)	0.0172 (7)	0.0005 (6)	0.0041 (6)	0.0016 (6)
C7	0.0211 (7)	0.0143 (7)	0.0148 (6)	-0.0027 (5)	0.0060 (6)	-0.0007 (5)
C6	0.0227 (7)	0.0157 (7)	0.0123 (6)	-0.0040 (6)	0.0042 (6)	-0.0002 (5)
C9	0.0225 (8)	0.0287 (9)	0.0283 (9)	0.0055 (7)	0.0098 (7)	0.0115 (7)
C14	0.0214 (8)	0.0232 (8)	0.0218 (8)	0.0059 (6)	0.0075 (6)	0.0091 (6)
C16	0.0363 (13)	0.0327 (12)	0.097 (2)	-0.0059 (10)	-0.0107 (13)	0.0361 (14)
C15	0.091 (2)	0.0425 (14)	0.0272 (11)	0.0238 (14)	-0.0041 (13)	0.0066 (10)
C2	0.0195 (7)	0.0135 (6)	0.0131 (6)	-0.0025 (5)	0.0054 (5)	-0.0011 (5)
C13	0.0707 (19)	0.115 (3)	0.0664 (18)	0.071 (2)	0.0532 (16)	0.0711 (19)
C8	0.0246 (8)	0.0265 (8)	0.0263 (9)	0.0063 (7)	0.0108 (7)	0.0070 (7)
C10	0.0556 (15)	0.0216 (10)	0.097 (2)	0.0112 (10)	0.0485 (16)	0.0155 (12)
C11	0.0293 (11)	0.099 (2)	0.0326 (12)	0.0073 (13)	0.0045 (9)	0.0261 (14)
N2	0.0229 (7)	0.0216 (7)	0.0193 (6)	0.0065 (5)	0.0075 (5)	0.0086 (5)
N1	0.0263 (7)	0.0266 (8)	0.0202 (7)	0.0060 (6)	0.0084 (6)	0.0076 (6)
O2	0.0269 (7)	0.0420 (8)	0.0269 (7)	0.0164 (6)	0.0143 (5)	0.0178 (6)
O1	0.0260 (6)	0.0299 (7)	0.0245 (6)	0.0090 (5)	0.0138 (5)	0.0126 (5)
Br1	0.02744 (9)	0.03482 (10)	0.02055 (9)	-0.00288 (7)	0.01352 (7)	0.00215 (7)
O3	0.039 (5)	0.040 (6)	0.024 (4)	-0.002 (4)	0.011 (4)	0.003 (4)

Geometric parameters (Å, °)

C1—C2	1.391 (2)	C14—C13	1.515 (3)
C1—C6	1.399 (2)	C16—H16A	0.9800
C1—C7	1.474 (2)	C16—H16B	0.9800
C3—C2	1.393 (2)	C16—H16C	0.9800
C3—C4	1.396 (2)	C15—H15A	0.9800
C3—C12	1.472 (2)	C15—H15B	0.9800
C5—C6	1.384 (2)	C15—H15C	0.9800
C5—C4	1.389 (2)	C2—H2	0.9500
C5—Br1	1.8898 (16)	C13—O2	1.443 (3)
C4—H4	0.9500	C13—H13A	0.9900
C12—N2	1.269 (2)	C13—H13B	0.9900
C12—O2	1.362 (2)	C8—O1	1.454 (2)
C7—N1	1.266 (2)	C8—H8A	0.9900
C7—O1	1.358 (2)	C8—H8B	0.9900
C6—H6	0.9500	C10—H10A	0.9800
C9—N1	1.485 (2)	C10—H10B	0.9800
C9—C10	1.520 (3)	C10—H10C	0.9800
C9—C11	1.520 (3)	C11—H11A	0.9800
C9—C8	1.537 (3)	C11—H11C	0.9800
C14—N2	1.487 (2)	C11—H11B	0.9800
C14—C16	1.498 (3)	O3—H3A	0.84 (2)
C14—C15	1.511 (3)	O3—H3B	0.84 (2)
C2—C1—C6	120.34 (15)	C14—C15—H15A	109.5
C2—C1—C7	120.67 (14)	C14—C15—H15B	109.5
C6—C1—C7	118.99 (14)	H15A—C15—H15B	109.5
C2—C3—C4	120.20 (14)	C14—C15—H15C	109.5
C2—C3—C12	119.44 (14)	H15A—C15—H15C	109.5
C4—C3—C12	120.35 (14)	H15B—C15—H15C	109.5
C6—C5—C4	122.04 (15)	C1—C2—C3	120.03 (15)
C6—C5—Br1	118.98 (12)	C1—C2—H2	120.0
C4—C5—Br1	118.98 (13)	C3—C2—H2	120.0
C5—C4—C3	118.75 (15)	O2—C13—C14	106.17 (16)
C5—C4—H4	120.6	O2—C13—H13A	110.5
C3—C4—H4	120.6	C14—C13—H13A	110.5
N2—C12—O2	118.75 (15)	O2—C13—H13B	110.5
N2—C12—C3	126.08 (15)	C14—C13—H13B	110.5
O2—C12—C3	115.16 (14)	H13A—C13—H13B	108.7
N1—C7—O1	118.85 (15)	O1—C8—C9	104.25 (14)
N1—C7—C1	126.06 (15)	O1—C8—H8A	110.9
O1—C7—C1	115.09 (13)	C9—C8—H8A	110.9
C5—C6—C1	118.59 (14)	O1—C8—H8B	110.9
C5—C6—H6	120.7	C9—C8—H8B	110.9
C1—C6—H6	120.7	H8A—C8—H8B	108.9
N1—C9—C10	108.11 (16)	C9—C10—H10A	109.5
N1—C9—C11	110.53 (19)	C9—C10—H10B	109.5

C10—C9—C11	111.9 (2)	H10A—C10—H10B	109.5
N1—C9—C8	103.30 (14)	C9—C10—H10C	109.5
C10—C9—C8	110.73 (19)	H10A—C10—H10C	109.5
C11—C9—C8	111.89 (17)	H10B—C10—H10C	109.5
N2—C14—C16	109.10 (15)	C9—C11—H11A	109.5
N2—C14—C15	109.82 (16)	C9—C11—H11C	109.5
C16—C14—C15	110.5 (2)	H11A—C11—H11C	109.5
N2—C14—C13	103.34 (15)	C9—C11—H11B	109.5
C16—C14—C13	113.3 (3)	H11A—C11—H11B	109.5
C15—C14—C13	110.6 (2)	H11C—C11—H11B	109.5
C14—C16—H16A	109.5	C12—N2—C14	106.82 (14)
C14—C16—H16B	109.5	C7—N1—C9	106.76 (15)
H16A—C16—H16B	109.5	C12—O2—C13	104.63 (15)
C14—C16—H16C	109.5	C7—O1—C8	105.11 (13)
H16A—C16—H16C	109.5	H3A—O3—H3B	105 (5)
H16B—C16—H16C	109.5		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C6—H6...O3	0.95	2.55 (1)	3.395 (10)	149 (1)
O3—H3A...N1	0.84 (2)	1.96 (2)	2.796 (10)	178 (15)
C16—H16B...O3 ⁱ	0.98	2.18 (1)	3.045 (11)	146 (1)
C13—H13B...O3 ⁱ	0.99	2.94 (1)	3.656 (12)	130 (1)
C8—H8B...Br1 ⁱⁱ	0.99	3.09 (1)	4.054 (2)	165 (1)
C11—H11C...O3 ⁱⁱⁱ	0.98	2.56 (1)	3.391 (12)	143 (1)
O3—H3B...N1 ⁱⁱⁱ	0.84 (2)	2.37 (9)	3.107 (11)	148 (15)
O3—H3B...O3 ⁱⁱⁱ	0.84 (2)	2.32 (13)	2.90 (2)	126 (13)

Symmetry codes: (i) $-x+1/2, y-1/2, -z+1/2$; (ii) $x-1, y, z$; (iii) $-x, -y+1, -z$.

(2) 3,5-Bis(4,4-dimethyl-1,3-oxazolin-2-yl)-1-iodobenzene

Crystal data

C₁₆H₁₉IN₂O₂
M_r = 398.23
 Monoclinic, *P*2₁/*n*
a = 9.6195 (2) Å
b = 9.9759 (2) Å
c = 17.2951 (4) Å
 β = 94.648 (1)°
V = 1654.23 (6) Å³
Z = 4

F(000) = 792
D_x = 1.599 Mg m⁻³
 Mo *K*α radiation, λ = 0.71073 Å
 Cell parameters from 9629 reflections
 θ = 2.3–34.2°
 μ = 1.94 mm⁻¹
T = 100 K
 Block, colorless
 0.22 × 0.12 × 0.08 mm

Data collection

Bruker SMART APEX CCD area-detector
 diffractometer
 Radiation source: Incoatec microfocus
 Mirror monochromator
 ω scans

Absorption correction: numerical
 (*SADABS*; Bruker, 2014)
T_{min} = 0.649, *T_{max}* = 0.747
 45346 measured reflections
 6463 independent reflections
 6035 reflections with *I* > 2σ(*I*)

$R_{\text{int}} = 0.019$
 $\theta_{\text{max}} = 33.5^\circ$, $\theta_{\text{min}} = 2.4^\circ$
 $h = -14 \rightarrow 14$

$k = -15 \rightarrow 15$
 $l = -26 \rightarrow 26$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.017$
 $wR(F^2) = 0.043$
 $S = 1.07$
 6463 reflections
 194 parameters
 0 restraints
 Primary atom site location: iterative

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0189P)^2 + 0.8336P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.007$
 $\Delta\rho_{\text{max}} = 0.82 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.62 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. All non-hydrogen atoms were refined with anisotropic displacement parameters. Carbon-bound hydrogen atoms were positioned geometrically and refined riding on their respective carbon atom. Bond lengths were fixed at 0.95 Å (aromatic H), 0.98 Å (methyl H), and 0.99 Å (methylene H). $U_{\text{iso}}(\text{H})$ was fixed at 1.5 $U_{\text{eq}}(\text{O})$ and $U_{\text{eq}}(\text{C})$ for hydroxyl and methyl H atoms or 1.2 $U_{\text{eq}}(\text{C})$ for the remaining H atoms. Methyl H atoms were fitted to the experimental electron density by allowing them to rotate around the C–C bond with a fixed angle (HFIX 137).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.56410 (10)	0.66623 (9)	0.41768 (5)	0.01116 (15)
C2	0.48728 (10)	0.55181 (9)	0.39654 (5)	0.01198 (15)
H2	0.5232	0.4878	0.3627	0.014*
C3	0.35754 (10)	0.53146 (9)	0.42507 (5)	0.01135 (15)
C4	0.30480 (10)	0.62363 (9)	0.47619 (6)	0.01244 (16)
H4	0.2179	0.6083	0.4971	0.015*
C5	0.38251 (10)	0.73850 (9)	0.49576 (6)	0.01210 (15)
C6	0.51110 (10)	0.76077 (9)	0.46704 (6)	0.01225 (15)
H6	0.5626	0.8397	0.4808	0.015*
C7	0.70012 (10)	0.69002 (9)	0.38670 (5)	0.01184 (15)
C8	0.89158 (12)	0.62676 (11)	0.33104 (8)	0.0221 (2)
H8A	0.8999	0.6078	0.2754	0.027*
H8B	0.9694	0.5822	0.3621	0.027*
C9	0.89278 (10)	0.77946 (10)	0.34634 (6)	0.01420 (16)
C10	0.88285 (14)	0.85997 (15)	0.27141 (7)	0.0273 (2)
H10A	0.8013	0.8304	0.2381	0.041*
H10B	0.9674	0.8460	0.2444	0.041*

H10C	0.8735	0.9554	0.2834	0.041*
C11	1.01869 (12)	0.82249 (13)	0.39920 (7)	0.0236 (2)
H11A	1.0107	0.9179	0.4116	0.035*
H11B	1.1038	0.8074	0.3729	0.035*
H11C	1.0227	0.7699	0.4472	0.035*
C12	0.27785 (10)	0.41157 (10)	0.39882 (6)	0.01243 (16)
C13	0.12395 (12)	0.24676 (11)	0.40919 (6)	0.0201 (2)
H13A	0.0211	0.2409	0.4004	0.024*
H13B	0.1575	0.1747	0.4454	0.024*
C14	0.19275 (12)	0.23620 (11)	0.33195 (6)	0.01761 (18)
C15	0.09128 (15)	0.27569 (14)	0.26324 (7)	0.0281 (3)
H15A	0.0521	0.3643	0.2727	0.042*
H15B	0.0158	0.2097	0.2570	0.042*
H15C	0.1408	0.2784	0.2159	0.042*
C16	0.25716 (16)	0.09954 (12)	0.31923 (9)	0.0299 (3)
H16A	0.3090	0.1023	0.2728	0.045*
H16B	0.1832	0.0320	0.3124	0.045*
H16C	0.3208	0.0762	0.3643	0.045*
N1	0.76407 (9)	0.80152 (9)	0.38642 (5)	0.01435 (15)
N2	0.30375 (10)	0.33974 (9)	0.34100 (5)	0.01682 (16)
O1	0.75825 (9)	0.58108 (8)	0.35446 (5)	0.01984 (15)
O2	0.16674 (9)	0.37812 (8)	0.43947 (5)	0.01763 (15)
II	0.30572 (2)	0.88951 (2)	0.56511 (2)	0.01650 (2)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0122 (4)	0.0101 (4)	0.0111 (4)	-0.0011 (3)	0.0008 (3)	0.0002 (3)
C2	0.0144 (4)	0.0106 (4)	0.0109 (4)	-0.0012 (3)	0.0007 (3)	-0.0008 (3)
C3	0.0130 (4)	0.0099 (4)	0.0108 (4)	-0.0018 (3)	-0.0007 (3)	0.0001 (3)
C4	0.0125 (4)	0.0113 (4)	0.0134 (4)	-0.0008 (3)	0.0006 (3)	0.0002 (3)
C5	0.0132 (4)	0.0102 (4)	0.0130 (4)	0.0003 (3)	0.0016 (3)	-0.0017 (3)
C6	0.0134 (4)	0.0101 (4)	0.0132 (4)	-0.0011 (3)	0.0008 (3)	-0.0012 (3)
C7	0.0133 (4)	0.0111 (4)	0.0113 (4)	0.0000 (3)	0.0017 (3)	-0.0010 (3)
C8	0.0184 (5)	0.0177 (5)	0.0320 (6)	-0.0007 (4)	0.0120 (4)	-0.0047 (4)
C9	0.0128 (4)	0.0151 (4)	0.0152 (4)	0.0001 (3)	0.0040 (3)	0.0016 (3)
C10	0.0277 (6)	0.0344 (6)	0.0208 (5)	0.0057 (5)	0.0073 (4)	0.0111 (5)
C11	0.0150 (5)	0.0285 (6)	0.0271 (5)	-0.0008 (4)	0.0000 (4)	-0.0028 (4)
C12	0.0129 (4)	0.0119 (4)	0.0124 (4)	-0.0030 (3)	0.0004 (3)	0.0004 (3)
C13	0.0224 (5)	0.0194 (5)	0.0191 (5)	-0.0115 (4)	0.0043 (4)	-0.0044 (4)
C14	0.0203 (5)	0.0153 (4)	0.0174 (4)	-0.0078 (4)	0.0028 (4)	-0.0047 (3)
C15	0.0329 (6)	0.0294 (6)	0.0207 (5)	-0.0104 (5)	-0.0061 (5)	-0.0043 (4)
C16	0.0319 (7)	0.0168 (5)	0.0419 (7)	-0.0062 (4)	0.0090 (6)	-0.0096 (5)
N1	0.0127 (3)	0.0128 (3)	0.0183 (4)	-0.0013 (3)	0.0051 (3)	-0.0004 (3)
N2	0.0191 (4)	0.0150 (4)	0.0168 (4)	-0.0070 (3)	0.0038 (3)	-0.0042 (3)
O1	0.0188 (4)	0.0137 (3)	0.0284 (4)	-0.0021 (3)	0.0107 (3)	-0.0070 (3)
O2	0.0175 (3)	0.0182 (4)	0.0178 (3)	-0.0083 (3)	0.0057 (3)	-0.0050 (3)
II	0.01488 (3)	0.01529 (3)	0.01976 (4)	-0.00079 (2)	0.00401 (2)	-0.00607 (2)

Geometric parameters (Å, °)

C1—C2	1.3924 (13)	C10—H10A	0.9800
C1—C6	1.3959 (13)	C10—H10B	0.9800
C1—C7	1.4728 (13)	C10—H10C	0.9800
C2—C3	1.3936 (13)	C11—H11A	0.9800
C2—H2	0.9500	C11—H11B	0.9800
C3—C4	1.3990 (13)	C11—H11C	0.9800
C3—C12	1.4724 (13)	C12—N2	1.2711 (13)
C4—C5	1.3948 (13)	C12—O2	1.3675 (12)
C4—H4	0.9500	C13—O2	1.4581 (13)
C5—C6	1.3878 (13)	C13—C14	1.5418 (15)
C5—I1	2.0968 (9)	C13—H13A	0.9900
C6—H6	0.9500	C13—H13B	0.9900
C7—N1	1.2713 (12)	C14—N2	1.4847 (13)
C7—O1	1.3620 (12)	C14—C16	1.5209 (17)
C8—O1	1.4494 (14)	C14—C15	1.5271 (17)
C8—C9	1.5460 (15)	C15—H15A	0.9800
C8—H8A	0.9900	C15—H15B	0.9800
C8—H8B	0.9900	C15—H15C	0.9800
C9—N1	1.4835 (13)	C16—H16A	0.9800
C9—C11	1.5191 (15)	C16—H16B	0.9800
C9—C10	1.5209 (15)	C16—H16C	0.9800
C2—C1—C6	120.14 (9)	H10B—C10—H10C	109.5
C2—C1—C7	120.32 (8)	C9—C11—H11A	109.5
C6—C1—C7	119.53 (8)	C9—C11—H11B	109.5
C1—C2—C3	119.86 (9)	H11A—C11—H11B	109.5
C1—C2—H2	120.1	C9—C11—H11C	109.5
C3—C2—H2	120.1	H11A—C11—H11C	109.5
C2—C3—C4	120.60 (9)	H11B—C11—H11C	109.5
C2—C3—C12	117.90 (8)	N2—C12—O2	118.56 (9)
C4—C3—C12	121.49 (9)	N2—C12—C3	124.76 (9)
C5—C4—C3	118.60 (9)	O2—C12—C3	116.67 (8)
C5—C4—H4	120.7	O2—C13—C14	104.13 (8)
C3—C4—H4	120.7	O2—C13—H13A	110.9
C6—C5—C4	121.35 (9)	C14—C13—H13A	110.9
C6—C5—I1	117.08 (7)	O2—C13—H13B	110.9
C4—C5—I1	121.49 (7)	C14—C13—H13B	110.9
C5—C6—C1	119.41 (9)	H13A—C13—H13B	108.9
C5—C6—H6	120.3	N2—C14—C16	109.93 (10)
C1—C6—H6	120.3	N2—C14—C15	108.15 (9)
N1—C7—O1	118.80 (9)	C16—C14—C15	111.15 (10)
N1—C7—C1	125.93 (9)	N2—C14—C13	102.51 (8)
O1—C7—C1	115.26 (8)	C16—C14—C13	113.27 (10)
O1—C8—C9	104.86 (8)	C15—C14—C13	111.38 (10)
O1—C8—H8A	110.8	C14—C15—H15A	109.5
C9—C8—H8A	110.8	C14—C15—H15B	109.5

O1—C8—H8B	110.8	H15A—C15—H15B	109.5
C9—C8—H8B	110.8	C14—C15—H15C	109.5
H8A—C8—H8B	108.9	H15A—C15—H15C	109.5
N1—C9—C11	109.38 (9)	H15B—C15—H15C	109.5
N1—C9—C10	108.82 (9)	C14—C16—H16A	109.5
C11—C9—C10	110.82 (10)	C14—C16—H16B	109.5
N1—C9—C8	103.37 (8)	H16A—C16—H16B	109.5
C11—C9—C8	112.08 (9)	C14—C16—H16C	109.5
C10—C9—C8	112.05 (10)	H16A—C16—H16C	109.5
C9—C10—H10A	109.5	H16B—C16—H16C	109.5
C9—C10—H10B	109.5	C7—N1—C9	107.12 (8)
H10A—C10—H10B	109.5	C12—N2—C14	106.86 (9)
C9—C10—H10C	109.5	C7—O1—C8	105.40 (8)
H10A—C10—H10C	109.5	C12—O2—C13	104.14 (8)
C6—C1—C2—C3	-0.31 (14)	C2—C3—C12—O2	166.31 (9)
C7—C1—C2—C3	-178.83 (9)	C4—C3—C12—O2	-14.56 (14)
C1—C2—C3—C4	-1.37 (14)	O2—C13—C14—N2	18.91 (11)
C1—C2—C3—C12	177.77 (9)	O2—C13—C14—C16	137.30 (10)
C2—C3—C4—C5	2.25 (14)	O2—C13—C14—C15	-96.53 (10)
C12—C3—C4—C5	-176.86 (9)	O1—C7—N1—C9	1.19 (13)
C3—C4—C5—C6	-1.49 (14)	C1—C7—N1—C9	-177.34 (9)
C3—C4—C5—I1	175.31 (7)	C11—C9—N1—C7	-124.42 (10)
C4—C5—C6—C1	-0.15 (14)	C10—C9—N1—C7	114.38 (10)
I1—C5—C6—C1	-177.09 (7)	C8—C9—N1—C7	-4.88 (11)
C2—C1—C6—C5	1.07 (14)	O2—C12—N2—C14	2.75 (13)
C7—C1—C6—C5	179.60 (9)	C3—C12—N2—C14	-175.98 (9)
C2—C1—C7—N1	162.51 (10)	C16—C14—N2—C12	-134.30 (11)
C6—C1—C7—N1	-16.02 (15)	C15—C14—N2—C12	104.17 (11)
C2—C1—C7—O1	-16.07 (13)	C13—C14—N2—C12	-13.58 (12)
C6—C1—C7—O1	165.40 (9)	N1—C7—O1—C8	3.37 (13)
O1—C8—C9—N1	6.63 (11)	C1—C7—O1—C8	-177.95 (9)
O1—C8—C9—C11	124.30 (10)	C9—C8—O1—C7	-6.05 (12)
O1—C8—C9—C10	-110.37 (11)	N2—C12—O2—C13	10.14 (13)
C2—C3—C12—N2	-14.94 (15)	C3—C12—O2—C13	-171.02 (9)
C4—C3—C12—N2	164.19 (10)	C14—C13—O2—C12	-17.51 (11)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C6—H6...I1 ⁱ	0.95	3.11	3.9679 (9)	150
C10—H10B...N2 ⁱⁱ	0.98	2.75	3.7228 (15)	172

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $-x+3/2, y+1/2, -z+1/2$.