



# Crystal structures of three new *N*-halomethylated quaternary ammonium salts

Carolina Múnera-Orozco,<sup>a</sup> Rogelio Ocampo-Cardona,<sup>a</sup> David L. Cedeño,<sup>b,‡</sup>  
Rubén A. Toscano<sup>c</sup> and Luz Amalia Ríos-Vásquez<sup>a\*</sup>

Received 22 July 2015  
Accepted 14 September 2015

Edited by H. Stoeckli-Evans, University of  
Neuchâtel, Switzerland

‡ Current address: Millennium Pain Center,  
Basic Science, 1015 S Mercer Ave Bloomington,  
IL 61701 USA.

**Keywords:** crystal structure; *N*-halomethylated  
quaternary ammonium salts; cation– $\pi$  inter-  
action; halogen bond; hydrogen bonding

**CCDC references:** 1424320; 1424319;  
1424318


**Supporting information:** this article has  
supporting information at journals.iucr.org/e

<sup>a</sup>Departamento de Química, Universidad de Caldas, Manizales, Caldas, Colombia, <sup>b</sup>Department of Chemistry, Illinois State University, Normal, Illinois, USA, and <sup>c</sup>Instituto de Química, UNAM, Circuito Exterior, Ciudad Universitaria, Delegación Coyoacán, C.P. 04510, México, D.F., Mexico. \*Correspondence e-mail: amalia.rios@ucaldas.edu.co


In the crystals of the title *N*-halomethylated quaternary ammonium salts,  $C_{19}H_{23}IN^+I^-$ , (I) [systematic name: *N*-(4,4-diphenylbut-3-en-1-yl)-*N*-iodomethyl-*N,N*-dimethylammonium iodide],  $C_{20}H_{25}IN^+I^-$ , (II) [systematic name: *N*-(5,5-diphenylpent-4-en-1-yl)-*N*-iodomethyl-*N,N*-dimethylammonium iodide], and  $C_{21}H_{27}IN^+I^-$ , (III) [systematic name: *N*-(6,6-diphenylhex-5-en-1-yl)-*N*-iodomethyl-*N,N*-dimethylammonium iodide], there are short  $I \cdots I^-$  interactions of 3.564 (4), 3.506 (1) and 3.557 (1) Å for compounds (I), (II) and (III), respectively. Compound (I) crystallizes in the Sohncke group  $P2_1$  as an ‘enantiopure’ compound and is therefore a potential material for NLO properties. In the crystal of compound (I), molecules are linked by  $C-H \cdots I^-$  and  $C-H \cdots \pi$  interactions which, together with the  $I \cdots I^-$  interactions, lead to the formation of ribbons along [100]. In (II), there are only  $C-H \cdots I^-$  interactions which, together with the  $I \cdots I^-$  interactions, lead to the formation of helices along [010]. In (III), apart from the  $I \cdots I^-$  interactions, there are no significant intermolecular interactions present.

## 1. Chemical context and background to halogen bonding and cation– $\pi$ interactions


Quaternary ammonium salts have been widely studied as anti-cancer (Wang *et al.*, 2012; Song *et al.*, 2013), anti-fungal (Ng *et al.*, 2006), anti-HIV-1 (Shiraishi *et al.*, 2000), anti-bacterial (Calvani *et al.*, 1998), anti-malarial (Calas *et al.*, 1997; Calas *et al.*, 2000) and anti-leishmanial (Mavromoustakos *et al.*, 2001) pharmaceuticals. Our research group has been working in the past few years on the activity of quaternary *N*-halomethyl ammonium salts for likely pharmaceutical purposes, specifically against axenic *L. (V) panamensis* and *L. (L) amazonensis* parasites, human pathogenic species that cause cutaneous and mucocutaneous leishmaniasis. The experiments proved that these compounds are very promising anti-leishmanial molecules, and very significant changes in their activity were observed upon a slight modification of the carbon skeleton by only a single methylene unit (Ríos-Vásquez *et al.*, 2015). A preliminary effort at understanding a structure–activity relationship with three *N*-iodomethyl quaternary ammonium salts (I), (II) and (III) of the form  $[ICH_2N(CH_3)_3(CH_2)_nCH=C(Ph)_2]^+I^-$  (with  $n = 2, 3$  and 4, respectively) is currently being carried out. One possible approach to understand the different activities is to establish what kind of interactions are present in compounds (I)–(III), for example whether  $C-I \cdots I^-$  (Desiraju *et al.*, 2013),  $C-H \cdots I^-$  (Glidewell *et al.*, 1994),  $C-H \cdots \pi$  (Nishio *et al.*, 1998) or cation– $\pi$  (Dougherty, 1996), and if so, how these interactions may affect their structure and biological properties.



As defined by International Union for Pure and Applied Chemistry (IUPAC): a halogen bond occurs when there is evidence of a net attractive interaction between an electrophilic region associated with a halogen atom in a molecular entity and a nucleophilic region in another, or the same, molecular entity (Desiraju *et al.*, 2013). Halogen bonds are characterized by  $X \cdots X$  distances that are clearly shorter than the van der Waals radii sum (Formigué, 2009; Awadi *et al.*; 2006); otherwise this interaction is neglected. In a similar way, the existence of  $C-H \cdots X$  hydrogen bonds ( $X = F, Cl, Br$  or  $I$ ) in neutral organic molecules (Aakerøy & Seddon, 1993) and even in organic salts has been recognized. On the other hand, a special kind of hydrogen bond, defined as a weak interaction between a soft acid (*i.e.* an  $sp^3$ ,  $sp^2$  or  $sp$  C–H system) and a soft base (*i.e.* an aromatic, olefinic or acetylenic  $p$  system), with a significant role on diverse chemical and biological phenomena has recently been described (Nishio, 2012). In particular, this interaction exerts an observable influence on host–guest recognition and crystal packing in the solid state. A related attraction is the cation– $\pi$  interaction, which is regarded as an electrostatic attraction between a positive charge and the quadrupole moment of an aromatic ring (Dougherty, 1996). A cation– $\pi$  interaction between aromatic and ammonium ions is known to play an important role in many biological systems (Ma & Dougherty, 1997; Dougherty, 2013; Sussman *et al.*, 1991; Chen *et al.*, 2011). Part of our research interest is focused not only in understanding the reactive nature of alpha ammonium distonic radical cations which are generated from *N*-halomethylated quaternary ammonium salts (Ríos *et al.*, 1996; Ríos, Bartberger *et al.*, 1997), but also in trying to understand how these salts behave against *Leishmania* parasites (Ríos-Vásquez *et al.*, 2015). The recognition of the occurrence of some supramolecular interactions in these salts may lead to a better understanding of the likely novel biological binding sites, and therefore to new suggestions about biocatalytic mechanisms.



The title *N*-iodomethyl quaternary ammonium salts, (I)–(III), were synthesized following standard procedures used for other related compounds (Newcomb *et al.*, 1993; Horner *et al.*, 1995) and suitable crystals were obtained (Múnera-Orozco, 2014). This paper reports a comparative crystal structure and supramolecular interactions analysis for the aforementioned compounds.




**Figure 1**  
The molecular structure of compound (I), showing the atom labelling. Displacement ellipsoids are drawn at the 50% probability level.


## 2. Structural commentary

Compound (I), Fig. 1, crystallizes in the non-centrosymmetric monoclinic space group  $P2_1$  and is therefore, a potential material for NLO properties. The asymmetric unit consists of an ammonium cation and an iodide anion. In the geminal-substituted diphenylethene unit, the phenyl rings (C5–C10 and C11–C16) are inclined to one another by  $74.6(2)^\circ$ , and are twisted from the mean plane of the central C=C bond fragment (C2–C5/C11) by  $33.2(2)$  and  $61.4(2)^\circ$ , respectively. Coplanarity of the olefin skeleton and the peripheral phenyl rings is prevented because of steric congestion between the associated phenyl rings. The conformation of the side chain reveals an *all-trans* extended conformation with the iodomethyl moiety on one side of the backbone chain, with bond lengths and angles in the expected ranges.

In compound (II), Fig. 2, the dihedral angles between the mean planes of the C=C double-bond fragment (C3–C6/C12) and the two phenyl rings (C6–C11 and C12–C17) are  $31.1(4)$



**Figure 2**  
The molecular structure of compound (II), showing the atom labelling. Displacement ellipsoids are drawn at the 50% probability level.




**Figure 3**  
The molecular structure of compound (III), showing the atom labelling. Displacement ellipsoids are drawn at the 50% probability level.

and  $58.6(4)^\circ$ , respectively, while the phenyl rings are inclined to one another by  $76.2(4)^\circ$ . The *N*-iodomethyl-*N,N*-dimethyl-*N*-propylammonium moiety adopts a fully extended conformation with one methyl group and the iodomethyl unit on opposite sides of the backbone of the side chain (Fig. 2). This conformation seems to be partially supported by a C–H···I<sup>−</sup> hydrogen bond (Table 2 and *Supramolecular features*).

In compound (III), Fig. 3, the phenyl rings are twisted out of the plane defined by the ethylene moiety (C4–C7/C13), making dihedral angles of  $38.7(4)$  and  $78.7(6)^\circ$  for the *trans* (C7–C12) and *cis* (C13–C18) phenyl rings, respectively. The phenyl rings are inclined to one another by  $78.5(6)^\circ$ . The alkylamino side chain is almost fully extended away from the geminal-substituted ethene group.

### 3. Supramolecular features

In the crystal of (I), ribbons are formed, by I1···I2<sup>i</sup> contacts [ $3.564(4)$  Å; symmetry code: (i)  $-x - 1, y - \frac{1}{2}, -z + 1$ ] and



**Figure 4**  
The crystal packing of compound (I), viewed along the *b* axis, showing the intermolecular contacts (dashed lines; see Table 1).

**Table 1**  
Hydrogen-bond geometry (Å, °) for (I).

C<sub>g</sub> is the centroid of the C11–C16 ring.

| <i>D</i> –H··· <i>A</i>              | <i>D</i> –H | H··· <i>A</i> | <i>D</i> ··· <i>A</i> | <i>D</i> –H··· <i>A</i> |
|--------------------------------------|-------------|---------------|-----------------------|-------------------------|
| C17–H17B···I2 <sup>i</sup>           | 0.97        | 3.00          | 3.919 (5)             | 159                     |
| C7–H7···C <sub>g</sub> <sup>ii</sup> | 0.93        | 2.84          | 3.030 (5)             | 143                     |


Symmetry codes: (i)  $-x, y - \frac{1}{2}, -z + 1$ ; (ii)  $x + 1, y, z$ .

**Table 2**  
Hydrogen-bond geometry (Å, °) for (II).


| <i>D</i> –H··· <i>A</i> | <i>D</i> –H | H··· <i>A</i> | <i>D</i> ··· <i>A</i> | <i>D</i> –H··· <i>A</i> |
|-------------------------|-------------|---------------|-----------------------|-------------------------|
| C2–H2B···I2             | 0.97        | 3.06          | 4.001 (7)             | 165                     |

C–H···I<sup>−</sup> hydrogen bonds, along the *a*-axis direction. The chains are reinforced by C–H···π interactions (Fig. 4 and Table 1).

In the crystal of (II), helical chains along the *b*-axis direction are formed by molecules linked *via* C–H···I<sup>−</sup> (Table 2) and I1···I2<sup>ii</sup> interactions [ $3.506(1)$  Å; symmetry code: (ii)  $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$ ]; as shown in Fig. 5. Here no C–H···π interactions are present in the crystal packing. The closest distance between the ammonium substituents and any of the phenyl rings is *ca* 7.18 Å. These features clearly rule out an intramolecular cation–π interaction for this molecule in the solid state. However, in studies of distonic radical cation (Ríos *et al.* 1996; Yates *et al.*, 1986), evidence is presented that the active conformation of the alkylamino side chain is oriented



**Figure 5**  
The crystal packing of compound (II), viewed along the *b* axis, showing the intermolecular contacts (dashed lines; see Table 2).




**Figure 6**  
The crystal packing of compound (III), viewed along the *b* axis.

toward and above the plane of the C=C double bond of the geminal-substituted ethene group. These results confirm that there is considerable freedom of rotation about the bonds separating the basic amino function and the tricyclic system, and thus numerous interconvertible side-chain conformations, differing only slightly in potential energy, may exist.

In the crystal of (III), apart from the  $I1 \cdots I2^{iii}$  contact of 3.557 (1) Å [symmetry code: (iii)  $-x, -y + 1, -z$ ], there are no other significant intermolecular contacts present (Fig. 6). The only possible conclusion regarding the crystal structure of (III) is that the steric requirements in this molecule outweigh the additional stabilization obtained by an intramolecular cation- $\pi$  interaction.

#### 4. Synthesis and crystallization

The general procedure for the preparation of the title quaternary ammonium salts is illustrated in Fig. 7 for



**Figure 7**  
The general procedure for the preparation of the title quaternary ammonium salts.

compounds (I)–(III). The reactions were carried out following a standard literature method (Ríos *et al.*, 1996) starting from the appropriate amine [*N,N*-dimethyl-4,4-diphenylbut-3-en-1-amine 1(a), *N,N*-dimethyl-5,5-diphenylpent-4-en-1-amine 1(b) and *N,N*-dimethyl-6,6-diphenylhex-5-en-1-amine 1(c)]. Typically,  $\text{CH}_2\text{I}_2$  (4 eq) and 1 eq of the starting tertiary amine [for example, compound 1(a) for the synthesis of (I); as shown in Fig. 7] were dissolved in acetonitrile. The reactions were allowed to run overnight at room temperature, and the precipitated salts were filtered off and washed several times with diethyl ether, and then recrystallized from a binary mixture water–isopropanol. The desired products were obtained as colourless crystals.

**Compound (I):** The product was obtained as a white solid in 74% yield; m.p. 425–427 K.  $^1\text{H}$  NMR (DMSO, 300 MHz,  $\delta$ , p.p.m.): 2.49 (*m*, 2H), 3.12 (*s*, 6H), 3.50 (*m*, 2H), 5.05 (*s*, 2H), 6.07 (*t*,  $J = 7.4$  Hz, 1H), 7.15–7.58 (*m*, 10H) p.p.m.  $^{13}\text{C}$  NMR (DMSO, 75 MHz, p.p.m.) 23.70, 31.49, 51.66, 63.58, 121.92, 127.19–129.51, 138.79, 141.62, 145.03 p.p.m. Elemental analysis calculated for  $\text{C}_{19}\text{H}_{23}\text{NI}_2$ : C, 43.95%; H, 4.46%; N, 2.70%; found, C, 43.48%; H, 4.35%; N, 2.68%. MS–ESI calculated for  $\text{C}_{19}\text{H}_{23}\text{NI}$ : 392.09, found: 391.95.

**Compound (II):** The product was obtained as a white solid in 77% yield; m.p. 430–437 K.  $^1\text{H}$  NMR (DMSO, 300 MHz,  $\delta$ , p.p.m.): 1.85 (*m*, 2H), 2.12 (*m*, 2H), 2.51 (*m*, 2H), 3.15 (*s*, 6H), 5.18 (*s*, 2H), 6.14 (*t*,  $J = 7.2$  Hz, 1H), 7.11–7.51 (10H).  $^{13}\text{C}$  NMR (DMSO, 75 MHz, p.p.m.): 22.30, 25.91, 39.01, 51.19, 63.84, 126.84–141.68. ESI–MS  $m/z$  calculated for  $\text{C}_{20}\text{H}_{25}\text{NI}$ : 406.10, found: 406.20.

**Compound (III):** The product was obtained as a white solid in 72% yield; m.p. 429–431 K.  $^1\text{H}$  NMR (DMSO, 300 MHz,  $\delta$ , p.p.m.): 1.45 (*m*, 2H), 1.68 (*m*, 2H), 2.12 (*m*, 2H), 2.51 (*m*, 2H), 3.10 (*s*, 6H), 5.14 (*s*, 2H), 6.14 (*t*,  $J = 7.3$  Hz, 1H), 7.06–7.51 (*m*, 10H) p.p.m.  $^{13}\text{C}$  NMR (DMSO, 75 MHz, p.p.m.): 25.07, 28.91, 31.91, 35.35, 54.29, 67.34, 129.89–132.50, 130.19, 142.45, 144.34, 145.02. Elemental analysis calculated for  $\text{C}_{21}\text{H}_{27}\text{NI}_2$ : C, 46.09%; H, 4.97%; N, 2.56%; found C, 45.91%; H, 4.93%; N, 2.58%. ESI–MS  $m/z$  calculated for  $\text{C}_{21}\text{H}_{27}\text{NI}$ : 420.12, found: 420.20.

#### 5. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. For all three compounds the C-bound H atoms were included in calculated positions and treated as riding atoms: C–H = 0.93–0.99 Å with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$  for methyl H atoms and  $1.2U_{\text{eq}}(\text{C})$  for other H atoms. Refining the structure of compound (I) in the non-centrosymmetric space group gives a value of 0.02 (3) for the Flack parameter (Flack & Bernardinelli, 1999), confirming that the direction of the polar axis has been correctly determined. The studied crystal of compound (III) was a nonmerohedral twin with a ratio of two major domains of 0.374 (2):0.626 (2). The two domains are rotated from each other by  $180.0^\circ$  about the reciprocal axis  $a^*$ , as determined by the *CELL NOW* program (Sheldrick, 2004). The final refinement was carried out using the twinned data set.



**Table 3**  
Experimental details.

|   | (I)   | (II)  | (III)   |
|---|---|---|---|
| Crystal data  |   |   |   |
| Chemical formula  | C <sub>19</sub> H <sub>23</sub> IN <sup>+</sup> ·I <sup>-</sup> | C <sub>20</sub> H <sub>25</sub> IN <sup>+</sup> ·I <sup>-</sup> | C <sub>21</sub> H <sub>27</sub> IN <sup>+</sup> ·I <sup>-</sup> |
| <i>M<sub>r</sub></i>  | 519.18  | 533.21  | 547.23  |
| Crystal system, space group   | Monoclinic, <i>P</i> 2 <sub>1</sub>                             | Monoclinic, <i>C</i> 2/ <i>c</i>                                | Monoclinic, <i>P</i> 2 <sub>1</sub> / <i>c</i>                  |
| Temperature (K)   | 298   | 298   | 298   |
| <i>a</i> , <i>b</i> , <i>c</i> (Å)  | 7.9254 (2), 13.6161 (3), 9.4632 (2)                             | 37.778 (7), 6.6323 (12), 17.021 (3)                             | 8.9423 (12), 24.058 (3), 10.3749 (13)                           |
| $\beta$ (°)   | 103.320 (1)   | 100.567 (4)   | 103.656 (3)   |
| <i>V</i> (Å <sup>3</sup> )  | 993.73 (4)  | 4192.3 (13)   | 2168.9 (5)  |
| <i>Z</i>  | 2   | 8   | 4   |
| Radiation type  | Mo <i>K</i> $\alpha$  | Mo <i>K</i> $\alpha$  | Mo <i>K</i> $\alpha$  |
| $\mu$ (mm <sup>-1</sup> )   | 3.16  | 3.00  | 2.90  |
| Crystal size (mm)   | 0.23 × 0.19 × 0.12  | 0.21 × 0.20 × 0.08  | 0.32 × 0.22 × 0.04  |
| Data collection   |   |   |   |
| Diffractometer  | Bruker SMART APEX CCD   | Bruker SMART APEX CCD   | Bruker SMART APEX CCD   |
| Absorption correction   | Multi-scan ( <i>SADABS</i> ; Bruker, 2012)                      | Multi-scan ( <i>SADABS</i> ; Bruker, 2012)                      | Multi-scan ( <i>TWINABS</i> ; Bruker, 2012)                     |
| <i>T<sub>min</sub></i> , <i>T<sub>max</sub></i>   | 0.624, 0.745  | 0.349, 0.745  | 0.273, 0.429  |
| No. of measured, independent and observed [ <i>I</i> > 2 $\sigma$ ( <i>I</i> )] reflections                             | 5791, 3085, 3013  | 16925, 3808, 3114   | 3961, 3961, 2941  |
| <i>R<sub>int</sub></i>  | 0.016   | 0.079   | 0.079   |
| ( <i>sin</i> $\theta$ / $\lambda$ ) <sub>max</sub> (Å <sup>-1</sup> )   | 0.602   | 0.602   | 0.603   |
| Refinement  |   |   |   |
| <i>R</i> [ <i>F</i> <sup>2</sup> > 2 $\sigma$ ( <i>F</i> <sup>2</sup> )], <i>wR</i> [ <i>F</i> <sup>2</sup> ], <i>S</i> | 0.017, 0.038, 1.08  | 0.052, 0.145, 1.05  | 0.060, 0.138, 1.05  |
| No. of reflections  | 3085  | 3808  | 3961  |
| No. of parameters   | 202   | 210   | 220   |
| No. of restraints   | 1   | 0   | 0   |
| H-atom treatment  | H-atom parameters constrained                                   | H-atom parameters constrained                                   | H-atom parameters constrained                                   |
| $\Delta\rho_{\max}$ , $\Delta\rho_{\min}$ (e Å <sup>-3</sup> )  | 0.27, -0.46   | 1.90, -1.98   | 0.82, -0.80   |
| Absolute structure  | Refined as an inversion twin                                    | —   | —   |
| Absolute structure parameter  | 0.02 (3)  | —   | —   |

Computer programs: *APEX2* and *SAINT* (Bruker, 2012), *SHELXS97* (Sheldrick, 2008), *SHELXL2014* (Sheldrick, 2015), *Mercury* (Macrae et al., 2008), *PLATON* (Spek, 2009) and *publCIF* (Westrip, 2010).

## Acknowledgements

Special thanks are extended to Dr Guillermo Delgado Lamas (Universidad Nacional Autónoma de México) and Dr Eunice Ríos Vásquez [Universidad Nacional Autónoma de México and Universidad del Quindío (Colombia)] for assistance with the structural analyses. The authors also acknowledge financial support from the Universidad de Caldas, Colombia, and Illinois State University, USA.

## References

- Aakeröy, C. B. & Seddon, K. R. (1993). *Z. Naturforsch Teil B*, **48**, 1023–1025.
- Awwadi, F. F., Willett, R. D., Peterson, K. A. & Twamley, B. (2006). *Chem. Eur. J.* **12**, 8952–8960.
- Bruker (2012). *APEX2*, *SAINT*, *SADABS* and *TWINABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Calas, M., Ancelin, M. L., Cordina, G., Portefaix, P., Piquet, G., Vidal-Sailhan, V. & Vial, H. (2000). *J. Med. Chem.* **43**, 505–516.
- Calas, M., Cordina, G., Bompart, J., Ben Bari, M., Jei, T., Ancelin, M. L. & Vial, H. (1997). *J. Med. Chem.* **40**, 3557–3566.
- Calvani, M., Critelli, L., Gallo, G., Giorgi, F., Gramiccioli, G., Santaniello, M., Scafetta, N., Tinti, M. O. & De Angelis, F. (1998). *J. Med. Chem.* **41**, 2227–2233.
- Chen, C. C., Hsu, W., Hwang, K. C., Hwu, J. R., Lin, C. C. & Horng, J. (2011). *Arch. Biochem. Biophys.* **508**, 46–53.
- Desiraju, G. R., Ho, P. S., Kloos, L., Legon, A. C., Marquardt, R., Metrangolo, P., Politzer, P., Resnati, G. & Rissanen, K. (2013). *Pure Appl. Chem.* **85**, 1711–1713.
- Dougherty, D. A. (1996). *Science*, **271**, 163–168.
- Dougherty, D. A. (2013). *Acc. Chem. Res.* **46**, 885–893.
- Flack, H. D. & Bernardinelli, G. (1999). *Acta Cryst.* **A55**, 908–915.
- Formigué, M. (2009). *Curr. Opin. Solid State Mater. Sci.* **13**, 36–45.
- Glidewell, C., Zakaria, C. M., Ferguson, G. & Gallagher, J. F. (1994). *Acta Cryst.* **C50**, 233–238.
- Horner, J. H., Martínez, F. N., Musa, O. M., Newcomb, M. & Shahin, H. E. (1995). *J. Am. Chem. Soc.* **117**, 11124–11133.
- Ma, J. C. & Dougherty, D. A. (1997). *Chem. Rev.* **97**, 1303–1324.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Mavromoustakos, T., Calogeropoulou, T., Koufaki, M., Kolocouris, A., Daliani, I., Demetzos, D., Meng, Z., Makriyannis, A., Balzarini, J. & De Clercq, E. (2001). *J. Med. Chem.* **44**, 1702–1709.
- Múnera-Orozco, C. (2014). MSc thesis, Universidad de Caldas, Manizales, Colombia.
- Newcomb, M., Horner, J. H. & Shahin, H. (1993). *Tetrahedron Lett.* **34**, 5523–5526.
- Ng, K. L. C., Obando, D., Widmer, F., Wright, L. C., Sorrell, T. C. & Jolliffe, K. A. (2006). *J. Med. Chem.* **49**, 811–816.
- Nishio, M. (2012). *J. Mol. Struct.* **1018**, 2–7.

- Nishio, M., Hirota, M. & Umezawa, Y. (1998). In *The C–H...p Interaction. Evidence, Nature, and Consequences*. New York: Wiley–VCH.
- Ríos, L. A., Bartberger, M. D., Dolbier, W. R. & Paredes, R. (1997). *Tetrahedron Lett.* **38**, 7041–7044.
- Ríos, L. A., Dolbier, W. R., Paredes, R., Lusztyk, J., Ingold, K. U. & Jonsson, M. (1996). *J. Am. Chem. Soc.* **118**, 11313–11314.
- Ríos-Vásquez, L. A., Ocampo-Cardona, R., Duque-Benítez, S. M., Cedeño, D. L., Jones, M., Robledo-Restrepo, S. M. & Vélez-Bernal, I. D. (2015). US Patent No. 9,145,352 (September 29, 2015).
- Sheldrick, G. M. (2004). *CELL NOW*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Shiraishi, M., Aramaki, Y., Seto, M., Imoto, H., Nishikawa, Y., Kanzaki, N., Okamoto, M., Sawada, H., Nishimura, O., Baba, M. & Fujino, M. (2000). *J. Med. Chem.* **43**, 2049–2063.
- Song, D., Yang, J. S., Oh, C., Cui, S., Kim, B. K., Won, M., Lee, J. I., Kim, H. M. & Han, G. (2013). *Eur. J. Med. Chem.* **69**, 670–677.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Sussman, J. L., Harel, M., Frolow, F., Oefner, C., Goldman, A., Toker, L. & Silman, I. (1991). *Science*, **253**, 872–879.
- Wang, W., Bai, Z., Zhang, F., Wang, C., Yuan, Y. & Shao, J. (2012). *Eur. J. Med. Chem.* **56**, 320–331.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Yates, B. F., Bouma, W. J. & Radom, L. (1986). *Tetrahedron*, **42**, 6225–6234.

## supporting information

*Acta Cryst.* (2015). E71, 1230-1235 [doi:10.1107/S2056989015017181]

## Crystal structures of three new *N*-halomethylated quaternary ammonium salts

Carolina Múnera-Orozco, Rogelio Ocampo-Cardona, David L. Cedeño, Rubén A. Toscano and Luz Amalia Ríos-Vásquez

### Computing details

For all compounds, data collection: *APEX2* (Bruker, 2012); cell refinement: *S SAINT* (Bruker, 2012); data reduction: *S SAINT* (Bruker, 2012); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL2014* (Sheldrick, 2015), *PLATON* (Spek, 2009) and *publCIF* (Westrip, 2010).

### (I) *N*-(4,4-Diphenylbut-3-en-1-yl)-*N*-iodomethyl-*N,N*-dimethylammonium iodide

#### Crystal data

$C_{19}H_{23}IN^+I^-$

$M_r = 519.18$

Monoclinic,  $P2_1$

$a = 7.9254$  (2) Å

$b = 13.6161$  (3) Å

$c = 9.4632$  (2) Å

$\beta = 103.320$  (1)°

$V = 993.73$  (4) Å<sup>3</sup>

$Z = 2$

$F(000) = 500$

$D_x = 1.735$  Mg m<sup>-3</sup>

Melting point = 425–427 K

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 4987 reflections

$\theta = 2.2$ – $25.3$ °

$\mu = 3.16$  mm<sup>-1</sup>

$T = 298$  K

Prism, colourless

$0.23 \times 0.19 \times 0.12$  mm

#### Data collection

Bruker SMART APEX CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 8.333 pixels mm<sup>-1</sup>

$\omega$ -scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2012)

$T_{\min} = 0.624$ ,  $T_{\max} = 0.745$

5791 measured reflections

3085 independent reflections

3013 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.016$

$\theta_{\max} = 25.3$ °,  $\theta_{\min} = 2.2$ °

$h = -9 \rightarrow 9$

$k = -16 \rightarrow 16$

$l = -11 \rightarrow 11$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.017$

$wR(F^2) = 0.038$

$S = 1.08$

3085 reflections

202 parameters

1 restraint

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0138P)^2 + 0.0203P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.27 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.46 \text{ e } \text{\AA}^{-3}$

Absolute structure: Refined as an inversion twin  
 Absolute structure parameter: 0.02 (3)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refined as a 2-component inversion twin.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

|     | <i>x</i>     | <i>y</i>    | <i>z</i>    | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|-------------|-------------|----------------------------------|
| I1  | -0.65685 (3) | 0.06224 (2) | 0.13054 (3) | 0.04839 (9)                      |
| I2  | 0.06519 (4)  | 0.44189 (2) | 0.91487 (3) | 0.04368 (9)                      |
| N1  | -0.3770 (4)  | 0.2023 (3)  | 0.0555 (4)  | 0.0376 (8)                       |
| C1  | -0.1849 (5)  | 0.2322 (4)  | 0.0948 (5)  | 0.0402 (10)                      |
| H1A | -0.1144      | 0.1733      | 0.1019      | 0.048*                           |
| H1B | -0.1609      | 0.2719      | 0.0166      | 0.048*                           |
| C2  | -0.1311 (5)  | 0.2891 (3)  | 0.2358 (5)  | 0.0409 (11)                      |
| H2A | -0.1888      | 0.3524      | 0.2249      | 0.049*                           |
| H2B | -0.1676      | 0.2536      | 0.3124      | 0.049*                           |
| C3  | 0.0613 (6)   | 0.3044 (3)  | 0.2779 (5)  | 0.0382 (10)                      |
| H3  | 0.1101       | 0.3408      | 0.2143      | 0.046*                           |
| C4  | 0.1686 (5)   | 0.2711 (3)  | 0.3970 (4)  | 0.0329 (9)                       |
| C5  | 0.3545 (5)   | 0.3021 (3)  | 0.4380 (4)  | 0.0349 (9)                       |
| C6  | 0.4815 (6)   | 0.2388 (4)  | 0.5116 (5)  | 0.0457 (11)                      |
| H6  | 0.4514       | 0.1752      | 0.5318      | 0.055*                           |
| C7  | 0.6528 (6)   | 0.2687 (4)  | 0.5557 (5)  | 0.0546 (13)                      |
| H7  | 0.7362       | 0.2248      | 0.6040      | 0.065*                           |
| C8  | 0.7001 (6)   | 0.3612 (5)  | 0.5292 (6)  | 0.0598 (15)                      |
| H8  | 0.8148       | 0.3812      | 0.5610      | 0.072*                           |
| C9  | 0.5768 (7)   | 0.4253 (4)  | 0.4549 (7)  | 0.0661 (16)                      |
| H9  | 0.6087       | 0.4887      | 0.4357      | 0.079*                           |
| C10 | 0.4056 (6)   | 0.3959 (4)  | 0.4085 (6)  | 0.0532 (13)                      |
| H10 | 0.3239       | 0.4395      | 0.3569      | 0.064*                           |
| C11 | 0.1128 (5)   | 0.2030 (3)  | 0.5013 (4)  | 0.0326 (9)                       |
| C12 | 0.0439 (6)   | 0.1106 (3)  | 0.4575 (5)  | 0.0422 (10)                      |
| H12 | 0.0314       | 0.0909      | 0.3616      | 0.051*                           |
| C13 | -0.0060 (6)  | 0.0480 (3)  | 0.5561 (5)  | 0.0500 (12)                      |
| H13 | -0.0522      | -0.0134     | 0.5258      | 0.060*                           |
| C14 | 0.0124 (6)   | 0.0761 (4)  | 0.6978 (5)  | 0.0523 (13)                      |
| H14 | -0.0198      | 0.0334      | 0.7637      | 0.063*                           |
| C15 | 0.0784 (6)   | 0.1674 (4)  | 0.7431 (5)  | 0.0467 (12)                      |
| H15 | 0.0883       | 0.1871      | 0.8388      | 0.056*                           |
| C16 | 0.1298 (5)   | 0.2295 (4)  | 0.6457 (5)  | 0.0403 (10)                      |
| H16 | 0.1768       | 0.2905      | 0.6774      | 0.048*                           |
| C17 | -0.4036 (5)  | 0.1302 (3)  | 0.1685 (5)  | 0.0411 (10)                      |



|      |             |            |             |             |
|------|-------------|------------|-------------|-------------|
| H17A | -0.3821     | 0.1637     | 0.2614      | 0.049*      |
| H17B | -0.3174     | 0.0787     | 0.1761      | 0.049*      |
| C18  | -0.4928 (7) | 0.2892 (4) | 0.0510 (7)  | 0.0633 (15) |
| H18A | -0.6106     | 0.2703     | 0.0100      | 0.095*      |
| H18B | -0.4590     | 0.3397     | -0.0076     | 0.095*      |
| H18C | -0.4832     | 0.3135     | 0.1478      | 0.095*      |
| C19  | -0.4089 (7) | 0.1546 (5) | -0.0910 (5) | 0.0596 (14) |
| H19A | -0.5290     | 0.1369     | -0.1218     | 0.089*      |
| H19B | -0.3385     | 0.0967     | -0.0855     | 0.089*      |
| H19C | -0.3796     | 0.1997     | -0.1596     | 0.089*      |

*Atomic displacement parameters (Å<sup>2</sup>)*

|     | $U^{11}$     | $U^{22}$     | $U^{33}$     | $U^{12}$      | $U^{13}$     | $U^{23}$      |
|-----|--------------|--------------|--------------|---------------|--------------|---------------|
| I1  | 0.03965 (16) | 0.05161 (19) | 0.05521 (18) | -0.00709 (15) | 0.01359 (13) | -0.00180 (15) |
| I2  | 0.04705 (17) | 0.03979 (15) | 0.04515 (16) | 0.00372 (15)  | 0.01257 (12) | 0.00767 (14)  |
| N1  | 0.0317 (17)  | 0.043 (2)    | 0.0351 (19)  | -0.0020 (17)  | 0.0010 (15)  | 0.0047 (16)   |
| C1  | 0.029 (2)    | 0.051 (3)    | 0.039 (2)    | -0.005 (2)    | 0.0042 (18)  | 0.006 (2)     |
| C2  | 0.035 (2)    | 0.043 (3)    | 0.042 (3)    | 0.001 (2)     | 0.004 (2)    | -0.001 (2)    |
| C3  | 0.038 (2)    | 0.037 (2)    | 0.038 (2)    | -0.004 (2)    | 0.0058 (19)  | 0.003 (2)     |
| C4  | 0.034 (2)    | 0.031 (2)    | 0.033 (2)    | -0.0022 (18)  | 0.0073 (18)  | -0.0027 (18)  |
| C5  | 0.033 (2)    | 0.037 (2)    | 0.035 (2)    | -0.002 (2)    | 0.0087 (18)  | -0.0047 (19)  |
| C6  | 0.037 (3)    | 0.055 (3)    | 0.046 (3)    | 0.003 (2)     | 0.012 (2)    | 0.011 (2)     |
| C7  | 0.033 (2)    | 0.083 (4)    | 0.049 (3)    | 0.007 (3)     | 0.010 (2)    | 0.009 (3)     |
| C8  | 0.034 (3)    | 0.080 (4)    | 0.065 (4)    | -0.009 (3)    | 0.011 (3)    | -0.014 (3)    |
| C9  | 0.049 (3)    | 0.048 (4)    | 0.103 (4)    | -0.019 (3)    | 0.021 (3)    | -0.017 (3)    |
| C10 | 0.040 (3)    | 0.037 (2)    | 0.082 (4)    | -0.003 (2)    | 0.012 (3)    | 0.003 (3)     |
| C11 | 0.0266 (19)  | 0.035 (2)    | 0.035 (2)    | 0.0031 (18)   | 0.0054 (17)  | 0.0010 (18)   |
| C12 | 0.043 (2)    | 0.040 (2)    | 0.045 (2)    | -0.002 (2)    | 0.013 (2)    | -0.004 (2)    |
| C13 | 0.054 (3)    | 0.032 (3)    | 0.068 (3)    | -0.001 (2)    | 0.022 (2)    | 0.004 (2)     |
| C14 | 0.046 (3)    | 0.054 (3)    | 0.061 (3)    | 0.009 (3)     | 0.019 (2)    | 0.023 (3)     |
| C15 | 0.043 (3)    | 0.065 (3)    | 0.031 (2)    | 0.007 (3)     | 0.008 (2)    | 0.007 (2)     |
| C16 | 0.031 (2)    | 0.048 (3)    | 0.040 (2)    | -0.004 (2)    | 0.0043 (19)  | -0.004 (2)    |
| C17 | 0.035 (2)    | 0.049 (3)    | 0.037 (2)    | -0.005 (2)    | 0.0035 (18)  | 0.005 (2)     |
| C18 | 0.041 (3)    | 0.049 (3)    | 0.091 (4)    | 0.009 (3)     | -0.002 (3)   | 0.013 (3)     |
| C19 | 0.055 (3)    | 0.084 (4)    | 0.036 (3)    | -0.014 (3)    | 0.003 (2)    | 0.001 (3)     |

*Geometric parameters (Å, °)*

|                    |            |         |           |
|--------------------|------------|---------|-----------|
| I1—C17             | 2.164 (4)  | C8—H8   | 0.9300    |
| I1—I2 <sup>i</sup> | 3.5640 (4) | C9—C10  | 1.386 (7) |
| N1—C18             | 1.492 (6)  | C9—H9   | 0.9300    |
| N1—C19             | 1.499 (6)  | C10—H10 | 0.9300    |
| N1—C17             | 1.502 (5)  | C11—C16 | 1.390 (6) |
| N1—C1              | 1.536 (5)  | C11—C12 | 1.396 (6) |
| C1—C2              | 1.517 (6)  | C12—C13 | 1.387 (6) |
| C1—H1A             | 0.9700     | C12—H12 | 0.9300    |
| C1—H1B             | 0.9700     | C13—C14 | 1.370 (7) |

|                        |             |               |           |
|------------------------|-------------|---------------|-----------|
| C2—C3                  | 1.499 (6)   | C13—H13       | 0.9300    |
| C2—H2A                 | 0.9700      | C14—C15       | 1.379 (7) |
| C2—H2B                 | 0.9700      | C14—H14       | 0.9300    |
| C3—C4                  | 1.326 (6)   | C15—C16       | 1.379 (6) |
| C3—H3                  | 0.9300      | C15—H15       | 0.9300    |
| C4—C11                 | 1.493 (6)   | C16—H16       | 0.9300    |
| C4—C5                  | 1.495 (6)   | C17—H17A      | 0.9700    |
| C5—C6                  | 1.385 (6)   | C17—H17B      | 0.9700    |
| C5—C10                 | 1.388 (6)   | C18—H18A      | 0.9600    |
| C6—C7                  | 1.387 (7)   | C18—H18B      | 0.9600    |
| C6—H6                  | 0.9300      | C18—H18C      | 0.9600    |
| C7—C8                  | 1.353 (8)   | C19—H19A      | 0.9600    |
| C7—H7                  | 0.9300      | C19—H19B      | 0.9600    |
| C8—C9                  | 1.376 (8)   | C19—H19C      | 0.9600    |
|                        |             |               |           |
| C17—H1—H2 <sup>i</sup> | 176.81 (12) | C9—C10—C5     | 120.8 (5) |
| C18—N1—C19             | 110.2 (4)   | C9—C10—H10    | 119.6     |
| C18—N1—C17             | 110.7 (4)   | C5—C10—H10    | 119.6     |
| C19—N1—C17             | 110.7 (4)   | C16—C11—C12   | 118.0 (4) |
| C18—N1—C1              | 111.4 (4)   | C16—C11—C4    | 120.8 (4) |
| C19—N1—C1              | 106.6 (3)   | C12—C11—C4    | 121.2 (4) |
| C17—N1—C1              | 107.1 (3)   | C13—C12—C11   | 120.5 (4) |
| C2—C1—N1               | 114.2 (3)   | C13—C12—H12   | 119.8     |
| C2—C1—H1A              | 108.7       | C11—C12—H12   | 119.8     |
| N1—C1—H1A              | 108.7       | C14—C13—C12   | 120.3 (5) |
| C2—C1—H1B              | 108.7       | C14—C13—H13   | 119.9     |
| N1—C1—H1B              | 108.7       | C12—C13—H13   | 119.9     |
| H1A—C1—H1B             | 107.6       | C13—C14—C15   | 120.2 (4) |
| C3—C2—C1               | 111.6 (4)   | C13—C14—H14   | 119.9     |
| C3—C2—H2A              | 109.3       | C15—C14—H14   | 119.9     |
| C1—C2—H2A              | 109.3       | C14—C15—C16   | 119.7 (4) |
| C3—C2—H2B              | 109.3       | C14—C15—H15   | 120.2     |
| C1—C2—H2B              | 109.3       | C16—C15—H15   | 120.2     |
| H2A—C2—H2B             | 108.0       | C15—C16—C11   | 121.4 (4) |
| C4—C3—C2               | 126.3 (4)   | C15—C16—H16   | 119.3     |
| C4—C3—H3               | 116.8       | C11—C16—H16   | 119.3     |
| C2—C3—H3               | 116.8       | N1—C17—H1     | 116.0 (3) |
| C3—C4—C11              | 123.0 (4)   | N1—C17—H17A   | 108.3     |
| C3—C4—C5               | 121.6 (4)   | H1—C17—H17A   | 108.3     |
| C11—C4—C5              | 115.3 (3)   | N1—C17—H17B   | 108.3     |
| C6—C5—C10              | 117.5 (4)   | H1—C17—H17B   | 108.3     |
| C6—C5—C4               | 120.9 (4)   | H17A—C17—H17B | 107.4     |
| C10—C5—C4              | 121.6 (4)   | N1—C18—H18A   | 109.5     |
| C5—C6—C7               | 121.1 (5)   | N1—C18—H18B   | 109.5     |
| C5—C6—H6               | 119.4       | H18A—C18—H18B | 109.5     |
| C7—C6—H6               | 119.4       | N1—C18—H18C   | 109.5     |
| C8—C7—C6               | 120.7 (5)   | H18A—C18—H18C | 109.5     |
| C8—C7—H7               | 119.6       | H18B—C18—H18C | 109.5     |

|               |            |                 |            |
|---------------|------------|-----------------|------------|
| C6—C7—H7      | 119.6      | N1—C19—H19A     | 109.5      |
| C7—C8—C9      | 119.4 (5)  | N1—C19—H19B     | 109.5      |
| C7—C8—H8      | 120.3      | H19A—C19—H19B   | 109.5      |
| C9—C8—H8      | 120.3      | N1—C19—H19C     | 109.5      |
| C8—C9—C10     | 120.5 (5)  | H19A—C19—H19C   | 109.5      |
| C8—C9—H9      | 119.8      | H19B—C19—H19C   | 109.5      |
| C10—C9—H9     | 119.8      |                 |            |
|               |            |                 |            |
| C18—N1—C1—C2  | 55.2 (5)   | C6—C5—C10—C9    | -1.8 (8)   |
| C19—N1—C1—C2  | 175.5 (4)  | C4—C5—C10—C9    | 176.1 (5)  |
| C17—N1—C1—C2  | -65.9 (5)  | C3—C4—C11—C16   | -120.5 (5) |
| N1—C1—C2—C3   | 172.5 (4)  | C5—C4—C11—C16   | 57.8 (5)   |
| C1—C2—C3—C4   | -117.9 (5) | C3—C4—C11—C12   | 60.2 (6)   |
| C2—C3—C4—C11  | 6.6 (7)    | C5—C4—C11—C12   | -121.5 (4) |
| C2—C3—C4—C5   | -171.6 (4) | C16—C11—C12—C13 | 0.2 (6)    |
| C3—C4—C5—C6   | -148.1 (5) | C4—C11—C12—C13  | 179.5 (4)  |
| C11—C4—C5—C6  | 33.5 (5)   | C11—C12—C13—C14 | -0.3 (7)   |
| C3—C4—C5—C10  | 34.0 (6)   | C12—C13—C14—C15 | 0.9 (7)    |
| C11—C4—C5—C10 | -144.3 (4) | C13—C14—C15—C16 | -1.5 (7)   |
| C10—C5—C6—C7  | 0.9 (7)    | C14—C15—C16—C11 | 1.5 (7)    |
| C4—C5—C6—C7   | -177.0 (4) | C12—C11—C16—C15 | -0.8 (6)   |
| C5—C6—C7—C8   | 0.7 (7)    | C4—C11—C16—C15  | 179.9 (4)  |
| C6—C7—C8—C9   | -1.4 (8)   | C18—N1—C17—I1   | 64.6 (4)   |
| C7—C8—C9—C10  | 0.5 (9)    | C19—N1—C17—I1   | -57.9 (4)  |
| C8—C9—C10—C5  | 1.1 (9)    | C1—N1—C17—I1    | -173.8 (3) |

Symmetry code: (i)  $-x-1, y-1/2, -z+1$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

Cg is the centroid of the C11—C16 ring.

| $D-H\cdots A$                      | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|------------------------------------|-------|-------------|-------------|---------------|
| C17—H17B $\cdots$ I2 <sup>ii</sup> | 0.97  | 3.00        | 3.919 (5)   | 159           |
| C7—H7 $\cdots$ Cg <sup>iii</sup>   | 0.93  | 2.84        | 3.030 (5)   | 143           |

Symmetry codes: (ii)  $-x, y-1/2, -z+1$ ; (iii)  $x+1, y, z$ .

### (II) *N*-(5,5-Diphenylpent-4-en-1-yl)-*N*-iodomethyl-*N,N*-dimethylammonium iodide

#### Crystal data

$C_{20}H_{25}IN^+I^-$

$M_r = 533.21$

Monoclinic,  $C2/c$

$a = 37.778$  (7)  $\text{\AA}$

$b = 6.6323$  (12)  $\text{\AA}$

$c = 17.021$  (3)  $\text{\AA}$

$\beta = 100.567$  (4) $^\circ$

$V = 4192.3$  (13)  $\text{\AA}^3$

$Z = 8$

$F(000) = 2064$

$D_x = 1.690$   $\text{Mg m}^{-3}$

Melting point = 430–431 K

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$   $\text{\AA}$

Cell parameters from 4726 reflections

$\theta = 2.2$ – $25.3^\circ$

$\mu = 3.00$   $\text{mm}^{-1}$

$T = 298$  K

Platy-prism, colourless

$0.21 \times 0.20 \times 0.08$  mm

*Data collection*

Bruker SMART APEX CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$ -scans  
Absorption correction: multi-scan  
(*SADABS*; Bruker, 2012)  
 $T_{\min} = 0.349$ ,  $T_{\max} = 0.745$

16925 measured reflections  
3808 independent reflections  
3114 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.079$   
 $\theta_{\max} = 25.4^\circ$ ,  $\theta_{\min} = 2.2^\circ$   
 $h = -45 \rightarrow 45$   
 $k = -7 \rightarrow 7$   
 $l = -20 \rightarrow 20$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.052$   
 $wR(F^2) = 0.145$   
 $S = 1.05$   
3808 reflections  
210 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0918P)^2]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 1.90 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -1.98 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

|     | <i>x</i>     | <i>y</i>     | <i>z</i>    | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|--------------|-------------|----------------------------------|
| I1  | 0.25244 (2)  | -0.01229 (6) | 0.13640 (2) | 0.04488 (19)                     |
| I2  | 0.30512 (2)  | 0.07869 (7)  | 0.42246 (3) | 0.04999 (19)                     |
| N1  | 0.31227 (12) | 0.3200 (8)   | 0.1496 (2)  | 0.0383 (11)                      |
| C1  | 0.32956 (18) | 0.4989 (9)   | 0.1981 (4)  | 0.0446 (15)                      |
| H1A | 0.3463       | 0.5629       | 0.1690      | 0.053*                           |
| H1B | 0.3109       | 0.5965       | 0.2029      | 0.053*                           |
| C2  | 0.34941 (19) | 0.4459 (10)  | 0.2810 (4)  | 0.0503 (16)                      |
| H2A | 0.3713       | 0.3736       | 0.2770      | 0.060*                           |
| H2B | 0.3345       | 0.3583       | 0.3069      | 0.060*                           |
| C3  | 0.35877 (18) | 0.6363 (11)  | 0.3312 (4)  | 0.0498 (15)                      |
| H3A | 0.3769       | 0.7134       | 0.3107      | 0.060*                           |
| H3B | 0.3375       | 0.7199       | 0.3284      | 0.060*                           |
| C4  | 0.37268 (18) | 0.5764 (12)  | 0.4160 (4)  | 0.0536 (17)                      |
| H4  | 0.3577       | 0.4935       | 0.4394      | 0.064*                           |
| C5  | 0.40427 (16) | 0.6276 (11)  | 0.4628 (3)  | 0.0476 (15)                      |
| C6  | 0.41456 (18) | 0.5452 (12)  | 0.5446 (4)  | 0.0537 (18)                      |
| C7  | 0.4042 (2)   | 0.3523 (14)  | 0.5653 (4)  | 0.066 (2)                        |
| H7  | 0.3905       | 0.2719       | 0.5265      | 0.080*                           |
| C8  | 0.4140 (2)   | 0.2791 (17)  | 0.6421 (5)  | 0.083 (3)                        |
| H8  | 0.4068       | 0.1506       | 0.6544      | 0.099*                           |

|      |              |             |            |             |
|------|--------------|-------------|------------|-------------|
| C9   | 0.4346 (2)   | 0.396 (2)   | 0.7012 (5) | 0.095 (4)   |
| H9   | 0.4411       | 0.3468      | 0.7530     | 0.114*      |
| C10  | 0.4453 (2)   | 0.586 (2)   | 0.6822 (5) | 0.098 (4)   |
| H10  | 0.4592       | 0.6652      | 0.7213     | 0.117*      |
| C11  | 0.43528 (19) | 0.6603 (16) | 0.6054 (4) | 0.073 (2)   |
| H11  | 0.4425       | 0.7893      | 0.5937     | 0.088*      |
| C12  | 0.42943 (16) | 0.7710 (12) | 0.4341 (3) | 0.0494 (16) |
| C13  | 0.4190 (2)   | 0.9644 (13) | 0.4097 (5) | 0.067 (2)   |
| H13  | 0.3957       | 1.0067      | 0.4121     | 0.080*      |
| C14  | 0.4417 (3)   | 1.0942 (16) | 0.3824 (6) | 0.087 (3)   |
| H14  | 0.4339       | 1.2226      | 0.3655     | 0.104*      |
| C15  | 0.4761 (3)   | 1.036 (2)   | 0.3796 (7) | 0.097 (4)   |
| H15  | 0.4914       | 1.1257      | 0.3604     | 0.117*      |
| C16  | 0.4882 (2)   | 0.849 (2)   | 0.4045 (5) | 0.091 (3)   |
| H16  | 0.5118       | 0.8115      | 0.4026     | 0.109*      |
| C17  | 0.46478 (18) | 0.7137 (15) | 0.4330 (4) | 0.067 (2)   |
| H17  | 0.4729       | 0.5865      | 0.4509     | 0.080*      |
| C18  | 0.28176 (16) | 0.2484 (10) | 0.1869 (3) | 0.0425 (14) |
| H18A | 0.2648       | 0.3587      | 0.1853     | 0.051*      |
| H18B | 0.2911       | 0.2201      | 0.2428     | 0.051*      |
| C19  | 0.33945 (17) | 0.1549 (11) | 0.1464 (4) | 0.0509 (16) |
| H19A | 0.3288       | 0.0516      | 0.1101     | 0.076*      |
| H19B | 0.3468       | 0.0982      | 0.1988     | 0.076*      |
| H19C | 0.3601       | 0.2098      | 0.1283     | 0.076*      |
| C20  | 0.29821 (18) | 0.3908 (10) | 0.0658 (3) | 0.0495 (16) |
| H20A | 0.2870       | 0.2801      | 0.0343     | 0.074*      |
| H20B | 0.3178       | 0.4416      | 0.0427     | 0.074*      |
| H20C | 0.2808       | 0.4959      | 0.0668     | 0.074*      |

*Atomic displacement parameters (Å<sup>2</sup>)*

|     | $U^{11}$   | $U^{22}$   | $U^{33}$   | $U^{12}$      | $U^{13}$      | $U^{23}$     |
|-----|------------|------------|------------|---------------|---------------|--------------|
| I1  | 0.0501 (3) | 0.0520 (3) | 0.0289 (3) | -0.01023 (17) | -0.00240 (18) | 0.00087 (16) |
| I2  | 0.0512 (3) | 0.0512 (3) | 0.0461 (3) | 0.00415 (19)  | 0.0052 (2)    | 0.01397 (19) |
| N1  | 0.041 (3)  | 0.050 (3)  | 0.021 (2)  | -0.003 (2)    | -0.0006 (19)  | -0.002 (2)   |
| C1  | 0.052 (4)  | 0.047 (4)  | 0.033 (3)  | -0.013 (3)    | 0.001 (3)     | -0.003 (3)   |
| C2  | 0.052 (4)  | 0.062 (4)  | 0.031 (3)  | -0.008 (3)    | -0.009 (3)    | -0.005 (3)   |
| C3  | 0.044 (3)  | 0.062 (4)  | 0.039 (3)  | -0.010 (3)    | -0.003 (3)    | -0.011 (3)   |
| C4  | 0.047 (4)  | 0.082 (5)  | 0.028 (3)  | -0.012 (3)    | -0.001 (3)    | -0.011 (3)   |
| C5  | 0.042 (3)  | 0.067 (4)  | 0.031 (3)  | -0.009 (3)    | 0.000 (2)     | -0.019 (3)   |
| C6  | 0.035 (3)  | 0.089 (5)  | 0.036 (3)  | 0.001 (3)     | 0.004 (3)     | -0.008 (3)   |
| C7  | 0.064 (5)  | 0.088 (6)  | 0.046 (4)  | 0.000 (4)     | 0.008 (3)     | -0.002 (4)   |
| C8  | 0.066 (5)  | 0.126 (8)  | 0.058 (5)  | 0.013 (5)     | 0.018 (4)     | 0.021 (5)    |
| C9  | 0.051 (5)  | 0.188 (12) | 0.044 (5)  | 0.007 (6)     | 0.002 (4)     | 0.014 (6)    |
| C10 | 0.059 (5)  | 0.190 (12) | 0.039 (4)  | -0.033 (7)    | -0.004 (4)    | -0.024 (6)   |
| C11 | 0.057 (4)  | 0.118 (7)  | 0.042 (4)  | -0.023 (5)    | 0.001 (3)     | -0.021 (4)   |
| C12 | 0.040 (3)  | 0.075 (5)  | 0.029 (3)  | -0.008 (3)    | -0.003 (2)    | -0.016 (3)   |
| C13 | 0.057 (5)  | 0.075 (5)  | 0.061 (5)  | -0.012 (4)    | -0.007 (4)    | -0.020 (4)   |

|     |           |            |           |            |            |            |
|-----|-----------|------------|-----------|------------|------------|------------|
| C14 | 0.079 (6) | 0.092 (7)  | 0.079 (6) | -0.027 (5) | -0.014 (5) | 0.009 (5)  |
| C15 | 0.073 (7) | 0.136 (10) | 0.076 (6) | -0.044 (6) | -0.004 (5) | 0.025 (7)  |
| C16 | 0.047 (4) | 0.167 (11) | 0.056 (5) | -0.019 (6) | 0.004 (4)  | -0.005 (6) |
| C17 | 0.043 (4) | 0.111 (7)  | 0.042 (4) | -0.004 (4) | -0.002 (3) | -0.004 (4) |
| C18 | 0.044 (3) | 0.058 (4)  | 0.025 (3) | -0.009 (3) | 0.003 (2)  | -0.004 (3) |
| C19 | 0.048 (4) | 0.058 (4)  | 0.044 (3) | 0.005 (3)  | 0.002 (3)  | -0.011 (3) |
| C20 | 0.057 (4) | 0.061 (4)  | 0.028 (3) | -0.004 (3) | 0.000 (3)  | 0.006 (3)  |

*Geometric parameters (Å, °)*

|                        |             |             |            |
|------------------------|-------------|-------------|------------|
| I1—C18                 | 2.146 (6)   | C9—C10      | 1.381 (16) |
| I1—I2 <sup>i</sup>     | 3.5058 (7)  | C9—H9       | 0.9300     |
| N1—C18                 | 1.492 (7)   | C10—C11     | 1.382 (12) |
| N1—C20                 | 1.504 (7)   | C10—H10     | 0.9300     |
| N1—C19                 | 1.509 (8)   | C11—H11     | 0.9300     |
| N1—C1                  | 1.522 (8)   | C12—C13     | 1.383 (11) |
| C1—C2                  | 1.513 (9)   | C12—C17     | 1.392 (9)  |
| C1—H1A                 | 0.9700      | C13—C14     | 1.357 (12) |
| C1—H1B                 | 0.9700      | C13—H13     | 0.9300     |
| C2—C3                  | 1.530 (9)   | C14—C15     | 1.361 (14) |
| C2—H2A                 | 0.9700      | C14—H14     | 0.9300     |
| C2—H2B                 | 0.9700      | C15—C16     | 1.363 (16) |
| C3—C4                  | 1.496 (9)   | C15—H15     | 0.9300     |
| C3—H3A                 | 0.9700      | C16—C17     | 1.410 (13) |
| C3—H3B                 | 0.9700      | C16—H16     | 0.9300     |
| C4—C5                  | 1.351 (9)   | C17—H17     | 0.9300     |
| C4—H4                  | 0.9300      | C18—H18A    | 0.9700     |
| C5—C6                  | 1.480 (10)  | C18—H18B    | 0.9700     |
| C5—C12                 | 1.489 (9)   | C19—H19A    | 0.9600     |
| C6—C7                  | 1.402 (12)  | C19—H19B    | 0.9600     |
| C6—C11                 | 1.404 (10)  | C19—H19C    | 0.9600     |
| C7—C8                  | 1.379 (11)  | C20—H20A    | 0.9600     |
| C7—H7                  | 0.9300      | C20—H20B    | 0.9600     |
| C8—C9                  | 1.390 (14)  | C20—H20C    | 0.9600     |
| C8—H8                  | 0.9300      |             |            |
| C18—I1—I2 <sup>i</sup> | 170.16 (15) | C9—C10—C11  | 120.3 (9)  |
| C18—N1—C20             | 109.7 (4)   | C9—C10—H10  | 119.9      |
| C18—N1—C19             | 111.6 (5)   | C11—C10—H10 | 119.9      |
| C20—N1—C19             | 108.5 (5)   | C10—C11—C6  | 121.6 (9)  |
| C18—N1—C1              | 107.8 (4)   | C10—C11—H11 | 119.2      |
| C20—N1—C1              | 108.2 (5)   | C6—C11—H11  | 119.2      |
| C19—N1—C1              | 111.0 (5)   | C13—C12—C17 | 118.1 (7)  |
| C2—C1—N1               | 114.4 (5)   | C13—C12—C5  | 121.8 (6)  |
| C2—C1—H1A              | 108.6       | C17—C12—C5  | 120.1 (7)  |
| N1—C1—H1A              | 108.6       | C14—C13—C12 | 121.8 (8)  |
| C2—C1—H1B              | 108.6       | C14—C13—H13 | 119.1      |
| N1—C1—H1B              | 108.6       | C12—C13—H13 | 119.1      |



|               |            |                 |            |
|---------------|------------|-----------------|------------|
| H1A—C1—H1B    | 107.6      | C13—C14—C15     | 120.0 (10) |
| C1—C2—C3      | 110.7 (6)  | C13—C14—H14     | 120.0      |
| C1—C2—H2A     | 109.5      | C15—C14—H14     | 120.0      |
| C3—C2—H2A     | 109.5      | C14—C15—C16     | 121.0 (9)  |
| C1—C2—H2B     | 109.5      | C14—C15—H15     | 119.5      |
| C3—C2—H2B     | 109.5      | C16—C15—H15     | 119.5      |
| H2A—C2—H2B    | 108.1      | C15—C16—C17     | 119.3 (9)  |
| C4—C3—C2      | 108.9 (6)  | C15—C16—H16     | 120.3      |
| C4—C3—H3A     | 109.9      | C17—C16—H16     | 120.3      |
| C2—C3—H3A     | 109.9      | C12—C17—C16     | 119.7 (9)  |
| C4—C3—H3B     | 109.9      | C12—C17—H17     | 120.1      |
| C2—C3—H3B     | 109.9      | C16—C17—H17     | 120.1      |
| H3A—C3—H3B    | 108.3      | N1—C18—I1       | 117.9 (4)  |
| C5—C4—C3      | 128.2 (7)  | N1—C18—H18A     | 107.8      |
| C5—C4—H4      | 115.9      | I1—C18—H18A     | 107.8      |
| C3—C4—H4      | 115.9      | N1—C18—H18B     | 107.8      |
| C4—C5—C6      | 120.9 (6)  | I1—C18—H18B     | 107.8      |
| C4—C5—C12     | 121.0 (6)  | H18A—C18—H18B   | 107.2      |
| C6—C5—C12     | 118.1 (5)  | N1—C19—H19A     | 109.5      |
| C7—C6—C11     | 117.0 (7)  | N1—C19—H19B     | 109.5      |
| C7—C6—C5      | 122.5 (6)  | H19A—C19—H19B   | 109.5      |
| C11—C6—C5     | 120.5 (7)  | N1—C19—H19C     | 109.5      |
| C8—C7—C6      | 121.4 (8)  | H19A—C19—H19C   | 109.5      |
| C8—C7—H7      | 119.3      | H19B—C19—H19C   | 109.5      |
| C6—C7—H7      | 119.3      | N1—C20—H20A     | 109.5      |
| C7—C8—C9      | 120.4 (10) | N1—C20—H20B     | 109.5      |
| C7—C8—H8      | 119.8      | H20A—C20—H20B   | 109.5      |
| C9—C8—H8      | 119.8      | N1—C20—H20C     | 109.5      |
| C10—C9—C8     | 119.3 (8)  | H20A—C20—H20C   | 109.5      |
| C10—C9—H9     | 120.3      | H20B—C20—H20C   | 109.5      |
| C8—C9—H9      | 120.3      |                 |            |
| C18—N1—C1—C2  | 69.2 (7)   | C7—C6—C11—C10   | 0.4 (12)   |
| C20—N1—C1—C2  | -172.2 (5) | C5—C6—C11—C10   | -179.9 (8) |
| C19—N1—C1—C2  | -53.3 (7)  | C4—C5—C12—C13   | 57.8 (9)   |
| N1—C1—C2—C3   | -167.8 (5) | C6—C5—C12—C13   | -121.2 (7) |
| C1—C2—C3—C4   | 170.6 (6)  | C4—C5—C12—C17   | -123.6 (8) |
| C2—C3—C4—C5   | 124.9 (8)  | C6—C5—C12—C17   | 57.4 (8)   |
| C3—C4—C5—C6   | -176.7 (7) | C17—C12—C13—C14 | 2.5 (11)   |
| C3—C4—C5—C12  | 4.3 (12)   | C5—C12—C13—C14  | -179.0 (7) |
| C4—C5—C6—C7   | 32.1 (11)  | C12—C13—C14—C15 | -1.0 (14)  |
| C12—C5—C6—C7  | -148.9 (7) | C13—C14—C15—C16 | -0.4 (16)  |
| C4—C5—C6—C11  | -147.5 (7) | C14—C15—C16—C17 | 0.4 (16)   |
| C12—C5—C6—C11 | 31.5 (10)  | C13—C12—C17—C16 | -2.5 (10)  |
| C11—C6—C7—C8  | 0.0 (11)   | C5—C12—C17—C16  | 179.0 (6)  |
| C5—C6—C7—C8   | -179.7 (7) | C15—C16—C17—C12 | 1.1 (13)   |
| C6—C7—C8—C9   | 0.0 (12)   | C20—N1—C18—I1   | 64.9 (6)   |
| C7—C8—C9—C10  | -0.2 (14)  | C19—N1—C18—I1   | -55.4 (5)  |

|               |           |              |            |
|---------------|-----------|--------------|------------|
| C8—C9—C10—C11 | 0.6 (15)  | C1—N1—C18—I1 | -177.5 (4) |
| C9—C10—C11—C6 | -0.7 (14) |              |            |

Symmetry code: (i)  $-x+1/2, y-1/2, -z+1/2$ .

#### Hydrogen-bond geometry ( $\text{\AA}, ^\circ$ )

| $D-H\cdots A$      | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|--------------------|-------|-------------|-------------|---------------|
| C2—H2B $\cdots$ I2 | 0.97  | 3.06        | 4.001 (7)   | 165           |

### (III) *N*-(6,6-Diphenylhex-5-en-1-yl)-*N*-iodomethyl-*N,N*-dimethylammonium iodide

#### Crystal data

$C_{21}H_{27}IN^+I^-$

$M_r = 547.23$

Monoclinic,  $P2_1/c$

$a = 8.9423$  (12)  $\text{\AA}$

$b = 24.058$  (3)  $\text{\AA}$

$c = 10.3749$  (13)  $\text{\AA}$

$\beta = 103.656$  (3) $^\circ$

$V = 2168.9$  (5)  $\text{\AA}^3$

$Z = 4$

$F(000) = 1064$

$D_x = 1.676$   $\text{Mg m}^{-3}$

Melting point = 429–431 K

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$   $\text{\AA}$

Cell parameters from 3046 reflections

$\theta = 2.3\text{--}24.4^\circ$

$\mu = 2.90$   $\text{mm}^{-1}$

$T = 298$  K

Prism, colourless

$0.32 \times 0.22 \times 0.04$  mm

#### Data collection

Bruker SMART APEX CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 8.333 pixels  $\text{mm}^{-1}$

$\omega$ -scans

Absorption correction: multi-scan

(*TWINABS*; Bruker, 2012)

$T_{\min} = 0.273$ ,  $T_{\max} = 0.429$

3961 measured reflections

3961 independent reflections

2941 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.079$

$\theta_{\max} = 25.4^\circ$ ,  $\theta_{\min} = 1.7^\circ$

$h = -10 \rightarrow 10$

$k = 0 \rightarrow 28$

$l = 0 \rightarrow 12$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.060$

$wR(F^2) = 0.138$

$S = 1.05$

3961 reflections

220 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0515P)^2 + 2.9321P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.82$   $\text{e \AA}^{-3}$

$\Delta\rho_{\min} = -0.80$   $\text{e \AA}^{-3}$

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refined as a 2-component twin. The studied crystal was a nonmerohedral twin with a ratio of two major domains of 0.374 (2):0.626 (2). The two domains were rotated from each other by 180.0° about the reciprocal axis (1 0 0), which was determined by the CELL NOW program (Sheldrick, 2004). The final refinement was carried out using twinned data set.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)*

|     | <i>x</i>     | <i>y</i>    | <i>z</i>    | <i>U</i> <sub>iso</sub> */ <i>U</i> <sub>eq</sub> |
|-----|--------------|-------------|-------------|---|
| I1  | −0.29957 (8) | 0.44339 (3) | 0.01740 (6) | 0.0597 (2)  |
| I2  | 0.17494 (8)  | 0.51476 (3) | 0.26901 (6) | 0.0598 (2)  |
| N1  | −0.3411 (8)  | 0.4415 (4)  | 0.3070 (7)  | 0.056 (2)   |
| C1  | −0.1733 (11) | 0.4319 (4)  | 0.3631 (10) | 0.063 (3)   |
| H1A | −0.1159      | 0.4525      | 0.3103      | 0.076*  |
| H1B | −0.1459      | 0.4469      | 0.4524      | 0.076*  |
| C2  | −0.1252 (12) | 0.3731 (4)  | 0.3678 (12) | 0.072 (3)   |
| H2A | −0.1407      | 0.3584      | 0.2786      | 0.087*  |
| H2B | −0.1860      | 0.3512      | 0.4153      | 0.087*  |
| C3  | 0.0467 (11)  | 0.3699 (4)  | 0.4394 (11) | 0.068 (3)   |
| H3A | 0.1029       | 0.3971      | 0.4001      | 0.082*  |
| H3B | 0.0578       | 0.3800      | 0.5317      | 0.082*  |
| C4  | 0.1182 (12)  | 0.3134 (4)  | 0.4333 (15) | 0.091 (4)   |
| H4A | 0.1013       | 0.3019      | 0.3413      | 0.109*  |
| H4B | 0.0681       | 0.2866      | 0.4789      | 0.109*  |
| C5  | 0.2892 (10)  | 0.3135 (4)  | 0.4957 (12) | 0.068 (3)   |
| H5  | 0.3381       | 0.3478      | 0.5120      | 0.082*  |
| C6  | 0.3743 (10)  | 0.2685 (3)  | 0.5287 (10) | 0.055 (2)   |
| C7  | 0.5451 (11)  | 0.2703 (3)  | 0.5813 (9)  | 0.052 (2)   |
| C8  | 0.6398 (11)  | 0.2322 (4)  | 0.5435 (11) | 0.066 (3)   |
| H8  | 0.5966       | 0.2031      | 0.4881      | 0.080*  |
| C9  | 0.8003 (12)  | 0.2359 (5)  | 0.5862 (14) | 0.091 (4)   |
| H9  | 0.8636       | 0.2109      | 0.5560      | 0.109*  |
| C10 | 0.8623 (15)  | 0.2778 (5)  | 0.6747 (14) | 0.097 (5)   |
| H10 | 0.9683       | 0.2798      | 0.7079      | 0.116*  |
| C11 | 0.7701 (16)  | 0.3158 (5)  | 0.7133 (14) | 0.094 (4)   |
| H11 | 0.8128       | 0.3446      | 0.7699      | 0.113*  |
| C12 | 0.6122 (14)  | 0.3116 (5)  | 0.6681 (11) | 0.079 (3)   |
| H12 | 0.5495       | 0.3373      | 0.6969      | 0.094*  |
| C13 | 0.3026 (11)  | 0.2115 (3)  | 0.5086 (10) | 0.055 (2)   |
| C14 | 0.2723 (13)  | 0.1847 (4)  | 0.6156 (12) | 0.074 (3)   |
| H14 | 0.2993       | 0.2012      | 0.6990      | 0.089*  |
| C15 | 0.2004 (14)  | 0.1323 (4)  | 0.6000 (16) | 0.088 (4)   |
| H15 | 0.1840       | 0.1135      | 0.6738      | 0.105*  |
| C16 | 0.1551 (14)  | 0.1091 (5)  | 0.4782 (18) | 0.095 (4)   |
| H16 | 0.1010       | 0.0757      | 0.4681      | 0.114*  |
| C17 | 0.1888 (18)  | 0.1347 (6)  | 0.3678 (16) | 0.116 (6)   |
| H17 | 0.1614       | 0.1184      | 0.2842      | 0.139*  |
| C18 | 0.2659 (17)  | 0.1861 (5)  | 0.3880 (12) | 0.091 (4)   |
| H18 | 0.2926       | 0.2034      | 0.3164      | 0.109*  |

|      |              |            |             |           |
|------|--------------|------------|-------------|-----------|
| C19  | −0.4077 (11) | 0.4192 (4) | 0.1718 (10) | 0.065 (3) |
| H19A | −0.5148      | 0.4304     | 0.1460      | 0.078*    |
| H19B | −0.4058      | 0.3789     | 0.1771      | 0.078*    |
| C20  | −0.3648 (13) | 0.5048 (4) | 0.3059 (11) | 0.074 (3) |
| H20A | −0.4723      | 0.5131     | 0.2740      | 0.111*    |
| H20B | −0.3078      | 0.5218     | 0.2488      | 0.111*    |
| H20C | −0.3294      | 0.5191     | 0.3943      | 0.111*    |
| C21  | −0.4312 (12) | 0.4165 (5) | 0.3995 (10) | 0.071 (3) |
| H21A | −0.5382      | 0.4256     | 0.3684      | 0.107*    |
| H21B | −0.3936      | 0.4313     | 0.4872      | 0.107*    |
| H21C | −0.4189      | 0.3769     | 0.4015      | 0.107*    |

*Atomic displacement parameters (Å<sup>2</sup>)*

|     | $U^{11}$   | $U^{22}$   | $U^{33}$   | $U^{12}$   | $U^{13}$    | $U^{23}$    |
|-----|------------|------------|------------|------------|-------------|-------------|
| I1  | 0.0559 (4) | 0.0659 (4) | 0.0605 (4) | 0.0039 (3) | 0.0202 (3)  | 0.0118 (3)  |
| I2  | 0.0586 (4) | 0.0618 (4) | 0.0632 (4) | 0.0015 (3) | 0.0228 (3)  | 0.0043 (3)  |
| N1  | 0.041 (4)  | 0.081 (6)  | 0.046 (4)  | 0.002 (4)  | 0.012 (4)   | 0.020 (4)   |
| C1  | 0.045 (5)  | 0.078 (7)  | 0.069 (6)  | −0.006 (5) | 0.015 (5)   | 0.014 (5)   |
| C2  | 0.058 (6)  | 0.056 (6)  | 0.106 (9)  | 0.003 (5)  | 0.024 (6)   | −0.001 (6)  |
| C3  | 0.057 (6)  | 0.069 (6)  | 0.075 (7)  | 0.012 (5)  | 0.007 (6)   | 0.017 (6)   |
| C4  | 0.054 (7)  | 0.065 (7)  | 0.158 (12) | 0.006 (5)  | 0.036 (8)   | 0.023 (8)   |
| C5  | 0.040 (5)  | 0.043 (5)  | 0.123 (9)  | 0.002 (4)  | 0.022 (6)   | 0.012 (6)   |
| C6  | 0.047 (5)  | 0.038 (5)  | 0.082 (7)  | 0.005 (4)  | 0.019 (5)   | 0.010 (5)   |
| C7  | 0.052 (6)  | 0.038 (5)  | 0.069 (6)  | −0.002 (4) | 0.019 (5)   | 0.010 (4)   |
| C8  | 0.051 (6)  | 0.055 (6)  | 0.090 (7)  | 0.004 (5)  | 0.009 (6)   | −0.006 (5)  |
| C9  | 0.051 (7)  | 0.083 (8)  | 0.140 (11) | 0.017 (6)  | 0.026 (8)   | 0.022 (8)   |
| C10 | 0.066 (8)  | 0.077 (8)  | 0.129 (11) | −0.028 (7) | −0.014 (8)  | 0.022 (8)   |
| C11 | 0.100 (10) | 0.054 (7)  | 0.116 (11) | −0.008 (7) | −0.001 (9)  | −0.007 (7)  |
| C12 | 0.075 (8)  | 0.074 (7)  | 0.087 (8)  | −0.007 (6) | 0.019 (7)   | −0.002 (7)  |
| C13 | 0.056 (6)  | 0.040 (5)  | 0.067 (6)  | 0.012 (4)  | 0.011 (5)   | 0.014 (5)   |
| C14 | 0.079 (8)  | 0.053 (6)  | 0.101 (9)  | −0.004 (6) | 0.043 (7)   | −0.012 (6)  |
| C15 | 0.085 (9)  | 0.041 (6)  | 0.152 (13) | −0.009 (6) | 0.058 (9)   | 0.011 (7)   |
| C16 | 0.053 (7)  | 0.063 (7)  | 0.155 (14) | −0.001 (6) | −0.004 (9)  | −0.005 (10) |
| C17 | 0.135 (14) | 0.072 (9)  | 0.108 (11) | 0.001 (9)  | −0.038 (10) | −0.002 (8)  |
| C18 | 0.116 (11) | 0.069 (8)  | 0.075 (8)  | −0.001 (8) | −0.002 (7)  | 0.001 (7)   |
| C19 | 0.049 (6)  | 0.080 (7)  | 0.073 (7)  | −0.005 (5) | 0.029 (5)   | 0.024 (6)   |
| C20 | 0.066 (7)  | 0.071 (7)  | 0.089 (8)  | 0.009 (6)  | 0.027 (6)   | 0.008 (6)   |
| C21 | 0.056 (6)  | 0.091 (8)  | 0.072 (7)  | −0.006 (6) | 0.025 (6)   | 0.032 (6)   |

*Geometric parameters (Å, °)*

|                    |            |         |            |
|--------------------|------------|---------|------------|
| I1—C19             | 2.138 (9)  | C9—C10  | 1.388 (17) |
| I1—I2 <sup>i</sup> | 3.5565 (9) | C9—H9   | 0.9300     |
| N1—C19             | 1.489 (12) | C10—C11 | 1.353 (18) |
| N1—C1              | 1.494 (12) | C10—H10 | 0.9300     |
| N1—C21             | 1.515 (11) | C11—C12 | 1.383 (16) |
| N1—C20             | 1.537 (12) | C11—H11 | 0.9300     |

|                        |            |             |            |
|------------------------|------------|-------------|------------|
| C1—C2                  | 1.477 (13) | C12—H12     | 0.9300     |
| C1—H1A                 | 0.9700     | C13—C18     | 1.360 (14) |
| C1—H1B                 | 0.9700     | C13—C14     | 1.365 (14) |
| C2—C3                  | 1.543 (13) | C14—C15     | 1.407 (14) |
| C2—H2A                 | 0.9700     | C14—H14     | 0.9300     |
| C2—H2B                 | 0.9700     | C15—C16     | 1.353 (18) |
| C3—C4                  | 1.510 (14) | C15—H15     | 0.9300     |
| C3—H3A                 | 0.9700     | C16—C17     | 1.39 (2)   |
| C3—H3B                 | 0.9700     | C16—H16     | 0.9300     |
| C4—C5                  | 1.514 (14) | C17—C18     | 1.409 (17) |
| C4—H4A                 | 0.9700     | C17—H17     | 0.9300     |
| C4—H4B                 | 0.9700     | C18—H18     | 0.9300     |
| C5—C6                  | 1.323 (12) | C19—H19A    | 0.9700     |
| C5—H5                  | 0.9300     | C19—H19B    | 0.9700     |
| C6—C7                  | 1.497 (12) | C20—H20A    | 0.9600     |
| C6—C13                 | 1.507 (12) | C20—H20B    | 0.9600     |
| C7—C8                  | 1.366 (12) | C20—H20C    | 0.9600     |
| C7—C12                 | 1.379 (14) | C21—H21A    | 0.9600     |
| C8—C9                  | 1.401 (14) | C21—H21B    | 0.9600     |
| C8—H8                  | 0.9300     | C21—H21C    | 0.9600     |
|                        |            |             |            |
| C19—I1—I2 <sup>i</sup> | 171.6 (3)  | C11—C10—C9  | 120.6 (11) |
| C19—N1—C1              | 116.9 (8)  | C11—C10—H10 | 119.7      |
| C19—N1—C21             | 107.4 (7)  | C9—C10—H10  | 119.7      |
| C1—N1—C21              | 109.1 (7)  | C10—C11—C12 | 119.7 (12) |
| C19—N1—C20             | 109.1 (7)  | C10—C11—H11 | 120.1      |
| C1—N1—C20              | 106.3 (7)  | C12—C11—H11 | 120.1      |
| C21—N1—C20             | 107.8 (8)  | C7—C12—C11  | 121.7 (12) |
| C2—C1—N1               | 114.8 (8)  | C7—C12—H12  | 119.1      |
| C2—C1—H1A              | 108.6      | C11—C12—H12 | 119.1      |
| N1—C1—H1A              | 108.6      | C18—C13—C14 | 119.1 (9)  |
| C2—C1—H1B              | 108.6      | C18—C13—C6  | 122.5 (9)  |
| N1—C1—H1B              | 108.6      | C14—C13—C6  | 118.4 (9)  |
| H1A—C1—H1B             | 107.5      | C13—C14—C15 | 120.1 (12) |
| C1—C2—C3               | 108.2 (8)  | C13—C14—H14 | 119.9      |
| C1—C2—H2A              | 110.1      | C15—C14—H14 | 119.9      |
| C3—C2—H2A              | 110.1      | C16—C15—C14 | 120.4 (12) |
| C1—C2—H2B              | 110.1      | C16—C15—H15 | 119.8      |
| C3—C2—H2B              | 110.1      | C14—C15—H15 | 119.8      |
| H2A—C2—H2B             | 108.4      | C15—C16—C17 | 120.6 (11) |
| C4—C3—C2               | 114.0 (9)  | C15—C16—H16 | 119.7      |
| C4—C3—H3A              | 108.7      | C17—C16—H16 | 119.7      |
| C2—C3—H3A              | 108.7      | C16—C17—C18 | 117.3 (13) |
| C4—C3—H3B              | 108.7      | C16—C17—H17 | 121.3      |
| C2—C3—H3B              | 108.7      | C18—C17—H17 | 121.3      |
| H3A—C3—H3B             | 107.6      | C13—C18—C17 | 122.2 (13) |
| C3—C4—C5               | 112.1 (9)  | C13—C18—H18 | 118.9      |
| C3—C4—H4A              | 109.2      | C17—C18—H18 | 118.9      |

|                |             |                 |             |
|----------------|-------------|-----------------|-------------|
| C5—C4—H4A      | 109.2       | N1—C19—I1       | 117.2 (6)   |
| C3—C4—H4B      | 109.2       | N1—C19—H19A     | 108.0       |
| C5—C4—H4B      | 109.2       | I1—C19—H19A     | 108.0       |
| H4A—C4—H4B     | 107.9       | N1—C19—H19B     | 108.0       |
| C6—C5—C4       | 124.8 (9)   | I1—C19—H19B     | 108.0       |
| C6—C5—H5       | 117.6       | H19A—C19—H19B   | 107.2       |
| C4—C5—H5       | 117.6       | N1—C20—H20A     | 109.5       |
| C5—C6—C7       | 123.1 (8)   | N1—C20—H20B     | 109.5       |
| C5—C6—C13      | 120.7 (8)   | H20A—C20—H20B   | 109.5       |
| C7—C6—C13      | 116.2 (7)   | N1—C20—H20C     | 109.5       |
| C8—C7—C12      | 117.9 (10)  | H20A—C20—H20C   | 109.5       |
| C8—C7—C6       | 121.6 (9)   | H20B—C20—H20C   | 109.5       |
| C12—C7—C6      | 120.5 (9)   | N1—C21—H21A     | 109.5       |
| C7—C8—C9       | 121.6 (10)  | N1—C21—H21B     | 109.5       |
| C7—C8—H8       | 119.2       | H21A—C21—H21B   | 109.5       |
| C9—C8—H8       | 119.2       | N1—C21—H21C     | 109.5       |
| C10—C9—C8      | 118.4 (11)  | H21A—C21—H21C   | 109.5       |
| C10—C9—H9      | 120.8       | H21B—C21—H21C   | 109.5       |
| C8—C9—H9       | 120.8       |                 |             |
|                |             |                 |             |
| C19—N1—C1—C2   | -55.7 (12)  | C8—C7—C12—C11   | 1.8 (16)    |
| C21—N1—C1—C2   | 66.3 (12)   | C6—C7—C12—C11   | -176.9 (10) |
| C20—N1—C1—C2   | -177.7 (9)  | C10—C11—C12—C7  | -2 (2)      |
| N1—C1—C2—C3    | -174.9 (8)  | C5—C6—C13—C18   | 78.5 (15)   |
| C1—C2—C3—C4    | -171.1 (10) | C7—C6—C13—C18   | -98.8 (13)  |
| C2—C3—C4—C5    | 175.7 (10)  | C5—C6—C13—C14   | -100.9 (13) |
| C3—C4—C5—C6    | 165.3 (12)  | C7—C6—C13—C14   | 81.9 (12)   |
| C4—C5—C6—C7    | 175.5 (11)  | C18—C13—C14—C15 | -1.6 (16)   |
| C4—C5—C6—C13   | -1.5 (18)   | C6—C13—C14—C15  | 177.7 (9)   |
| C5—C6—C7—C8    | -140.0 (11) | C13—C14—C15—C16 | -2.7 (18)   |
| C13—C6—C7—C8   | 37.2 (13)   | C14—C15—C16—C17 | 4.7 (19)    |
| C5—C6—C7—C12   | 38.6 (15)   | C15—C16—C17—C18 | -2 (2)      |
| C13—C6—C7—C12  | -144.2 (10) | C14—C13—C18—C17 | 4.0 (19)    |
| C12—C7—C8—C9   | -2.7 (16)   | C6—C13—C18—C17  | -175.3 (11) |
| C6—C7—C8—C9    | 175.9 (10)  | C16—C17—C18—C13 | -2 (2)      |
| C7—C8—C9—C10   | 3.5 (18)    | C1—N1—C19—I1    | -54.3 (10)  |
| C8—C9—C10—C11  | -3.4 (19)   | C21—N1—C19—I1   | -177.2 (7)  |
| C9—C10—C11—C12 | 3 (2)       | C20—N1—C19—I1   | 66.3 (9)    |

Symmetry code: (i)  $-x, -y+1, -z$ .