



Crystal structure of 2-amino-4-(4-chlorophenyl)-1-(4-methylphenyl)-5-oxo-1,4,5,6,7,8-hexahydroquinoline-3-carbonitrile

Shaaban K. Mohamed,^{a,b} Mehmet Akkurt,^c Jerry P. Jasinski,^d Omya A. A. Abd Allah^e and Mustafa R. Albayati^{f*}

^aChemistry and Environmental Division, Manchester Metropolitan University, Manchester M1 5GD, England, ^bChemistry Department, Faculty of Science, Minia University, 61519 El-Minia, Egypt, ^cDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^dDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, ^eChemistry Department, Faculty of Science, Sohag University, 82524 Sohag, Egypt, and ^fKirkuk University, College of Science, Department of Chemistry, Kirkuk, Iraq. *Correspondence e-mail: shaabankamel@yahoo.com

Received 9 November 2015; accepted 10 November 2015

Edited by E. R. T. Tiekink, University of Malaya, Malaysia


In the title compound, C₂₃H₂₀ClN₃O, each of the cyclohexene and 1,4-dihydropyridine rings of the 1,4,5,6,7,8-hexahydroquinoline ring system adopts a twisted-boat conformation. The dihedral angle between the two benzene rings is 11.52 (7)°. In the crystal, molecules are linked through a pair of amino–nitrile N–H···N hydrogen bonds, forming inversion dimers. These assemble into a three-dimensional network via C–H···O and C–H···π interactions.

Keywords: crystal structure; dihydropyridines; annelated dihydropyridines; hydrogen bonding; inversion dimers.

CCDC reference: 1436038

1. Related literature

For the synthesis and pharmaceutical applications of dihydropyridines, see: Kumar & Maurya (2007); Kendre *et al.* (2008); Heydari *et al.* (2009).



2. Experimental

2.1. Crystal data

C₂₃H₂₀ClN₃O
M_r = 389.87
Monoclinic, P2₁/c
a = 8.7759 (3) Å
b = 10.6399 (3) Å
c = 20.7929 (7) Å
β = 93.842 (3)°

V = 1937.17 (11) Å³
Z = 4
Mo Kα radiation
μ = 0.22 mm⁻¹
T = 293 K
0.46 × 0.42 × 0.38 mm

2.2. Data collection

Agilent Xcalibur Eos Gemini
diffractometer
Absorption correction: multi-scan
(CrysAlis PRO; Agilent, 2014)
T_{min} = 0.841, T_{max} = 1.000

15538 measured reflections
6453 independent reflections
4853 reflections with I > 2σ(I)
R_{int} = 0.027

2.3. Refinement

R[F² > 2σ(F²)] = 0.053
wR(F²) = 0.144
S = 1.04
6453 reflections

254 parameters
H-atom parameters constrained
Δρ_{max} = 0.50 e Å⁻³
Δρ_{min} = -0.48 e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg3 and Cg4 are the centroids of the methyl- and chloro-benzene rings (C10–C15 and C18–C23), respectively.

D–H···A	D–H	H···A	D···A	D–H···A
N2–H2N···N3 ⁱ	0.91	2.17	3.013 (2)	153
C14–H14···O1 ⁱⁱ	0.93	2.58	3.4577 (19)	156
C3–H3A···Cg4 ⁱⁱⁱ	0.97	2.93	3.7210 (16)	139
C16–H16B···Cg3 ^{iv}	0.96	2.81	3.6464 (18)	146

Symmetry codes: (i) $-x, -y + 2, -z + 1$; (ii) $x, y - 1, z$; (iii) $x + 1, y, z$; (iv) $-x + 1, -y + 1, -z + 1$.

Data collection: CrysAlis PRO (Agilent, 2014); cell refinement: CrysAlis PRO; data reduction: CrysAlis PRO; program(s) used to solve structure: SHELXS2014 (Sheldrick, 2008); program(s) used to refine structure: SHELXL2014 (Sheldrick, 2015); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: PLATON (Spek, 2009).

Acknowledgements

JPJ acknowledges the NSF–MRI program (grant No. CHE-1039027) for funds to purchase the X-ray diffractometer.

Supporting information for this paper is available from the IUCr electronic archives (Reference: TK5407).

References

Agilent (2014). *CrysAlis PRO*. Agilent Technologies Ltd, Yarnton, England.
Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.

Heydari, A., Khaksar, S., Tajbakhsh, M. & Bijanzadeh, H. R. (2009). *J. Fluor. Chem.* **130**, 609–614.
Kendre, D. B., Toche, R. B. & Jachak, M. N. (2008). *J. Heterocycl. Chem.* **45**, 667–671.
Kumar, A. & Maurya, R. A. (2007). *Tetrahedron*, **63**, 1946–1952.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2015). E71, o949–o950 [https://doi.org/10.1107/S2056989015021313]

Crystal structure of 2-amino-4-(4-chlorophenyl)-1-(4-methylphenyl)-5-oxo-1,4,5,6,7,8-hexahydroquinoline-3-carbonitrile

Shaaban K. Mohamed, Mehmet Akkurt, Jerry P. Jasinski, Omya A. A. Abd Allah and Mustafa R. Albayati

S1. Comment

Dihydropyridines (DHP's) are a significant class of heterocyclic molecules due to their important biological activities (Kumar & Maurya, 2007). Dihydropyridine drugs, such as nifedipine, nicardipine and amlodipine are well known as cardiovascular agents for the treatment of hypertension (Kendre *et al.*, 2008), calcium channel modulators and for cardiovascular disease treatment (Heydari *et al.*, 2009). In this context, we report here the synthesis and crystal structure of the title compound.

In the title compound (Fig. 1), the cyclohexene (C1–C6) and 1,4-dihydropyridine (N1/C5–C9) rings of the 1,4,5,6,7,8-hexahydroquinoline ring system (N1/C1–C9) each adopt a twisted-boat conformation (the puckering parameters are $Q_T = 0.4813(16)$ Å, $\theta = 123.31(18)^\circ$, $\varphi = 283.9(2)^\circ$ and $Q_T = 0.2182(14)$ Å, $\theta = 103.7(4)^\circ$, $\varphi = 353.5(4)^\circ$, respectively). The dihedral angle between the methyl- and chloro-benzene rings is $11.52(7)^\circ$.

In the crystal, a pair of N—H \cdots N intermolecular hydrogen bonds connects two molecules by an $R_2^2(12)$ ring motif (Table 1), forming a centrosymmetric dimer (Fig. 2). Weak C—H \cdots O and C—H \cdots π interactions connect the dimers to each other, forming a three-dimensional network.

S2. Experimental

To a solution of 1.3-cyclohexanedione (3.36 g, 0.03 mol) in 40 ml ethanol, *p*-toluidine (3.21 g, 0.03 mol) and catalytic amount of triethylamine were added. The mixture was heated under reflux for 3 h. (4-Chlorobenzylidene)malononitrile (5.68 g, 0.03 mol) was added to the reaction mixture and refluxed for another 3 h. The separated solid was filtered off while hot, dried and crystallized from DMF as colourless crystals. Yield: 49.7%; m.p. 573 K, IR (λ_{\max} , cm^{-1}): 3472, 3325 (NH₂), 3209, 3032 (CH_{arom.}), 2972–2879 (CH_{aliph.}), 2177 (C \equiv N), 1631 (C=O); ¹H-NMR (DMSO-*d*₆), δ ppm: 7.39–7.29 (m, 8H, CH_{arom.}), 5.32 (s, 2H, NH₂, masked by D₂O), 4.51 (s, 1H, CH), 2.4 (s, 3H, CH₃), 2.23–2.19 (t, 2H, CH₂—C=O), 1.93–1.81 (t, 2H, CH₂—C=C), 1.67–1.62 (m, 2H, CH₂—CH₂—CH₂); ¹³C-NMR (DMSO-*d*₆), δ ppm: 195.45, 153.14, 151.70, 146.07, 139.86, 133.96, 131.26, 131.03, 130.10, 129.13, 128.79, 121.78, 112.72, 60.26, 36.45, 36.21, 28.23, 21.22, 21.06.

S3. Refinement

All H atoms were placed in calculated positions with N2—H1N = 0.90 Å, N2—H2N = 0.91 Å, and C—H = 0.93 - 1.04 Å, and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}(\text{C}, \text{N})$.




Figure 1

Molecular structure of the title compound with displacement ellipsoids drawn at the 30% probability level.




Figure 2

View of the dimers formed by N—H...N hydrogen bonds down the *b* axis.

2-Amino-4-(4-chlorophenyl)-1-(4-methylphenyl)-5-oxo-1,4,5,6,7,8-hexahydroquinoline-3-carbonitrile

Crystal data

$C_{23}H_{20}ClN_3O$

$M_r = 389.87$

Monoclinic, $P2_1/c$

Hall symbol: $-P 2_1/c$

$a = 8.7759 (3) \text{ \AA}$

$b = 10.6399 (3) \text{ \AA}$

$c = 20.7929 (7) \text{ \AA}$

$\beta = 93.842 (3)^\circ$

$V = 1937.17 (11) \text{ \AA}^3$

$Z = 4$

$F(000) = 816$

$D_x = 1.337 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4336 reflections

$\theta = 3.5\text{--}32.1^\circ$

$\mu = 0.22 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Irregular, colourless

$0.46 \times 0.42 \times 0.38 \text{ mm}$

Data collection

Agilent Xcalibur Eos Gemini diffractometer	15538 measured reflections
Radiation source: Enhance (Mo) X-ray Source	6453 independent reflections
Graphite monochromator	4853 reflections with $I > 2\sigma(I)$
Detector resolution: 16.0416 pixels mm ⁻¹	$R_{\text{int}} = 0.027$
ω scans	$\theta_{\text{max}} = 32.8^\circ$, $\theta_{\text{min}} = 3.5^\circ$
Absorption correction: multi-scan (CrysAlis PRO; Agilent, 2014)	$h = -13 \rightarrow 12$
$T_{\text{min}} = 0.841$, $T_{\text{max}} = 1.000$	$k = -15 \rightarrow 13$
	$l = -31 \rightarrow 29$

Refinement

Refinement on F^2	Hydrogen site location: mixed
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.053$	$w = 1/[\sigma^2(F_o^2) + (0.0563P)^2 + 0.8789P]$
$wR(F^2) = 0.144$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\text{max}} < 0.001$
6453 reflections	$\Delta\rho_{\text{max}} = 0.50 \text{ e } \text{\AA}^{-3}$
254 parameters	$\Delta\rho_{\text{min}} = -0.48 \text{ e } \text{\AA}^{-3}$
0 restraints	

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	-0.06881 (6)	1.08244 (7)	0.09288 (3)	0.0623 (2)
O1	0.50713 (13)	1.30415 (10)	0.28910 (6)	0.0326 (3)
N1	0.45035 (14)	0.91350 (11)	0.39374 (6)	0.0244 (3)
N2	0.26036 (17)	0.87555 (14)	0.46421 (7)	0.0396 (4)
N3	-0.00227 (17)	1.12750 (16)	0.44726 (8)	0.0423 (5)
C1	0.56260 (16)	1.20335 (12)	0.30796 (7)	0.0232 (3)
C2	0.72897 (16)	1.17417 (14)	0.30258 (8)	0.0276 (4)
C3	0.75310 (17)	1.03530 (14)	0.29019 (8)	0.0291 (4)
C4	0.69031 (16)	0.95720 (14)	0.34366 (8)	0.0282 (4)
C5	0.53218 (15)	0.99782 (12)	0.35801 (6)	0.0213 (3)
C6	0.47011 (15)	1.10911 (12)	0.33829 (6)	0.0206 (3)
C7	0.30491 (15)	1.14205 (12)	0.34566 (6)	0.0212 (3)
C8	0.24453 (15)	1.06023 (13)	0.39764 (6)	0.0221 (3)
C9	0.31330 (16)	0.95091 (13)	0.41841 (7)	0.0236 (3)
C10	0.50999 (16)	0.78952 (12)	0.40891 (7)	0.0223 (3)
C11	0.61730 (19)	0.77207 (14)	0.45979 (8)	0.0315 (4)
C12	0.67482 (19)	0.65278 (16)	0.47317 (8)	0.0340 (4)

C13	0.62493 (17)	0.54973 (14)	0.43663 (7)	0.0268 (4)
C14	0.51702 (19)	0.56892 (14)	0.38571 (7)	0.0296 (4)
C15	0.45943 (17)	0.68829 (14)	0.37163 (7)	0.0270 (4)
C16	0.6846 (2)	0.42010 (16)	0.45269 (8)	0.0369 (5)
C17	0.10810 (17)	1.09671 (14)	0.42500 (7)	0.0267 (4)
C18	0.21047 (15)	1.12918 (13)	0.28167 (7)	0.0215 (3)
C19	0.12277 (17)	1.22793 (14)	0.25608 (8)	0.0290 (4)
C20	0.03681 (18)	1.21482 (16)	0.19772 (8)	0.0351 (4)
C21	0.03801 (18)	1.10148 (18)	0.16578 (8)	0.0341 (5)
C22	0.12380 (19)	1.00150 (17)	0.19015 (8)	0.0334 (4)
C23	0.21025 (17)	1.01654 (14)	0.24800 (7)	0.0284 (4)
H1N	0.31260	0.80530	0.47550	0.0480*
H2A	0.78620	1.19830	0.34220	0.0330*
H2B	0.76720	1.22270	0.26760	0.0330*
H2N	0.17130	0.89510	0.48190	0.0480*
H3A	0.86130	1.01850	0.28810	0.0350*
H3B	0.70190	1.01210	0.24910	0.0350*
H4A	0.68820	0.86940	0.33100	0.0340*
H4B	0.75760	0.96510	0.38240	0.0340*
H7	0.29660	1.23540	0.36000	0.0250*
H11	0.65090	0.84010	0.48500	0.0380*
H12	0.74800	0.64170	0.50720	0.0410*
H14	0.48270	0.50090	0.36060	0.0350*
H15	0.38730	0.70000	0.33730	0.0320*
H16A	0.63750	0.36050	0.42290	0.0550*
H16B	0.66120	0.39850	0.49580	0.0550*
H16C	0.79330	0.41860	0.44960	0.0550*
H19	0.12140	1.30390	0.27820	0.0350*
H20	-0.02070	1.28170	0.18050	0.0420*
H22	0.12370	0.92540	0.16820	0.0400*
H23	0.26920	0.94990	0.26450	0.0340*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0483 (3)	0.0933 (5)	0.0424 (3)	0.0171 (3)	-0.0178 (2)	-0.0039 (3)
O1	0.0312 (6)	0.0215 (5)	0.0457 (6)	-0.0005 (4)	0.0073 (5)	0.0058 (4)
N1	0.0234 (6)	0.0192 (5)	0.0318 (6)	0.0035 (4)	0.0106 (4)	0.0033 (4)
N2	0.0389 (8)	0.0354 (7)	0.0476 (8)	0.0117 (6)	0.0251 (6)	0.0171 (6)
N3	0.0319 (7)	0.0531 (9)	0.0439 (8)	0.0113 (7)	0.0168 (6)	0.0056 (7)
C1	0.0228 (6)	0.0203 (6)	0.0267 (6)	-0.0028 (5)	0.0036 (5)	-0.0022 (5)
C2	0.0213 (6)	0.0253 (6)	0.0370 (8)	-0.0037 (5)	0.0069 (5)	0.0027 (6)
C3	0.0231 (7)	0.0277 (7)	0.0377 (8)	0.0004 (5)	0.0121 (6)	0.0016 (6)
C4	0.0222 (6)	0.0250 (6)	0.0383 (8)	0.0032 (5)	0.0095 (5)	0.0040 (6)
C5	0.0206 (6)	0.0200 (6)	0.0238 (6)	-0.0014 (5)	0.0058 (5)	-0.0011 (5)
C6	0.0190 (6)	0.0192 (6)	0.0242 (6)	-0.0001 (4)	0.0050 (4)	-0.0012 (5)
C7	0.0199 (6)	0.0179 (5)	0.0264 (6)	0.0012 (5)	0.0068 (5)	-0.0016 (5)
C8	0.0210 (6)	0.0237 (6)	0.0224 (6)	0.0007 (5)	0.0065 (5)	-0.0017 (5)

C9	0.0229 (6)	0.0235 (6)	0.0253 (6)	0.0004 (5)	0.0084 (5)	0.0001 (5)
C10	0.0235 (6)	0.0197 (6)	0.0242 (6)	0.0023 (5)	0.0063 (5)	0.0025 (5)
C11	0.0362 (8)	0.0255 (7)	0.0319 (7)	0.0001 (6)	-0.0049 (6)	-0.0045 (6)
C12	0.0364 (8)	0.0326 (8)	0.0315 (7)	0.0062 (6)	-0.0081 (6)	0.0014 (6)
C13	0.0302 (7)	0.0257 (6)	0.0249 (6)	0.0060 (5)	0.0054 (5)	0.0042 (5)
C14	0.0371 (8)	0.0230 (6)	0.0282 (7)	0.0024 (6)	-0.0007 (6)	-0.0032 (5)
C15	0.0296 (7)	0.0259 (7)	0.0250 (6)	0.0034 (5)	-0.0016 (5)	-0.0003 (5)
C16	0.0449 (10)	0.0304 (8)	0.0357 (8)	0.0127 (7)	0.0049 (7)	0.0060 (6)
C17	0.0242 (7)	0.0304 (7)	0.0260 (6)	0.0031 (5)	0.0054 (5)	0.0021 (5)
C18	0.0184 (6)	0.0213 (6)	0.0255 (6)	0.0017 (5)	0.0063 (5)	0.0028 (5)
C19	0.0257 (7)	0.0232 (6)	0.0385 (8)	0.0037 (5)	0.0059 (6)	0.0060 (6)
C20	0.0265 (7)	0.0384 (8)	0.0406 (8)	0.0081 (6)	0.0030 (6)	0.0150 (7)
C21	0.0234 (7)	0.0500 (10)	0.0287 (7)	0.0041 (7)	0.0007 (5)	0.0050 (7)
C22	0.0323 (8)	0.0378 (8)	0.0298 (7)	0.0057 (6)	0.0006 (6)	-0.0057 (6)
C23	0.0307 (7)	0.0267 (7)	0.0277 (7)	0.0074 (6)	0.0004 (5)	-0.0005 (5)

Geometric parameters (Å, °)

C11—C21	1.7399 (18)	C14—C15	1.391 (2)
O1—C1	1.2310 (17)	C18—C23	1.388 (2)
N1—C5	1.3943 (18)	C18—C19	1.387 (2)
N1—C9	1.3967 (19)	C19—C20	1.392 (2)
N1—C10	1.4462 (18)	C20—C21	1.377 (3)
N2—C9	1.351 (2)	C21—C22	1.380 (3)
N3—C17	1.149 (2)	C22—C23	1.388 (2)
C1—C2	1.504 (2)	C2—H2A	0.9700
C1—C6	1.4596 (19)	C2—H2B	0.9700
C2—C3	1.517 (2)	C3—H3A	0.9700
N2—H2N	0.9100	C3—H3B	0.9700
N2—H1N	0.9000	C4—H4A	0.9700
C3—C4	1.521 (2)	C4—H4B	0.9700
C4—C5	1.5024 (19)	C7—H7	1.0400
C5—C6	1.3554 (18)	C11—H11	0.9300
C6—C7	1.5093 (19)	C12—H12	0.9300
C7—C18	1.5259 (19)	C14—H14	0.9300
C7—C8	1.5112 (18)	C15—H15	0.9300
C8—C17	1.414 (2)	C16—H16A	0.9600
C8—C9	1.3670 (19)	C16—H16B	0.9600
C10—C15	1.383 (2)	C16—H16C	0.9600
C10—C11	1.381 (2)	C19—H19	0.9300
C11—C12	1.387 (2)	C20—H20	0.9300
C12—C13	1.388 (2)	C22—H22	0.9300
C13—C16	1.505 (2)	C23—H23	0.9300
C13—C14	1.388 (2)		
C5—N1—C9	119.95 (11)	C11—C21—C20	120.07 (14)
C5—N1—C10	120.75 (12)	C20—C21—C22	121.31 (15)
C9—N1—C10	119.20 (12)	C21—C22—C23	118.85 (16)

O1—C1—C2	121.42 (13)	C18—C23—C22	121.27 (14)
O1—C1—C6	121.11 (13)	C1—C2—H2A	109.00
C2—C1—C6	117.45 (12)	C1—C2—H2B	109.00
C1—C2—C3	111.15 (12)	C3—C2—H2A	109.00
C9—N2—H1N	119.00	C3—C2—H2B	109.00
C9—N2—H2N	120.00	H2A—C2—H2B	108.00
H1N—N2—H2N	121.00	C2—C3—H3A	110.00
C2—C3—C4	110.27 (13)	C2—C3—H3B	110.00
C3—C4—C5	111.89 (12)	C4—C3—H3A	110.00
N1—C5—C6	120.83 (12)	C4—C3—H3B	110.00
N1—C5—C4	115.87 (12)	H3A—C3—H3B	108.00
C4—C5—C6	123.31 (12)	C3—C4—H4A	109.00
C1—C6—C5	120.47 (12)	C3—C4—H4B	109.00
C1—C6—C7	116.75 (11)	C5—C4—H4A	109.00
C5—C6—C7	122.78 (12)	C5—C4—H4B	109.00
C6—C7—C18	111.10 (10)	H4A—C4—H4B	108.00
C8—C7—C18	112.08 (11)	C6—C7—H7	110.00
C6—C7—C8	108.91 (11)	C8—C7—H7	108.00
C7—C8—C17	118.46 (12)	C18—C7—H7	107.00
C7—C8—C9	123.20 (12)	C10—C11—H11	120.00
C9—C8—C17	118.31 (13)	C12—C11—H11	120.00
N1—C9—N2	115.79 (13)	C11—C12—H12	119.00
N1—C9—C8	119.95 (13)	C13—C12—H12	119.00
N2—C9—C8	124.22 (13)	C13—C14—H14	119.00
C11—C10—C15	120.10 (13)	C15—C14—H14	120.00
N1—C10—C11	120.53 (12)	C10—C15—H15	120.00
N1—C10—C15	119.37 (13)	C14—C15—H15	120.00
C10—C11—C12	119.80 (14)	C13—C16—H16A	109.00
C11—C12—C13	121.05 (15)	C13—C16—H16B	109.00
C12—C13—C14	118.40 (14)	C13—C16—H16C	109.00
C14—C13—C16	120.92 (14)	H16A—C16—H16B	110.00
C12—C13—C16	120.68 (14)	H16A—C16—H16C	109.00
C13—C14—C15	120.99 (14)	H16B—C16—H16C	109.00
C10—C15—C14	119.66 (13)	C18—C19—H19	120.00
N3—C17—C8	179.36 (17)	C20—C19—H19	120.00
C7—C18—C19	121.43 (13)	C19—C20—H20	120.00
C7—C18—C23	119.98 (12)	C21—C20—H20	120.00
C19—C18—C23	118.59 (14)	C21—C22—H22	121.00
C18—C19—C20	120.84 (14)	C23—C22—H22	121.00
C19—C20—C21	119.14 (15)	C18—C23—H23	119.00
C11—C21—C22	118.62 (14)	C22—C23—H23	119.00
C5—N1—C10—C11	81.13 (18)	C6—C7—C8—C17	162.97 (12)
C9—N1—C10—C11	-95.26 (17)	C18—C7—C8—C9	104.62 (15)
C5—N1—C10—C15	-98.38 (16)	C18—C7—C8—C17	-73.69 (15)
C5—N1—C9—N2	-165.14 (13)	C6—C7—C18—C23	55.07 (16)
C10—N1—C9—N2	11.28 (19)	C8—C7—C18—C19	112.49 (15)
C5—N1—C9—C8	12.5 (2)	C8—C7—C18—C23	-67.03 (16)

C10—N1—C9—C8	-171.05 (13)	C6—C7—C18—C19	-125.41 (14)
C9—N1—C5—C4	169.55 (13)	C7—C8—C9—N1	3.4 (2)
C10—N1—C5—C4	-6.82 (18)	C17—C8—C9—N1	-178.31 (13)
C9—N1—C5—C6	-10.23 (19)	C17—C8—C9—N2	-0.8 (2)
C10—N1—C5—C6	173.41 (12)	C7—C8—C9—N2	-179.15 (13)
C9—N1—C10—C15	85.23 (17)	N1—C10—C11—C12	-179.16 (14)
C6—C1—C2—C3	35.47 (18)	C15—C10—C11—C12	0.4 (2)
O1—C1—C6—C5	179.08 (13)	N1—C10—C15—C14	179.60 (13)
C2—C1—C6—C7	177.66 (12)	C11—C10—C15—C14	0.1 (2)
O1—C1—C2—C3	-146.41 (15)	C10—C11—C12—C13	-0.8 (2)
C2—C1—C6—C5	-2.80 (19)	C11—C12—C13—C14	0.7 (2)
O1—C1—C6—C7	-0.5 (2)	C11—C12—C13—C16	-178.22 (15)
C1—C2—C3—C4	-57.68 (17)	C12—C13—C14—C15	-0.3 (2)
C2—C3—C4—C5	47.83 (16)	C16—C13—C14—C15	178.66 (14)
C3—C4—C5—C6	-16.04 (19)	C13—C14—C15—C10	-0.1 (2)
C3—C4—C5—N1	164.19 (12)	C7—C18—C19—C20	-179.83 (14)
N1—C5—C6—C1	172.36 (12)	C23—C18—C19—C20	-0.3 (2)
N1—C5—C6—C7	-8.13 (19)	C7—C18—C23—C22	179.08 (14)
C4—C5—C6—C7	172.11 (12)	C19—C18—C23—C22	-0.5 (2)
C4—C5—C6—C1	-7.4 (2)	C18—C19—C20—C21	0.9 (2)
C1—C6—C7—C8	-159.39 (11)	C19—C20—C21—C11	179.82 (13)
C1—C6—C7—C18	76.69 (14)	C19—C20—C21—C22	-0.7 (2)
C5—C6—C7—C8	21.09 (17)	C11—C21—C22—C23	179.45 (12)
C5—C6—C7—C18	-102.84 (14)	C20—C21—C22—C23	-0.1 (2)
C6—C7—C8—C9	-18.72 (17)	C21—C22—C23—C18	0.6 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg3 and Cg4 are the centroids of the methyl- and chloro-benzene rings (C10–C15 and C18–C23), respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2N \cdots N3 ⁱ	0.91	2.17	3.013 (2)	153
C14—H14 \cdots O1 ⁱⁱ	0.93	2.58	3.4577 (19)	156
C3—H3A \cdots Cg4 ⁱⁱⁱ	0.97	2.93	3.7210 (16)	139
C16—H16B \cdots Cg3 ^{iv}	0.96	2.81	3.6464 (18)	146

Symmetry codes: (i) $-x, -y+2, -z+1$; (ii) $x, y-1, z$; (iii) $x+1, y, z$; (iv) $-x+1, -y+1, -z+1$.