

Crystal structure of tris(*trans*-1,2-cyclohexane-diamine- κ^2N,N')chromium(III) tetrachloridozincate chloride trihydrate from synchrotron data

Dohyun Moon^a and Jong-Ha Choi^{b*}

^aPohang Accelerator Laboratory, POSTECH, Pohang 37673, Republic of Korea, and ^bDepartment of Chemistry, Andong National University, Andong 36729, Republic of Korea. *Correspondence e-mail: jhchoi@anu.ac.kr

Received 1 April 2016
Accepted 7 April 2016

Edited by M. Weil, Vienna University of Technology, Austria

Keywords: crystal structure; 1,2-cyclohexanediamine; tetrachloridozincate chloride double salt; chromium(III) complex; hydrogen bonding; synchrotron radiation.


CCDC reference: 1472901

Supporting information: this article has supporting information at journals.iucr.org/e


The structure of the title double salt, $[\text{Cr}(\text{rac-chxn})_3][\text{ZnCl}_4]\text{Cl}\cdot 3\text{H}_2\text{O}$ (chxn is *trans*-1,2-cyclohexanediamine; $\text{C}_6\text{H}_{14}\text{N}_2$), has been determined from synchrotron data. The Cr^{III} ion is coordinated by six N atoms of three chelating chxn ligands, displaying a slightly distorted octahedral coordination environment. The distorted tetrahedral $[\text{ZnCl}_4]^{2-}$ anion, the isolated Cl^- anion and three lattice water molecules remain outside the coordination sphere. The $\text{Cr}-\text{N}(\text{chxn})$ bond lengths are in a narrow range between 2.0737 (12) and 2.0928 (12) Å; the mean $\text{N}-\text{Cr}-\text{N}$ bite angle is 82.1 (4)°. The crystal packing is stabilized by hydrogen-bonding interactions between the amino groups of the chxn ligands and the water molecules as donor groups, and O atoms of the water molecules, chloride anions and Cl atoms of the $[\text{ZnCl}_4]^{2-}$ anions as acceptor groups, leading to the formation of a three-dimensional network. The $[\text{ZnCl}_4]^{2-}$ anion is disordered over two sets of sites with an occupancy ratio of 0.94:0.06.

1. Chemical context

trans-1,2-Cyclohexanediamine (chxn) can coordinate to a central metal ion as a bidentate ligand *via* the two nitrogen atoms, forming a five-membered chelate ring. The synthetic procedures, crystal structures and detailed spectroscopic properties of such $[\text{Cr}(\text{chxn})_3]^{3+}$ complexes with chloride or nitrate anions have been reported previously (Morooka *et al.*, 1992; Choi, 1994; Kalf *et al.*, 2002). Since counter-anionic species play a very important role in coordination chemistry and supramolecular chemistry (Fabrizzi & Poggi, 2013; Santos-Figueroa *et al.*, 2013), changing the type of anion can also result in different structural properties. With respect to the tetrachloridozincate anion, $[\text{ZnCl}_4]^{2-}$, the crystal structures of complexes with trivalent chromium have been determined for $[\text{Cr}(\text{NH}_3)_6][\text{ZnCl}_4]\text{Cl}$ (Clegg, 1976), $[\text{Cr}(\text{en})_3][\text{ZnCl}_4]\text{Cl}$ (en is ethylenediamine; Pons *et al.*, 1988) and *trans*- $[\text{Cr}(\text{NH}_3)_2(\text{cyclam})][\text{ZnCl}_4]\text{Cl}\cdot\text{H}_2\text{O}$ (cyclam is 1,4,8,11-tetraazacyclotetradecane; Moon & Choi, 2016). However, a combination of the $[\text{Cr}(\text{chxn})_3]^{3+}$ cation with $[\text{ZnCl}_4]^{2-}$ and Cl^- as anions is unreported. In order to confirm that the resulting structure belongs to a double salt with $[\text{ZnCl}_4]^{2-}$ and Cl^- anions and does not contain a $[\text{ZnCl}_5]^{3-}$ anion, we prepared this material and report here on the molecular and crystal structure of $[\text{Cr}(\text{rac-chxn})_3][\text{ZnCl}_4]\text{Cl}\cdot 3\text{H}_2\text{O}$, (I).



OPEN ACCESS



2. Structural commentary

First of all we performed a single-crystal structure analysis of the starting complex $[\text{Cr}(\text{chxn})_3]\text{Cl}_3 \cdot 2\text{H}_2\text{O}$ with 98 K synchrotron data to determine the exact composition and coordination geometry of the Cr^{III} ion. The complex crystallizes in the space group $\bar{4}2d$ with eight formula units in a cell of dimensions $a = 18.893(3)$ and $c = 14.069(3)$ Å. The $\text{Cr}-\text{N}(\text{chxn})$ bond lengths are in the range 2.0723 (19) to 2.0937 (19) Å, and the $\text{N}-\text{Cr}-\text{N}$ bite angles are in the range 82.53 (7) to 82.69 (10)°. In comparison with the bond lengths and angles of the structure of this complex determined with 223 K data (Kalf *et al.*, 2002), there are no remarkable differences, and also the the crystal packing has virtually identical features.

Fig. 1 shows the molecular components of the title compound, (I), which consists of a discrete complex cation $[\text{Cr}(\text{rac-chxn})_3]^{3+}$, three lattice water molecules, together with one tetrahedral $[\text{ZnCl}_4]^{2-}$ and one isolated Cl^- counter-ion. The nitrogen atoms of the three 1,2-cyclohexanediamine ligands define a distorted octahedral coordination environment around the $\text{Cr}(\text{III})$ ion with a mean $\text{N}-\text{Cr}-\text{N}$ bite angle of 82.1 (4)°. The resulting five-membered chelate rings of chxn ligands have the expected stable *gauche* conformation. The $\text{Cr}-\text{N}(\text{chxn})$ bond lengths are in the range 2.0737 (12) to 2.0928 (12) Å, in good agreement with those determined in




Figure 1

The structures of the molecular components of the title double salt, drawn with displacement parameters at the 50% probability level. Dashed lines represent hydrogen-bonding interactions.

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N1}-\text{H1A} \cdots \text{Cl5}^{\text{i}}$	0.91	2.40	3.2535 (15)	157
$\text{N1}-\text{H1B} \cdots \text{O3W}$	0.91	2.36	3.0178 (16)	129
$\text{N2}-\text{H2A} \cdots \text{O2W}$	0.91	2.01	2.9051 (17)	166
$\text{N2}-\text{H2B} \cdots \text{Cl2A}^{\text{ii}}$	0.91	2.45	3.2197 (14)	142
$\text{N2}-\text{H2B} \cdots \text{Cl3B}^{\text{ii}}$	0.91	2.36	3.180 (18)	150
$\text{N3}-\text{H3A} \cdots \text{O2W}$	0.91	2.13	2.9832 (16)	156
$\text{N3}-\text{H3B} \cdots \text{Cl1A}^{\text{iii}}$	0.91	2.52	3.2574 (13)	138
$\text{N3}-\text{H3B} \cdots \text{Cl3A}^{\text{iii}}$	0.91	2.77	3.4547 (16)	133
$\text{N3}-\text{H3B} \cdots \text{Cl2B}^{\text{iii}}$	0.91	2.67	3.471 (10)	147
$\text{N3}-\text{H3B} \cdots \text{Cl4B}^{\text{iii}}$	0.91	2.68	3.35 (2)	131
$\text{N4}-\text{H4A} \cdots \text{Cl1B}^{\text{iv}}$	0.91	2.74	3.473 (11)	138
$\text{N4}-\text{H4B} \cdots \text{Cl2A}^{\text{ii}}$	0.91	2.64	3.4267 (15)	146
$\text{N4}-\text{H4B} \cdots \text{O1W}^{\text{ii}}$	0.91	2.39	2.9804 (17)	123
$\text{N5}-\text{H5A} \cdots \text{Cl3A}^{\text{iv}}$	0.91	2.51	3.4245 (14)	178
$\text{N5}-\text{H5A} \cdots \text{Cl4B}^{\text{iv}}$	0.91	2.73	3.634 (19)	173
$\text{N5}-\text{H5B} \cdots \text{Cl1A}^{\text{iii}}$	0.91	2.74	3.3664 (16)	127
$\text{N5}-\text{H5B} \cdots \text{O3W}$	0.91	2.22	2.9724 (17)	140
$\text{N6}-\text{H6A} \cdots \text{Cl5}^{\text{i}}$	0.91	2.39	3.2474 (14)	158
$\text{O1W}-\text{H1O1} \cdots \text{Cl5}$	0.85 (1)	2.24 (1)	3.0878 (17)	179 (2)
$\text{O1W}-\text{H2O1} \cdots \text{Cl4A}^{\text{ii}}$	0.84 (1)	2.28 (1)	3.1170 (13)	174 (2)
$\text{O2W}-\text{H1O2} \cdots \text{Cl1A}$	0.83 (1)	2.28 (1)	3.1140 (12)	175 (2)
$\text{O2W}-\text{H1O2} \cdots \text{Cl2B}$	0.83 (1)	2.45 (1)	3.271 (9)	167 (2)
$\text{O2W}-\text{H2O2} \cdots \text{O1W}$	0.83 (1)	1.92 (1)	2.7468 (19)	177 (2)
$\text{O3W}-\text{H1O3} \cdots \text{Cl5}^{\text{iii}}$	0.84 (1)	2.41 (1)	3.2139 (13)	159 (2)
$\text{O3W}-\text{H2O3} \cdots \text{Cl2A}^{\text{iii}}$	0.84 (1)	2.38 (1)	3.2153 (17)	175 (2)
$\text{O3W}-\text{H2O3} \cdots \text{Cl3B}^{\text{iii}}$	0.84 (1)	2.23 (2)	3.05 (2)	167 (2)

Symmetry codes: (i) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (ii) $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iv) $x, y + 1, z$.

$[\text{Cr}(\text{RR-chxn})_3](\text{NO}_3)_3 \cdot 3\text{H}_2\text{O}$ (Morooka *et al.*, 1992) and $[\text{Cr}(\text{rac-chxn})_3]\text{Cl}_3 \cdot 2\text{H}_2\text{O}$ (Kalf *et al.*, 2002). The disordered tetrahedral $[\text{ZnCl}_4]^{2-}$ anion, the isolated Cl^- anion and the three water molecules remain outside the coordination sphere of Cr^{III} . The complex $[\text{ZnCl}_4]^{2-}$ anion is distorted due to its involvement in hydrogen-bonding interactions. The $[\text{ZnCl}_4]^{2-}$ and Cl^- anions are well separated by van der Waals contacts and consequently there is no basis for describing the Zn^{II} species as a distorted $[\text{ZnCl}_5]^{3-}$ anion.

3. Supramolecular features

Extensive hydrogen-bonding interactions occur in the crystal structure (Table 1), involving the $\text{N}-\text{H}$ groups of the chxn ligands and the $\text{O}-\text{H}$ groups of the lattice water molecules as donors, and the chloride ions and Cl atoms of the disordered $[\text{ZnCl}_4]^{2-}$ anions and water O atoms as acceptors. The supramolecular architecture gives rise to a three-dimensional network structure (Fig. 2).

4. Database survey

A search of the Cambridge Structural Database (Version 5.36, May 2015 with last update; Groom *et al.*, 2016) shows that there are three previous reports for Cr^{III} complexes containing three chelating chxn ligands, *viz.* $[\text{Cr}(\text{RR-chxn})_3](\text{NO}_3)_3 \cdot 3\text{H}_2\text{O}$ (Morooka *et al.*, 1992), $[\text{Cr}(\text{rac-chxn})_3]\text{Cl}_3 \cdot 2\text{H}_2\text{O}$ (Kalf *et al.*, 2002), and $[\text{Cr}(\text{RR-chxn})_3][\text{Co}(\text{SS-chxn})_3]\text{Cl}_6 \cdot 4\text{H}_2\text{O}$ (Kalf *et al.*, 2002). The structure of any double salt of $[\text{Cr}(\text{chxn})_3]^{3+}$ with an additional $[\text{ZnCl}_4]^{2-}$ anion has not been deposited.




Figure 2

The crystal packing in the title double salt viewed perpendicular to the bc plane. Dashed lines represent hydrogen-bonding interactions: $\text{N-H}\cdots\text{Cl}$ (pink), $\text{N-H}\cdots\text{O}$ (cyan), $\text{O-H}\cdots\text{O}$ (light green) and $\text{O-H}\cdots\text{Cl}$ (orange). The minor disorder components of the $[\text{ZnCl}_4]^{2-}$ anion have been omitted for clarity.

5. Synthesis and crystallization

Commercially available (Aldrich) *racemic trans*-1,2-cyclohexanediamine was used as provided. All other chemicals with the best analytical grade available were used. The starting material, $[\text{Cr}(\text{rac-chxn})_3]\text{Cl}_3\cdot 2\text{H}_2\text{O}$ was prepared according to the literature (Pedersen, 1970). The crude trichloride salt (0.22 g) was dissolved in 10 mL of 1 M HCl at 313 K and 5 mL of 1 M HCl containing 0.5 g of solid ZnCl_2 were added to this solution. The resulting solution was filtered and allowed to stand at room temperature for one week to give block-like yellow crystals of the tetrachloridozincate(II) chloride salt suitable for X-ray structural analysis.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. All H atoms were found from difference maps but were placed in geometrically idealized positions and constrained to ride on their parent atoms, with $\text{C-H} = 0.99\text{--}1.00$ Å and $\text{N-H} = 0.91$ Å, and with $U_{\text{iso}}(\text{H})$ values of 1.2 or $1.5U_{\text{eq}}$ of the parent atoms. The hydrogen atoms of water molecules were restrained using DFIX and DANG commands during the least-squares refinement (Sheldrick, 2015b). The $[\text{ZnCl}_4]^{2-}$ anion was refined as positionally disordered over two sets of sites with a refined occupancy ratio constrained to 0.94:0.06 in the last refinement cycles.

Table 2

Experimental details.

Crystal data	
Chemical formula	$[\text{Cr}(\text{C}_6\text{H}_{14}\text{N}_2)_3][\text{ZnCl}_4]\text{Cl}\cdot 3\text{H}_2\text{O}$
M_r	691.24
Crystal system, space group	Monoclinic, $P2_1/c$
Temperature (K)	100
a, b, c (Å)	10.594 (2), 13.075 (3), 22.384 (5)
β (°)	100.87 (3)
V (Å ³)	3045.0 (11)
Z	4
Radiation type	Synchrotron, $\lambda = 0.62998$ Å
μ (mm ⁻¹)	1.15
Crystal size (mm)	$0.25 \times 0.15 \times 0.05$
Data collection	
Diffractionmeter	ADSC Q210 CCD area detector
Absorption correction	Empirical (using intensity measurements) (<i>HKL3000sm SCALEPACK</i> ; Otwinowski & Minor, 1997)
$T_{\text{min}}, T_{\text{max}}$	0.762, 0.945
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	23113, 8090, 7647
R_{int}	0.034
$(\sin \theta/\lambda)_{\text{max}}$ (Å ⁻¹)	0.696
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.027, 0.073, 1.05
No. of reflections	8090
No. of parameters	371
No. of restraints	15
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ⁻³)	1.07, -1.14

Computer programs: *PAL BL2D-SMDC* (Shin *et al.*, 2016), *HKL3000sm* (Otwinowski & Minor, 1997), *SHELXT2014* (Sheldrick, 2015a), *SHELXL2014* (Sheldrick, 2015b), *DIAMOND* (Putz & Brandenburg, 2014) and *pubCIF* (Westrip, 2010).

Acknowledgements

This work was supported by a grant from 2016 Research Funds of Andong National University. The X-ray crystallography experiment at PLS-II BL2D-SMC beamline was supported in part by MSIP and POSTECH.

References

- Choi, J.-H. (1994). *Bull. Korean Chem. Soc.* **15**, 145–150.
- Clegg, W. (1976). *Acta Cryst.* **B32**, 2907–2909.
- Fabbrizzi, L. & Poggi, A. (2013). *Chem. Soc. Rev.* **42**, 1681–1699.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Kalf, I., Calmuschi, B. & Englert, U. (2002). *CrystEngComm*, **4**, 548–551.
- Moon, D. & Choi, J.-H. (2016). *Acta Cryst.* **E72**, 456–459.
- Morooka, M., Ohba, S. & Miyamae, H. (1992). *Acta Cryst.* **B48**, 667–672.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Pedersen, E. (1970). *Acta Chem. Scand.* **24**, 3362–3372.
- Pons, J., Casabó, J., Palacio, F., Morón, M. C., Solans, X. & Carlin, R. L. (1988). *Inorg. Chim. Acta*, **146**, 161–165.
- Putz, H. & Brandenburg, K. (2014). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Santos-Figueroa, L. E., Moragues, M. E., Climent, E., Agostini, A., Martínez-Mañez, R. & Sancenón, F. (2013). *Chem. Soc. Rev.* **42**, 3489–3613.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Shin, J. W., Eom, K. & Moon, D. (2016). *J. Synchrotron Rad.* **23**, 369–373.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2016). E72, 671-674 [doi:10.1107/S2056989016005788]

Crystal structure of tris(*trans*-1,2-cyclohexanediamine- κ^2N,N')chromium(III) tetrachloridozincate chloride trihydrate from synchrotron data

Dohyun Moon and Jong-Ha Choi

Computing details

Data collection: *PAL BL2D-SMDC* (Shin *et al.*, 2016); cell refinement: *HKL3000sm* (Otwinowski & Minor, 1997); data reduction: *HKL3000sm* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXT2014* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015b); molecular graphics: *DIAMOND* (Putz & Brandenburg, 2014); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Tris(*trans*-1,2-cyclohexanediamine- κ^2N,N')chromium(III) tetrachloridozincate chloride trihydrate

Crystal data

$[\text{Cr}(\text{C}_6\text{H}_{14}\text{N}_2)_3][\text{ZnCl}_4]\text{Cl}\cdot 3\text{H}_2\text{O}$

$M_r = 691.24$

Monoclinic, $P2_1/c$

$a = 10.594$ (2) Å

$b = 13.075$ (3) Å

$c = 22.384$ (5) Å

$\beta = 100.87$ (3)°

$V = 3045.0$ (11) Å³

$Z = 4$

$F(000) = 1444$

$D_x = 1.508$ Mg m⁻³

Synchrotron radiation, $\lambda = 0.62998$ Å

Cell parameters from 39315 reflections

$\theta = 0.4\text{--}33.6^\circ$

$\mu = 1.15$ mm⁻¹

$T = 100$ K

Block, yellow

0.25 × 0.15 × 0.05 mm

Data collection

ADSC Q210 CCD area detector
diffractometer

Radiation source: PLSII 2D bending magnet
 ω scan

Absorption correction: empirical (using
intensity measurements)

(*HKL3000sm SCALEPACK*; Otwinowski &
Minor, 1997)

$T_{\min} = 0.762$, $T_{\max} = 0.945$

23113 measured reflections

8090 independent reflections

7647 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.034$

$\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 1.7^\circ$

$h = -14 \rightarrow 13$

$k = -18 \rightarrow 18$

$l = -31 \rightarrow 23$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.027$

$wR(F^2) = 0.073$

$S = 1.05$

8090 reflections

371 parameters

15 restraints

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0353P)^2 + 1.582P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 1.07$ e Å⁻³

$\Delta\rho_{\min} = -1.14$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cr1	0.74469 (2)	0.93268 (2)	0.34326 (2)	0.00945 (5)	
N1	0.64213 (11)	0.83929 (8)	0.39272 (5)	0.0148 (2)	
H1A	0.6540	0.8610	0.4320	0.018*	
H1B	0.5567	0.8427	0.3766	0.018*	
N2	0.86651 (10)	0.80619 (8)	0.35251 (5)	0.01418 (19)	
H2A	0.8570	0.7719	0.3166	0.017*	
H2B	0.9497	0.8270	0.3629	0.017*	
N3	0.64039 (10)	0.87915 (8)	0.26146 (5)	0.01276 (18)	
H3A	0.6636	0.8136	0.2552	0.015*	
H3B	0.5550	0.8799	0.2625	0.015*	
N4	0.84308 (11)	1.01181 (8)	0.28504 (5)	0.01452 (19)	
H4A	0.8202	1.0790	0.2836	0.017*	
H4B	0.9293	1.0076	0.2991	0.017*	
N5	0.61717 (10)	1.05117 (7)	0.35045 (5)	0.01305 (18)	
H5A	0.6204	1.0989	0.3212	0.016*	
H5B	0.5355	1.0265	0.3453	0.016*	
N6	0.85476 (11)	1.00563 (8)	0.41871 (5)	0.0163 (2)	
H6A	0.8540	0.9680	0.4529	0.020*	
H6B	0.9376	1.0115	0.4135	0.020*	
C1	0.68832 (12)	0.73168 (8)	0.39046 (6)	0.0125 (2)	
H1	0.6558	0.7035	0.3489	0.015*	
C2	0.64271 (14)	0.66233 (9)	0.43714 (6)	0.0179 (2)	
H2C	0.5480	0.6558	0.4273	0.021*	
H2D	0.6665	0.6929	0.4782	0.021*	
C3	0.70450 (15)	0.55615 (9)	0.43674 (6)	0.0208 (3)	
H3C	0.6773	0.5126	0.4682	0.025*	
H3D	0.6748	0.5234	0.3967	0.025*	
C4	0.85043 (15)	0.56437 (10)	0.44909 (6)	0.0220 (3)	
H4C	0.8885	0.4953	0.4484	0.026*	
H4D	0.8806	0.5941	0.4900	0.026*	
C5	0.89466 (13)	0.63175 (9)	0.40110 (6)	0.0175 (2)	
H5C	0.9895	0.6380	0.4102	0.021*	
H5D	0.8694	0.5998	0.3605	0.021*	
C6	0.83391 (12)	0.73752 (9)	0.40071 (5)	0.0126 (2)	
H6	0.8654	0.7702	0.4412	0.015*	
C7	0.66650 (12)	0.94593 (8)	0.21109 (5)	0.0120 (2)	
H7	0.6215	1.0126	0.2134	0.014*	
C8	0.61868 (13)	0.89995 (10)	0.14856 (6)	0.0169 (2)	
H8A	0.5241	0.8921	0.1417	0.020*	

H8B	0.6570	0.8313	0.1464	0.020*	
C9	0.65520 (14)	0.96868 (11)	0.09937 (6)	0.0221 (3)	
H9A	0.6104	1.0351	0.0992	0.026*	
H9B	0.6273	0.9363	0.0590	0.026*	
C10	0.80021 (14)	0.98656 (11)	0.11056 (6)	0.0208 (3)	
H10A	0.8213	1.0329	0.0789	0.025*	
H10B	0.8446	0.9207	0.1075	0.025*	
C11	0.84846 (13)	1.03355 (9)	0.17347 (6)	0.0162 (2)	
H11A	0.9432	1.0405	0.1805	0.019*	
H11B	0.8110	1.1026	0.1752	0.019*	
C12	0.81033 (11)	0.96599 (9)	0.22289 (5)	0.0123 (2)	
H12	0.8560	0.8990	0.2232	0.015*	
C13	0.65454 (12)	1.09834 (9)	0.41220 (6)	0.0133 (2)	
H13	0.6297	1.0503	0.4428	0.016*	
C14	0.58832 (14)	1.20051 (9)	0.41731 (7)	0.0199 (2)	
H14A	0.4940	1.1905	0.4100	0.024*	
H14B	0.6084	1.2482	0.3861	0.024*	
C15	0.63376 (16)	1.24644 (10)	0.48076 (7)	0.0253 (3)	
H15A	0.5931	1.3142	0.4829	0.030*	
H15B	0.6064	1.2016	0.5116	0.030*	
C16	0.77959 (17)	1.25828 (12)	0.49508 (8)	0.0303 (3)	
H16A	0.8063	1.2837	0.5372	0.036*	
H16B	0.8060	1.3094	0.4672	0.036*	
C17	0.84742 (15)	1.15638 (11)	0.48828 (7)	0.0255 (3)	
H17A	0.9413	1.1678	0.4942	0.031*	
H17B	0.8309	1.1081	0.5200	0.031*	
C18	0.79971 (13)	1.10982 (9)	0.42553 (6)	0.0164 (2)	
H18	0.8250	1.1561	0.3942	0.020*	
Zn1A	0.76131 (2)	0.36294 (2)	0.25952 (2)	0.01355 (6)	0.94
Cl1A	0.63241 (4)	0.50140 (3)	0.26301 (2)	0.01450 (9)	0.94
Cl2A	0.86823 (7)	0.38639 (5)	0.18121 (3)	0.02463 (12)	0.94
Cl3A	0.62548 (10)	0.22658 (7)	0.23764 (4)	0.01813 (15)	0.94
Cl4A	0.89500 (4)	0.34751 (3)	0.34937 (2)	0.02371 (8)	0.94
Zn1B	0.7716 (6)	0.3345 (4)	0.2463 (3)	0.0287 (10)	0.06
Cl1B	0.9326 (9)	0.2641 (8)	0.3181 (5)	0.056 (3)	0.06
Cl2B	0.6709 (11)	0.4742 (7)	0.2722 (6)	0.040 (2)	0.06
Cl3B	0.8504 (18)	0.3625 (15)	0.1645 (9)	0.062 (5)	0.06
Cl4B	0.6062 (18)	0.2271 (16)	0.2239 (8)	0.031 (4)	0.06
Cl5	0.76791 (6)	0.60816 (3)	0.03343 (2)	0.04180 (13)	
O1W	0.94431 (12)	0.64959 (9)	0.15743 (6)	0.0304 (3)	
H1O1	0.8960 (19)	0.6389 (15)	0.1230 (6)	0.037*	
H2O1	0.982 (2)	0.7057 (11)	0.1558 (10)	0.037*	
O2W	0.78942 (13)	0.69376 (8)	0.23958 (5)	0.0276 (2)	
H1O2	0.7480 (18)	0.6407 (11)	0.2436 (10)	0.033*	
H2O2	0.8389 (18)	0.6809 (15)	0.2159 (9)	0.033*	
O3W	0.40117 (10)	0.95853 (9)	0.39920 (5)	0.0259 (2)	
H1O3	0.3764 (19)	1.0020 (13)	0.4223 (8)	0.031*	
H2O3	0.3341 (14)	0.9374 (15)	0.3769 (8)	0.031*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cr1	0.00941 (9)	0.00891 (8)	0.01117 (9)	-0.00020 (6)	0.00483 (7)	0.00071 (6)
N1	0.0136 (5)	0.0139 (4)	0.0196 (5)	0.0054 (3)	0.0102 (4)	0.0066 (4)
N2	0.0111 (5)	0.0135 (4)	0.0204 (5)	0.0001 (3)	0.0094 (4)	0.0001 (4)
N3	0.0125 (5)	0.0125 (4)	0.0143 (5)	-0.0047 (3)	0.0050 (4)	0.0004 (3)
N4	0.0151 (5)	0.0154 (4)	0.0142 (5)	-0.0072 (4)	0.0057 (4)	-0.0016 (4)
N5	0.0126 (5)	0.0117 (4)	0.0145 (5)	0.0015 (3)	0.0019 (4)	0.0012 (3)
N6	0.0112 (5)	0.0196 (5)	0.0178 (5)	0.0035 (4)	0.0023 (4)	-0.0034 (4)
C1	0.0119 (5)	0.0119 (5)	0.0154 (5)	0.0019 (4)	0.0066 (4)	0.0035 (4)
C2	0.0235 (7)	0.0145 (5)	0.0191 (6)	0.0009 (4)	0.0124 (5)	0.0052 (4)
C3	0.0355 (8)	0.0122 (5)	0.0169 (6)	0.0012 (5)	0.0109 (5)	0.0025 (4)
C4	0.0328 (8)	0.0146 (5)	0.0184 (6)	0.0089 (5)	0.0042 (5)	0.0027 (4)
C5	0.0185 (6)	0.0141 (5)	0.0205 (6)	0.0070 (4)	0.0053 (5)	-0.0009 (4)
C6	0.0118 (5)	0.0117 (5)	0.0152 (5)	0.0026 (4)	0.0045 (4)	0.0004 (4)
C7	0.0127 (5)	0.0122 (4)	0.0121 (5)	-0.0024 (4)	0.0050 (4)	0.0008 (4)
C8	0.0162 (6)	0.0203 (6)	0.0143 (5)	-0.0045 (4)	0.0033 (4)	-0.0025 (4)
C9	0.0229 (7)	0.0294 (7)	0.0139 (6)	-0.0025 (5)	0.0038 (5)	0.0017 (5)
C10	0.0225 (7)	0.0270 (6)	0.0149 (6)	-0.0026 (5)	0.0089 (5)	0.0008 (5)
C11	0.0170 (6)	0.0176 (5)	0.0162 (5)	-0.0049 (4)	0.0088 (4)	0.0019 (4)
C12	0.0127 (5)	0.0126 (5)	0.0130 (5)	-0.0030 (4)	0.0058 (4)	-0.0004 (4)
C13	0.0140 (5)	0.0108 (5)	0.0156 (5)	0.0015 (4)	0.0041 (4)	-0.0001 (4)
C14	0.0227 (6)	0.0123 (5)	0.0265 (7)	0.0055 (4)	0.0096 (5)	0.0000 (4)
C15	0.0329 (8)	0.0169 (6)	0.0294 (7)	0.0023 (5)	0.0145 (6)	-0.0063 (5)
C16	0.0357 (9)	0.0223 (6)	0.0345 (8)	-0.0063 (6)	0.0104 (7)	-0.0146 (6)
C17	0.0231 (7)	0.0269 (7)	0.0256 (7)	-0.0030 (5)	0.0020 (6)	-0.0132 (5)
C18	0.0149 (6)	0.0145 (5)	0.0202 (6)	-0.0010 (4)	0.0047 (4)	-0.0048 (4)
Zn1A	0.00992 (9)	0.01085 (11)	0.02095 (13)	0.00062 (7)	0.00562 (8)	0.00178 (7)
Cl1A	0.01152 (18)	0.01180 (19)	0.02058 (16)	0.00198 (12)	0.00407 (13)	-0.00062 (13)
Cl2A	0.0129 (2)	0.0329 (3)	0.0317 (3)	0.00436 (17)	0.0133 (2)	0.0052 (2)
Cl3A	0.0171 (4)	0.01019 (19)	0.0277 (4)	-0.0013 (2)	0.0058 (3)	-0.0002 (2)
Cl4A	0.02115 (17)	0.01976 (15)	0.02746 (18)	0.00188 (11)	-0.00247 (13)	0.00251 (12)
Zn1B	0.029 (2)	0.027 (2)	0.034 (3)	-0.0094 (19)	0.0149 (16)	-0.0010 (17)
Cl1B	0.037 (4)	0.069 (6)	0.059 (6)	-0.013 (4)	-0.002 (4)	0.028 (5)
Cl2B	0.037 (5)	0.019 (4)	0.071 (7)	-0.017 (3)	0.030 (5)	-0.020 (4)
Cl3B	0.041 (9)	0.085 (12)	0.066 (11)	0.026 (7)	0.028 (8)	-0.001 (7)
Cl4B	0.010 (4)	0.034 (5)	0.045 (8)	0.010 (3)	-0.008 (4)	-0.004 (5)
Cl5	0.0815 (4)	0.03036 (19)	0.01572 (16)	-0.0234 (2)	0.01466 (19)	-0.00282 (13)
O1W	0.0211 (5)	0.0298 (5)	0.0381 (7)	-0.0013 (4)	0.0000 (5)	0.0111 (5)
O2W	0.0423 (7)	0.0146 (4)	0.0300 (6)	-0.0058 (4)	0.0168 (5)	-0.0031 (4)
O3W	0.0156 (5)	0.0347 (6)	0.0285 (5)	0.0001 (4)	0.0071 (4)	-0.0024 (4)

Geometric parameters (\AA , $^\circ$)

Cr1—N3	2.0737 (12)	C8—H8A	0.9900
Cr1—N5	2.0817 (10)	C8—H8B	0.9900
Cr1—N2	2.0839 (11)	C9—C10	1.527 (2)

Cr1—N1	2.0859 (11)	C9—H9A	0.9900
Cr1—N4	2.0899 (11)	C9—H9B	0.9900
Cr1—N6	2.0928 (12)	C10—C11	1.5330 (19)
N1—C1	1.4937 (15)	C10—H10A	0.9900
N1—H1A	0.9100	C10—H10B	0.9900
N1—H1B	0.9100	C11—C12	1.5283 (16)
N2—C6	1.4932 (15)	C11—H11A	0.9900
N2—H2A	0.9100	C11—H11B	0.9900
N2—H2B	0.9100	C12—H12	1.0000
N3—C7	1.4927 (15)	C13—C18	1.5178 (18)
N3—H3A	0.9100	C13—C14	1.5232 (16)
N3—H3B	0.9100	C13—H13	1.0000
N4—C12	1.4942 (15)	C14—C15	1.533 (2)
N4—H4A	0.9100	C14—H14A	0.9900
N4—H4B	0.9100	C14—H14B	0.9900
N5—C13	1.4971 (16)	C15—C16	1.525 (2)
N5—H5A	0.9100	C15—H15A	0.9900
N5—H5B	0.9100	C15—H15B	0.9900
N6—C18	1.5009 (16)	C16—C17	1.535 (2)
N6—H6A	0.9100	C16—H16A	0.9900
N6—H6B	0.9100	C16—H16B	0.9900
C1—C6	1.5179 (17)	C17—C18	1.5270 (19)
C1—C2	1.5289 (16)	C17—H17A	0.9900
C1—H1	1.0000	C17—H17B	0.9900
C2—C3	1.5356 (18)	C18—H18	1.0000
C2—H2C	0.9900	Zn1A—C14A	2.2400 (9)
C2—H2D	0.9900	Zn1A—C11A	2.2779 (6)
C3—C4	1.522 (2)	Zn1A—C12A	2.2799 (9)
C3—H3C	0.9900	Zn1A—C13A	2.2856 (10)
C3—H3D	0.9900	Zn1B—C13B	2.18 (2)
C4—C5	1.529 (2)	Zn1B—C14B	2.23 (2)
C4—H4C	0.9900	Zn1B—C12B	2.246 (11)
C4—H4D	0.9900	Zn1B—C11B	2.304 (11)
C5—C6	1.5247 (16)	C11B—O1W ⁱ	1.993 (11)
C5—H5C	0.9900	O1W—C11B ⁱⁱ	1.993 (11)
C5—H5D	0.9900	O1W—H1O1	0.853 (9)
C6—H6	1.0000	O1W—H2O1	0.841 (9)
C7—C12	1.5194 (17)	O2W—H1O2	0.834 (9)
C7—C8	1.5197 (17)	O2W—H2O2	0.829 (9)
C7—H7	1.0000	O3W—H1O3	0.843 (9)
C8—C9	1.5267 (19)	O3W—H2O3	0.835 (9)
N3—Cr1—N5	94.24 (5)	C12—C7—C8	112.19 (10)
N3—Cr1—N2	92.11 (5)	N3—C7—H7	108.2
N5—Cr1—N2	169.03 (4)	C12—C7—H7	108.2
N3—Cr1—N1	91.55 (5)	C8—C7—H7	108.2
N5—Cr1—N1	89.09 (4)	C7—C8—C9	110.21 (10)
N2—Cr1—N1	81.81 (4)	C7—C8—H8A	109.6

N3—Cr1—N4	82.06 (4)	C9—C8—H8A	109.6
N5—Cr1—N4	94.99 (5)	C7—C8—H8B	109.6
N2—Cr1—N4	94.74 (5)	C9—C8—H8B	109.6
N1—Cr1—N4	172.64 (4)	H8A—C8—H8B	108.1
N3—Cr1—N6	171.68 (4)	C8—C9—C10	110.83 (12)
N5—Cr1—N6	82.49 (5)	C8—C9—H9A	109.5
N2—Cr1—N6	92.35 (5)	C10—C9—H9A	109.5
N1—Cr1—N6	96.04 (5)	C8—C9—H9B	109.5
N4—Cr1—N6	90.58 (5)	C10—C9—H9B	109.5
C1—N1—Cr1	109.07 (7)	H9A—C9—H9B	108.1
C1—N1—H1A	109.9	C9—C10—C11	111.28 (11)
Cr1—N1—H1A	109.9	C9—C10—H10A	109.4
C1—N1—H1B	109.9	C11—C10—H10A	109.4
Cr1—N1—H1B	109.9	C9—C10—H10B	109.4
H1A—N1—H1B	108.3	C11—C10—H10B	109.4
C6—N2—Cr1	109.01 (7)	H10A—C10—H10B	108.0
C6—N2—H2A	109.9	C12—C11—C10	110.21 (10)
Cr1—N2—H2A	109.9	C12—C11—H11A	109.6
C6—N2—H2B	109.9	C10—C11—H11A	109.6
Cr1—N2—H2B	109.9	C12—C11—H11B	109.6
H2A—N2—H2B	108.3	C10—C11—H11B	109.6
C7—N3—Cr1	109.01 (7)	H11A—C11—H11B	108.1
C7—N3—H3A	109.9	N4—C12—C7	106.28 (10)
Cr1—N3—H3A	109.9	N4—C12—C11	113.29 (9)
C7—N3—H3B	109.9	C7—C12—C11	111.54 (10)
Cr1—N3—H3B	109.9	N4—C12—H12	108.5
H3A—N3—H3B	108.3	C7—C12—H12	108.5
C12—N4—Cr1	109.05 (7)	C11—C12—H12	108.5
C12—N4—H4A	109.9	N5—C13—C18	107.70 (10)
Cr1—N4—H4A	109.9	N5—C13—C14	112.66 (10)
C12—N4—H4B	109.9	C18—C13—C14	111.25 (10)
Cr1—N4—H4B	109.9	N5—C13—H13	108.4
H4A—N4—H4B	108.3	C18—C13—H13	108.4
C13—N5—Cr1	108.39 (7)	C14—C13—H13	108.4
C13—N5—H5A	110.0	C13—C14—C15	110.15 (11)
Cr1—N5—H5A	110.0	C13—C14—H14A	109.6
C13—N5—H5B	110.0	C15—C14—H14A	109.6
Cr1—N5—H5B	110.0	C13—C14—H14B	109.6
H5A—N5—H5B	108.4	C15—C14—H14B	109.6
C18—N6—Cr1	109.12 (8)	H14A—C14—H14B	108.1
C18—N6—H6A	109.9	C16—C15—C14	111.29 (12)
Cr1—N6—H6A	109.9	C16—C15—H15A	109.4
C18—N6—H6B	109.9	C14—C15—H15A	109.4
Cr1—N6—H6B	109.9	C16—C15—H15B	109.4
H6A—N6—H6B	108.3	C14—C15—H15B	109.4
N1—C1—C6	106.11 (9)	H15A—C15—H15B	108.0
N1—C1—C2	112.82 (10)	C15—C16—C17	111.45 (12)
C6—C1—C2	111.69 (10)	C15—C16—H16A	109.3

N1—C1—H1	108.7	C17—C16—H16A	109.3
C6—C1—H1	108.7	C15—C16—H16B	109.3
C2—C1—H1	108.7	C17—C16—H16B	109.3
C1—C2—C3	110.02 (10)	H16A—C16—H16B	108.0
C1—C2—H2C	109.7	C18—C17—C16	110.94 (13)
C3—C2—H2C	109.7	C18—C17—H17A	109.5
C1—C2—H2D	109.7	C16—C17—H17A	109.5
C3—C2—H2D	109.7	C18—C17—H17B	109.5
H2C—C2—H2D	108.2	C16—C17—H17B	109.5
C4—C3—C2	110.76 (11)	H17A—C17—H17B	108.0
C4—C3—H3C	109.5	N6—C18—C13	106.80 (10)
C2—C3—H3C	109.5	N6—C18—C17	112.82 (11)
C4—C3—H3D	109.5	C13—C18—C17	111.57 (11)
C2—C3—H3D	109.5	N6—C18—H18	108.5
H3C—C3—H3D	108.1	C13—C18—H18	108.5
C3—C4—C5	110.38 (11)	C17—C18—H18	108.5
C3—C4—H4C	109.6	Cl4A—Zn1A—Cl1A	108.81 (2)
C5—C4—H4C	109.6	Cl4A—Zn1A—Cl2A	112.38 (3)
C3—C4—H4D	109.6	Cl1A—Zn1A—Cl2A	107.91 (3)
C5—C4—H4D	109.6	Cl4A—Zn1A—Cl3A	112.84 (3)
H4C—C4—H4D	108.1	Cl1A—Zn1A—Cl3A	105.66 (4)
C6—C5—C4	109.98 (11)	Cl2A—Zn1A—Cl3A	108.92 (3)
C6—C5—H5C	109.7	Cl3B—Zn1B—Cl4B	109.0 (8)
C4—C5—H5C	109.7	Cl3B—Zn1B—Cl2B	110.8 (6)
C6—C5—H5D	109.7	Cl4B—Zn1B—Cl2B	100.3 (7)
C4—C5—H5D	109.7	Cl3B—Zn1B—Cl1B	107.6 (6)
H5C—C5—H5D	108.2	Cl4B—Zn1B—Cl1B	110.7 (6)
N2—C6—C1	106.74 (10)	Cl2B—Zn1B—Cl1B	118.1 (5)
N2—C6—C5	113.24 (10)	O1W ⁱ —Cl1B—Zn1B	148.4 (7)
C1—C6—C5	111.78 (10)	Cl1B ⁱⁱ —O1W—H1O1	128.5 (15)
N2—C6—H6	108.3	Cl1B ⁱⁱ —O1W—H2O1	20.4 (15)
C1—C6—H6	108.3	H1O1—O1W—H2O1	108.4 (17)
C5—C6—H6	108.3	H1O2—O2W—H2O2	108.0 (17)
N3—C7—C12	106.99 (10)	H1O3—O3W—H2O3	105.2 (17)
N3—C7—C8	112.78 (9)		
Cr1—N1—C1—C6	43.74 (11)	Cr1—N4—C12—C11	165.21 (8)
Cr1—N1—C1—C2	166.35 (9)	N3—C7—C12—N4	-56.47 (11)
N1—C1—C2—C3	-174.32 (11)	C8—C7—C12—N4	179.35 (9)
C6—C1—C2—C3	-54.88 (14)	N3—C7—C12—C11	179.60 (9)
C1—C2—C3—C4	56.88 (15)	C8—C7—C12—C11	55.42 (13)
C2—C3—C4—C5	-58.90 (14)	C10—C11—C12—N4	-174.49 (11)
C3—C4—C5—C6	57.83 (14)	C10—C11—C12—C7	-54.61 (14)
Cr1—N2—C6—C1	43.07 (10)	Cr1—N5—C13—C18	43.53 (10)
Cr1—N2—C6—C5	166.49 (8)	Cr1—N5—C13—C14	166.59 (9)
N1—C1—C6—N2	-57.08 (12)	N5—C13—C14—C15	-178.16 (11)
C2—C1—C6—N2	179.59 (10)	C18—C13—C14—C15	-57.11 (15)
N1—C1—C6—C5	178.59 (10)	C13—C14—C15—C16	56.49 (16)

C2—C1—C6—C5	55.27 (13)	C14—C15—C16—C17	-55.32 (18)
C4—C5—C6—N2	-176.70 (11)	C15—C16—C17—C18	54.00 (18)
C4—C5—C6—C1	-56.10 (14)	Cr1—N6—C18—C13	40.46 (11)
Cr1—N3—C7—C12	43.41 (10)	Cr1—N6—C18—C17	163.39 (10)
Cr1—N3—C7—C8	167.23 (8)	N5—C13—C18—N6	-55.59 (12)
N3—C7—C8—C9	-176.66 (11)	C14—C13—C18—N6	-179.51 (10)
C12—C7—C8—C9	-55.76 (14)	N5—C13—C18—C17	-179.30 (10)
C7—C8—C9—C10	56.43 (15)	C14—C13—C18—C17	56.79 (14)
C8—C9—C10—C11	-57.35 (15)	C16—C17—C18—N6	-174.93 (12)
C9—C10—C11—C12	55.91 (15)	C16—C17—C18—C13	-54.69 (16)
Cr1—N4—C12—C7	42.38 (10)		

Symmetry codes: (i) $-x+2, y-1/2, -z+1/2$; (ii) $-x+2, y+1/2, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots C15 ⁱⁱⁱ	0.91	2.40	3.2535 (15)	157
N1—H1B \cdots O3W	0.91	2.36	3.0178 (16)	129
N2—H2A \cdots O2W	0.91	2.01	2.9051 (17)	166
N2—H2B \cdots C12A ⁱⁱ	0.91	2.45	3.2197 (14)	142
N2—H2B \cdots C13B ⁱⁱ	0.91	2.36	3.180 (18)	150
N3—H3A \cdots O2W	0.91	2.13	2.9832 (16)	156
N3—H3B \cdots C11A ^{iv}	0.91	2.52	3.2574 (13)	138
N3—H3B \cdots C13A ^{iv}	0.91	2.77	3.4547 (16)	133
N3—H3B \cdots C12B ^{iv}	0.91	2.67	3.471 (10)	147
N3—H3B \cdots C14B ^{iv}	0.91	2.68	3.35 (2)	131
N4—H4A \cdots C11B ^v	0.91	2.74	3.473 (11)	138
N4—H4B \cdots C12A ⁱⁱ	0.91	2.64	3.4267 (15)	146
N4—H4B \cdots O1W ^{vi}	0.91	2.39	2.9804 (17)	123
N5—H5A \cdots C13A ^v	0.91	2.51	3.4245 (14)	178
N5—H5A \cdots C14B ^v	0.91	2.73	3.634 (19)	173
N5—H5B \cdots C11A ^{iv}	0.91	2.74	3.3664 (16)	127
N5—H5B \cdots O3W	0.91	2.22	2.9724 (17)	140
N6—H6A \cdots C15 ⁱⁱⁱ	0.91	2.39	3.2474 (14)	158
O1W—H1O1 \cdots C15	0.85 (1)	2.24 (1)	3.0878 (17)	179 (2)
O1W—H2O1 \cdots C14A ⁱⁱ	0.84 (1)	2.28 (1)	3.1170 (13)	174 (2)
O2W—H1O2 \cdots C11A	0.83 (1)	2.28 (1)	3.1140 (12)	175 (2)
O2W—H1O2 \cdots C12B	0.83 (1)	2.45 (1)	3.271 (9)	167 (2)
O2W—H2O2 \cdots O1W	0.83 (1)	1.92 (1)	2.7468 (19)	177 (2)
O3W—H1O3 \cdots C15 ^{iv}	0.84 (1)	2.41 (1)	3.2139 (13)	159 (2)
O3W—H2O3 \cdots C12A ^{iv}	0.84 (1)	2.38 (1)	3.2153 (17)	175 (2)
O3W—H2O3 \cdots C13B ^{iv}	0.84 (1)	2.23 (2)	3.05 (2)	167 (2)

Symmetry codes: (ii) $-x+2, y+1/2, -z+1/2$; (iii) $x, -y+3/2, z+1/2$; (iv) $-x+1, y+1/2, -z+1/2$; (v) $x, y+1, z$.