

Received 15 July 2016 Accepted 16 July 2016

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland

Keywords: crystal structure; synthesis; chalcones; molecular structure; molecular conformation; conformational disorder; hydrogen bonding; supramolecular assembly.

CCDC references: 1494027; 1494026

Supporting information: this article has supporting information at journals.iucr.org/e



OPEN d ACCESS

The crystal structures of two chalcones: (2*E*)-1-(5chlorothiophen-2-yl)-3-(2-methylphenyl)prop-2-en-1-one and (2*E*)-1-(anthracen-9-yl)-3-[4-(propan-2yl)phenyl]prop-2-en-1-one

Marisiddaiah Girisha,^a Hemmige S. Yathirajan,^a* Jerry P. Jasinski^b and Christopher Glidewell^c

^aDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysuru 570 006, India, ^bDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, and ^cSchool of Chemistry, University of St Andrews, Fife KY16 9ST, UK. *Correspondence e-mail: yathirajan@hotmail.com

In the crystal of compound (I), $C_{14}H_{11}ClOS$, molecules are linked by $C-H\cdots O$ hydrogen bonds to form simple C(5) chains. Compound (II), $C_{26}H_{22}O$, crystallizes with Z' = 2 in space group $P\overline{1}$; one of the molecules is fully ordered but the other is disordered over two sets of atomic sites having occupancies 0.644 (3) and 0.356 (3). The two disordered components differ from one another in the orientation of the isopropyl substituents, and both differ from the ordered molecules in the arrangement of the central propenone spacer unit, so that the crystal of (II) contains three distinct conformers. The ordered and disordered conformers each form a C(8) chain built from a single type of $C-H\cdots O$ hydrogen bond but those formed by the disordered conformers differ from that formed by the ordered form.

1. Chemical context

Chalcones, $R^1 - C(=O) - CH = CH - R^2$, are versatile intermediates in synthesis (Baktır et al., 2011; Samshuddin et al., 2011, 2012, 2014; Nayak et al., 2014; Salian et al., 2015; Mohan et al., 2016). Compounds of this class also exhibit a wide range of biological activity, including anti-bacterial (Tran et al., 2012), anti-cancer (Syam et al., 2012; Kumar et al., 2014), antifungal (López et al., 2001), anti-inflammatory (Fang et al., 2015), anti-malarial (Agarwal et al., 2005) and antitubercular activities (Dimmock et al., 1999). Accordingly, the synthesis and characterization of new examples of this type is of interest and potentially of value and herein we report on the synthesis and crystal structures of two further examples; (2E)-1-(5chlorothiophen-2-yl)-3-(2-methylphenyl)-prop-2-en-1-one (I), and (2E)-1-(anthracen-9-yl)-3-[4-(propan-2-yl)phenyl]prop-2en-1-one (II). Compounds (I) and (II) were prepared by baseinduced condensation of an aryl aldehyde, 2-methylbenzaldehyde in the case of (I) or 4-isopropylbenzaldehyde for (II) with, respectively, 2-acetyl-5-chlorothiophene or 9acetylanthracene.

2. Structural commentary

In the molecule of compound (I), Fig. 1, the central spacer unit comprising the atoms (C12,C1,C2,C3,C31) is effectively planar: the maximum deviation from the mean plane of these atoms is 0.21 (2) Å, with an r.m.s. deviation of 0.025 Å. The

research communications

heterocyclic ring is nearly co-planar with the spacer unit, making with it a dihedral angle of $1.41 (1)^{\circ}$. The dihedral angles between the phenyl group and the spacer unit, and between the two rings are 10.95 (11) and 9.81 (10)°, respectively. The bond distances within the molecule of (I) show clearly the localized double bond between atoms C2 and C3, and the distances within the thiophene ring clearly rule out the possibility of any orientational disorder of the type sometimes found in thiophene rings (Cobo *et al.*, 2005; Trilleras *et al.*, 2005, 2009; Insuasty *et al.*, 2014).



(II), conformers 2 and 3

Compound (II) crystallizes with Z' = 2 in space group $P\overline{1}$. The molecule containing atom O11 (Fig. 2) is fully ordered, but the other molecule is disordered over two sets of atomic sites: the major-disorder component containing atom O21 (Fig. 3) has occupancy 0.644 (3) while the minor-disorder component containing atom O31 (Fig. 4) has occupancy 0.356 (3). All three forms exhibit different conformations, as discussed below, and it will be convenient to refer to the molecules containing atoms O11, O21 or O31 as conformers of types 1, 2 or 3, respectively.





The molecular structure of compound (I), with atom labelling and displacement ellipsoids drawn at the 30% probability level.

In the fully ordered molecule containing atom O11 the torsional angle C119-C11-C12-C13 is 177.72 (16)° whereas in the two disordered components containing atoms O21 and O31 the values of the corresponding torsional angles Cn19-Cn1-Cn2-Cn3 are 11 (3)° and 12 (5)° for n = 2 and 3 respectively, corresponding to a rotation of approximately 180° about the bond Cn1-Cn2 in conformers 2 and 3 as compared with conformer 1. In addition, in conformers 2 and 3 the torsional angles Cn33-Cn34-Cn37-Hn37 are -14° and -170° for n = 2 and 3, respectively, so that the orientation of the CHMe₂ group in these two forms differs by a rotation of approximately 180° about the bond Cn34-Cn37; the corresponding value in conformer 1 is *ca* 162°. Hence three different conformations of compound (II) co-exist in the same





The molecular structure of conformer 1 in compound (II), with atom labelling and displacement ellipsoids drawn at the 30% probability level.



Figure 3

The molecular structure of the major-disorder component, conformer 2 having occupancy 0.644 (3), in compound (II), with atom labelling and displacement ellipsoids drawn at the 30% probability level.

crystal (*cf.* Figs. 2–4) with relative abundances 1.000:0.644 (3):0.356 (3) in the crystal selected for data collection. Conformer 1 thus differs from conformers 2 and 3 in the arrangement of the central spacer unit, while conformers 1 and 3 exhibit similar orientations of the isopropyl unit relative to the adjacent phenyl ring, but different from that in conformer 2.



Figure 4

The molecular structure of the minor-disorder component, conformer 3 having occupancy 0.356 (3), in compound (II), with atom labelling and displacement ellipsoids drawn at the 30% probability level.

Table 1	
---------	--

-x + 1, -y + 1, -z + 1.

Hydrogen-bond geometry (Å, °) for (I).

$D - \mathbf{H} \cdot \cdot \cdot A$	$D-\mathrm{H}$	$H \cdot \cdot \cdot A$	$D \cdots A$	$D - \mathbf{H} \cdot \cdot \cdot A$
$C13-H13\cdots O1^{i}$	0.93	2.55	3.467 (2)	169

Symmetry code: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Table 2

Hydrogen-bond geometry (Å, °) for (II).

Cg1 and Cg2 are the centroids of rings (C111–C114/C122/C111) and (C114–C118/C124/C113), respectively.

$D - H \cdots A$	D-H	$H \cdot \cdot \cdot A$	$D \cdots A$	$D - \mathbf{H} \cdot \cdot \cdot A$
C133-H133···O11 ⁱ	0.93	2.49	3.336 (2)	151
$C216-H216\cdots O21^{i}$	0.93	2.61	3.41 (2)	144
$C216-H216\cdots O31^{i}$	0.93	2.58	3.36 (4)	142
$C316-H316\cdots O21^{i}$	0.93	2.51	3.30 (3)	144
$C316-H316\cdots O31^{i}$	0.93	2.47	3.25 (4)	142
$C233-H233\cdots Cg1^{ii}$	0.93	2.64	3.355 (4)	134
$C236-H236\cdots Cg2^{iii}$	0.93	2.75	3.519 (4)	140
$C336-H336\cdots Cg2^{iii}$	0.93	2.80	3.354 (7)	119
Symmetry codes: (i) $x - 1, y$,	z; (ii)	-x + 2, -y + 1,	-z + 1; (iii)

In each conformer, the central spacer unit encompassing

atoms Cn19, Cn1, Cn2, Cn3, Cn31 is effectively planar with r.m.s. deviations from the mean planes of 0.011, 0.036 and 0.043 Å for n = 1-3, respectively. This spacer unit makes dihedral angles with the central ring of the anthracene unit of 67.70 (11), 65.7 (5) and 71.7 (10)° for n = 1-3, respectively, and the corresponding dihedral angles with the adjacent aryl rings Cn31-Cn36 are 6.26 (18), 1.5 (11) and 7(2)°, respectively. These values confirm that the principal difference between conformer 1 and conformers 2 and 3 is simply a rotation about the bond Cn1-Cn2.

Within each of the anthracene units, the distances Cn11 - Cn12, Cn13 - Cn14, Cn15 - Cn16 and Cn17 - Cn18 are very much shorter than the other C-C bonds in these units, while the C-C distances in the central rings show rather little variation. These observations are consistent with an electronic structure for the anthracene units where a central ring displaying aromatic delocalization is flanked by two isolated diene units (Glidewell & Lloyd, 1984,1986).

3. Supramolecular features

In the crystal of compound (I), molecule related by a *c*-glide plane are linked by a single $C-H\cdots O$ hydrogen bond (Table 1) to form a C(5) chain running parallel to the [001] direction (Fig. 5). In the crystal of compound (II), molecules are also linked into chains by $C-H\cdots O$ hydrogen bonds (Table 2), but the chains formed by the ordered and disordered forms are different, in that in the chain of ordered molecules the donor is a phenyl C-H unit, while in the disordered forms the donors are part of the anthracene units. In both types of chain molecules related by translation form C(8) chains running parallel to the [100] direction (Fig. 6). In

research communications



Figure 5

Part of the crystal structure of compound (I), showing the formation of a hydrogen-bonded C(5) chain parallel to [001]. Hydrogen bonds are shown as dashed lines and, for the sake of clarity, the H atoms not involved in the motif shown are omitted. The atoms marked with an asterisk (*) or a hash (#) are at the symmetry positions $(x, \frac{1}{2} - y, -\frac{1}{2} + z)$ and $(x, \frac{1}{2} - y, \frac{1}{2} + z)$, respectively.

addition, inversion-related pairs of the chains built from the disordered components are weakly linked by $C-H\cdots\pi$ interactions (Table 2).

4. Database survey

The structures of a number of chalcones containing substituted thiophene units, and thus closely related to compound (I) have been reported recently (Naik, Shettigar et al., 2015; Naik, Yathirajan et al., 2015). There are no hydrogen bonds of any kind in the crystals of the isostructural compounds (2E)-1-(5-chlorothiophen-2-yl)-3-(4-ethylphenyl)prop-2-en-1-one and (2E)-1-(5-bromothiophen-2-yl)-3-(4-ethylphenyl)prop-2en-1-one, but in the isostructural compounds (2E)-1-(5chlorothiophen-2-yl)-3-(4-ethoxyphenyl)prop-2-en-1-one and (2*E*)-1-(5-bromothiophen-2-yl)-3-(4-ethoxyphenyl)prop-2en-1-one the molecules are linked by C-H···O hydrogen bonds to form simple C(7) chains, while C(5) chains are present in the structure of (2E)-1-(5-bromothiophen-2-yl)-3-(3-methoxyphenyl)prop-2-en-1-one. In the structure of (2E, 2'E)-3, 3'-(1,3-phenylene)-bis(1-(anthracene-9-yl)prop-2-en-1-one), which is related to compound (II), inversion-related pairs of molecules are linked by multiple $C-H \cdots O$ hydrogen bonds to form centrosymmetric dimers (Kant et al., 2015). In the recently reported structure of (2E)-3-(2,4-dichlorophenyl)-1-(2-methoxyphenyl)prop-2-en-1-one (Salian et al., 2016),



Figure 6

Part of the crystal structure of compound (II), showing the formation of two different types of hydrogen-bonded C(8) chain. Hydrogen bonds are shown as dashed lines and, for the sake of clarity, the minor-disorder component and the H atoms not involved in the motifs shown have been omitted. The atoms marked with an asterisk (*) or a hash (#) are at the symmetry positions (-1 + x, y, z) and (1 + x, y, z), respectively.

there are no hydrogen bonds of any kind, but the molecules are linked into chains by π - π stacking interactions.

5. Synthesis and crystallization

For the synthesis of compound (I), a solution of 2-methylbenzaldehyde (0.075 g, 0.625 mol) in methanol (20 ml) was added to solution of with 2-acetyl-5-chlorothiophene (0.100 g, 0.625 mol) in methanol (10 ml) and to this mixture was added aqueous sodium hydroxide solution (40% w/v, 5 ml). The reaction mixture was then stirred at 301 K for 4 h, when the resulting solid product was collected by filtration, washed with cold water and dried. Crystals suitable for single-crystal X-ray diffraction were grown by slow evaporation, at ambient temperature, of a solution in acetone-dimethylformamide (1:1, v/v): m. p. 387–389 K. For the synthesis of compound (II), aqueous sodium hydroxide solution (10%, w/v, 15 ml) was added to a mixture of 4-isopropylbenzaldehyde (1.5 ml,

Table 3Experimental details.

	(I)	(II)
Crystal data		
Chemical formula	C ₁₄ H ₁₁ ClOS	$C_{26}H_{22}O$
M_r	262.74	350.44
Crystal system, space group	Monoclinic, $P2_1/c$	Triclinic, $P\overline{1}$
Temperature (K)	298	296
a, b, c (Å)	14.7179 (7), 7.5783 (4), 11.5451 (5)	9.0150 (4), 14.0601 (6), 16.2611 (8)
$\alpha, \beta, \gamma(\circ)$	90, 102.999 (4), 90	105.146 (4), 95.967 (4), 98.057 (4)
$V(\text{\AA}^3)$	1254.70 (11)	1948.46 (16)
Ζ	4	4
Radiation type	Μο Κα	Cu Ka
$\mu (\text{mm}^{-1})$	0.45	0.55
Crystal size (mm)	$0.26 \times 0.22 \times 0.15$	$0.21 \times 0.14 \times 0.10$
Data collection		
Diffractometer	Agilent Xcalibur Eos Gemini	Agilent Xcalibur Eos Gemini
Absorption correction	Multi-scan (CrysAlis PRO; Agilent, 2014)	Multi-scan (CrysAlis PRO; Agilent, 2014)
T_{\min}, T_{\max}	0.832, 0.935	0.737, 0.947
No. of measured, independent and observed $[I > 2\sigma(I)]$ reflections	6097, 2785, 2053	12946, 7079, 5485
R _{int}	0.028	0.035
$(\sin \theta / \lambda)_{\max} (\text{\AA}^{-1})$	0.651	0.601
Refinement		
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.040, 0.094, 1.04	0.050, 0.143, 1.04
No. of reflections	2785	7079
No. of parameters	156	575
No. of restraints	0	72
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained
$\Delta \rho_{\rm max}, \Delta \rho_{\rm min} \ ({ m e} \ { m \AA}^{-3})$	0.18, -0.19	0.22, -0.26

Computer programs: CrysAlis PRO (Agilent, 2014), SHELXS97 (Sheldrick, 2008), SHELXL2014 (Sheldrick, 2015) and PLATON (Spek, 2009).

0.01 mol) and 9-acetylanthracene (2.2 g, 0.01 mol) in ethanol (50 ml), and the resulting mixture was stirred at 278 K for 3 h. The resulting solid product was collected by filtration and recrystallized from ethanol solution: m.p. 369–371 K. Crystals suitable for single-crystal X-ray diffraction were grown by slow evaporation, at ambient temperature, of a solution in dimethylformamide.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. It was obvious from an early stage in the refinement of compound (II) that one of the two independent molecules was disordered over two sets of atomic sites having unequal occupancies. For the minor-disorder component the bonded distances and the one-angle nonbonded distances were restrained to be the same as the corresponding distances in the major component, subject to s.u.s of 0.005 and 0.01 Å respectively. In addition, the anisotropic displacement parameters for corresponding atomic pairs of atomic sites in the two disorder components were constrained to be identical and, subject to these conditions the occupancies for the two components refined to values of 0.645 (4) and 0.355 (4). The H atoms in all but the minordisorder component of compound (II) were located in difference maps and then treated as riding atoms in geometrically idealized positions with C-H distances 0.93 Å

(alkenyl, aromatic and heteroaromatic), 0.96 Å (methyl) or 0.98 Å (aliphatic C–H) and with $U_{iso}(H) = kU_{eq}(C)$ where k = 1.5 for the methyl groups, which were permitted to rotate but not to tilt, and 1.2 for all other H atoms. The H atoms in the minor-disorder component were included in the refinement in calculated positions under exactly the same conditions. For compound (II), 16 bad outliers of low intensity were omitted from the final refinements. In the final analysis of variance for compound (I) there was a fairly large value, 2.583, of K = mean $(F_o^2)/\text{mean}$ (F_c^2) for the group of 287 very weak reflections having $F_c/F_c(\text{max})$ in the range 0.000 < $F_c/F_c(\text{max})$
in the group of 733 very weak reflections having $F_c/F_c(\text{max})$ in the range 0.000.

Acknowledgements

MG thanks the UGC for the award of a Rajeev Gandhi fellowship and HSY thanks the University of Mysore for research facilities. JPJ acknowledges the NSF–MRI program (grant No. 1039027) for funds to purchase the X-ray diffract-ometer.

References

Agarwal, A., Srivastava, K., Puri, S. K. & Chauhan, P. M. S. (2005). Bioorg. Med. Chem. 13, 4645–4650.

Agilent (2014). CrysAlis PRO. Agilent Technologies Ltd, Yarnton, England.

research communications

- Baktır, Z., Akkurt, M., Samshuddin, S., Narayana, B. & Yathirajan, H. S. (2011). Acta Cryst. E67, 01292–01293.
- Cobo, D., Quiroga, J., Cobo, J., Low, J. N. & Glidewell, C. (2005). Acta Cryst. E61, 03639–03641.
- Dimmock, J. R., Elias, D. W., Beazely, M. A. & Kandepu, N. M. (1999). Curr. Med. Chem. 6, 1125–1149.
- Fang, Q., Zhao, L., Wang, Y., Zhang, Y., Li, Z., Pan, Y., Kanchana, K., Wang, J., Tong, C., Li, D. & Liang, G. (2015). *Toxicol. Appl. Pharmacol.* 282, 129–138.
- Glidewell, C. & Lloyd, D. (1984). Tetrahedron, 40, 4455-4472.
- Glidewell, C. & Lloyd, D. (1986). J. Chem. Educ. 63, 306-309.
- Insuasty, H., Insuasty, B., Castro, E., Cobo, J. & Glidewell, C. (2014). Acta Cryst. C70, 908–911.
- Kant, R., Sharma, R., Salian, V. V., Sarojini, B. K. & Narayana, B. (2015). Eur. Chem. Bull. 4, 327–330.
- Kumar, D., Maruthi Kumar, N., Tantak, M. P., Ogura, M., Kusaka, E. & Ito, T. (2014). *Bioorg. Med. Chem. Lett.* 24, 5170–5174.
- López, S. N., Castelli, M. V., Zacchino, S. A., Domínguez, J. N., Lobo, G., Charris-Charris, J., Cortés, J. C. G., Ribas, J. C., Devia, C., Rodríguez, A. M. & Enriz, R. D. (2001). *Bioorg. Med. Chem.* 9, 1999–2013.
- Mohan, B. J., Sarojini, B. K., Yathirajan, H. S., Rathore, R. & Glidewell, C. (2016). Acta Cryst. E72, 31–34.
- Naik, V. S., Shettigar, V., Berglin, T. S., Coburn, J. S., Jasinski, J. P. & Yathirajan, H. S. (2015). *Acta Cryst.* E**71**, 965–971.
- Naik, V. S., Yathirajan, H. S., Jasinski, J. P., Smolenski, V. A. & Glidewell, C. (2015). Acta Cryst. E71, 1093–1099.

- Nayak, P. S., Narayana, B., Yathirajan, H. S., Hosten, E. C., Betz, R. & Glidewell, C. (2014). Acta Cryst. C70, 1011–1016.
- Salian, V. V., Narayana, B., Yathirajan, H. S., Akkurt, M., Çelik, Ö., Ersanlı, C. C. & Glidewell, C. (2015). Acta Cryst. C71, 610–617.
- Salian, V. V., Narayana, B., Yathirajan, H. S., Rathore, R. S., Jasinski, J. P. & Glidewell, C. (2016). Private communication (refcode: CCDC 1481314). CCDC, Cambridge, England.
- Samshuddin, S., Jasinski, J. P., Butcher, R. J., Neuhardt, E. A., Narayana, B., Yathirajan, H. S. & Glidewell, C. (2014). Acta Cryst. C70, 953–959.
- Samshuddin, S., Narayana, B., Sarojini, B. K., Yathirajan, H. S. & Raghavendra, R. (2012). *Pharma Chemica*, **4**, 1445–1457.
- Samshuddin, S., Narayana, B., Shetty, D. N. & Raghavendra, R. (2011). *Pharma Chemica*, **3**, 232–240.
- Sheldrick, G. M. (2008). Acta Cryst. A64, 112–122.
- Sheldrick, G. M. (2005). Acta Cryst. **C71**, 3–8.
- Spek, A. L. (2009). Acta Cryst. D65, 148-155.
- Syam, S., Abdelwahab, S. I., Al-Mamary, M. A. & Mohan, S. (2012). *Molecules*, **17**, 6179–6195.
- Tran, T.-D., Nguyen, T.-T.-N., Do, T.-H., Huynh, T.-N.-P., Tran, C.-D. & Thai, K.-M. (2012). *Molecules*, **17**, 6684–6696.
- Trilleras, J., Quiroga, J., Cobo, J. & Glidewell, C. (2009). *Acta Cryst.* C65, o284–o286.
- Trilleras, J., Quiroga, J., Cobo, J., Low, J. N. & Glidewell, C. (2005). Acta Cryst. E61, o1892–o1894.

Acta Cryst. (2016). E72, 1153-1158 [https://doi.org/10.1107/S2056989016011592]

The crystal structures of two chalcones: (2*E*)-1-(5-chlorothiophen-2-yl)-3-(2methylphenyl)prop-2-en-1-one and (2*E*)-1-(anthracen-9-yl)-3-[4-(propan-2yl)phenyl]prop-2-en-1-one

Marisiddaiah Girisha, Hemmige S. Yathirajan, Jerry P. Jasinski and Christopher Glidewell

Computing details

For both compounds, data collection: *CrysAlis PRO* (Agilent, 2014); cell refinement: *CrysAlis PRO* (Agilent, 2014); data reduction: *CrysAlis PRO* (Agilent, 2014); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL2014* (Sheldrick, 2015) and *PLATON* (Spek, 2009).

(I) (2E)-1-(5-Chlorothiophen-2-yl)-3-(2-methylphenyl)prop-2-en-1-one

Crystal data

C₁₄H₁₁ClOS $M_r = 262.74$ Monoclinic, $P2_1/c$ a = 14.7179 (7) Å b = 7.5783 (4) Å c = 11.5451 (5) Å $\beta = 102.999$ (4)° V = 1254.70 (11) Å³ Z = 4

Data collection

Agilent Xcalibur Eos Gemini diffractometer Detector resolution: 16.0416 pixels mm⁻¹ φ and ω scans Absorption correction: multi-scan (CrysAlis PRO; Agilent, 2014) $T_{\min} = 0.832$, $T_{\max} = 0.935$ 6097 measured reflections

Refinement

Refinement on F^2 Least-squares matrix: full $R[F^2 > 2\sigma(F^2)] = 0.040$ $wR(F^2) = 0.094$ S = 1.042785 reflections 156 parameters 0 restraints F(000) = 544 $D_x = 1.391 \text{ Mg m}^{-3}$ Mo K\alpha radiation, $\lambda = 0.71073 \text{ Å}$ Cell parameters from 2913 reflections $\theta = 3.0-29.4^{\circ}$ $\mu = 0.45 \text{ mm}^{-1}$ T = 298 KBlock, colourless $0.26 \times 0.22 \times 0.15 \text{ mm}$

2785 independent reflections 2053 reflections with $I > 2\sigma(I)$ $R_{int} = 0.028$ $\theta_{max} = 27.6^{\circ}, \ \theta_{min} = 3.0^{\circ}$ $h = -17 \rightarrow 18$ $k = -5 \rightarrow 9$ $l = -14 \rightarrow 14$

Hydrogen site location: inferred from neighbouring sites H-atom parameters constrained $w = 1/[\sigma^2(F_o^2) + (0.0304P)^2 + 0.3588P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{max} = 0.001$ $\Delta\rho_{max} = 0.18 \text{ e} \text{ Å}^{-3}$ $\Delta\rho_{min} = -0.19 \text{ e} \text{ Å}^{-3}$ Extinction correction: SHELXL2014 (Sheldrick, 2015), $Fc^*=kFc[1+0.001xFc^2\lambda^3/sin(2\theta)]^{-1/4}$ Extinction coefficient: 0.0028 (8)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

	x	у	Ζ	$U_{ m iso}$ */ $U_{ m eq}$
C1	0.43382 (14)	0.2903 (3)	0.56191 (17)	0.0443 (5)
01	0.41785 (10)	0.3013 (2)	0.66164 (12)	0.0649 (5)
C2	0.36322 (13)	0.3328 (3)	0.45350 (17)	0.0451 (5)
H2	0.3771	0.3155	0.3796	0.054*
C3	0.28068 (14)	0.3950 (3)	0.45918 (17)	0.0458 (5)
H3	0.2706	0.4129	0.5350	0.055*
S11	0.60913 (4)	0.18403 (8)	0.67339 (4)	0.04994 (18)
C12	0.52502 (13)	0.2307 (3)	0.54715 (16)	0.0398 (5)
C13	0.55665 (13)	0.1979 (3)	0.44730 (16)	0.0441 (5)
H13	0.5207	0.2148	0.3709	0.053*
C14	0.64871 (14)	0.1362 (3)	0.47103 (17)	0.0485 (5)
H14	0.6808	0.1076	0.4128	0.058*
C15	0.68504 (13)	0.1233 (3)	0.58937 (17)	0.0434 (5)
Cl15	0.79457 (4)	0.05248 (8)	0.65700 (5)	0.0609 (2)
C31	0.20305 (13)	0.4390 (3)	0.35977 (16)	0.0418 (5)
C32	0.12392 (14)	0.5248 (3)	0.38106 (19)	0.0474 (5)
C33	0.05096 (15)	0.5622 (3)	0.2844 (2)	0.0569 (6)
H33	-0.0020	0.6185	0.2974	0.068*
C34	0.05518 (16)	0.5185 (3)	0.1710 (2)	0.0623 (7)
H34	0.0054	0.5449	0.1081	0.075*
C35	0.13302 (15)	0.4352 (3)	0.14948 (19)	0.0588 (6)
H35	0.1362	0.4061	0.0722	0.071*
C36	0.20586 (14)	0.3957 (3)	0.24326 (17)	0.0499 (5)
H36	0.2582	0.3388	0.2287	0.060*
C37	0.11721 (16)	0.5819 (4)	0.5038 (2)	0.0667 (7)
H37A	0.1232	0.4808	0.5550	0.100*
H37B	0.0579	0.6372	0.4999	0.100*
H37C	0.1662	0.6643	0.5347	0.100*

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters $(Å^2)$

Atomic displacement parameters $(Å^2)$

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0463 (11)	0.0518 (13)	0.0340 (10)	-0.0035 (10)	0.0071 (8)	-0.0033 (9)
01	0.0549 (9)	0.1060 (14)	0.0336 (8)	0.0117 (9)	0.0097 (6)	-0.0032 (8)
C2	0.0449 (11)	0.0563 (13)	0.0330 (10)	-0.0021 (10)	0.0063 (8)	-0.0011 (9)

C3	0.0497 (12)	0.0521 (13)	0.0365 (10)	0.0002 (10)	0.0113 (8)	-0.0001 (9)
S11	0.0493 (3)	0.0704 (4)	0.0272 (2)	0.0019 (3)	0.0024 (2)	-0.0003 (2)
C12	0.0404 (10)	0.0463 (12)	0.0304 (9)	-0.0043 (9)	0.0031 (7)	0.0003 (8)
C13	0.0439 (11)	0.0570 (13)	0.0293 (9)	-0.0010 (10)	0.0039 (8)	0.0033 (9)
C14	0.0465 (11)	0.0648 (14)	0.0350 (10)	0.0018 (11)	0.0107 (8)	0.0006 (10)
C15	0.0406 (10)	0.0469 (12)	0.0405 (11)	-0.0031 (10)	0.0042 (8)	0.0019 (9)
Cl15	0.0479 (3)	0.0704 (4)	0.0579 (4)	0.0077 (3)	-0.0016 (2)	0.0077 (3)
C31	0.0420 (11)	0.0417 (11)	0.0407 (11)	-0.0055 (9)	0.0074 (8)	0.0037 (9)
C32	0.0450 (11)	0.0456 (12)	0.0523 (12)	-0.0067 (10)	0.0124 (9)	0.0000 (10)
C33	0.0407 (11)	0.0606 (15)	0.0679 (15)	0.0019 (11)	0.0090 (10)	0.0045 (12)
C34	0.0505 (13)	0.0740 (17)	0.0557 (14)	-0.0022 (13)	-0.0021 (10)	0.0132 (13)
C35	0.0564 (13)	0.0760 (17)	0.0413 (12)	-0.0036 (13)	0.0048 (9)	0.0076 (11)
C36	0.0469 (11)	0.0598 (14)	0.0425 (11)	0.0013 (11)	0.0090 (9)	0.0039 (10)
C37	0.0559 (14)	0.0811 (19)	0.0661 (16)	0.0043 (13)	0.0199 (11)	-0.0140 (14)

Geometric parameters (Å, °)

C1—01	1.229 (2)	C31—C36	1.394 (3)	
C1-C12	1.462 (3)	C31—C32	1.403 (3)	
C1—C2	1.471 (3)	C32—C33	1.393 (3)	
С2—С3	1.318 (3)	C32—C37	1.506 (3)	
С2—Н2	0.9300	C33—C34	1.365 (3)	
C3—C31	1.464 (3)	С33—Н33	0.9300	
С3—Н3	0.9300	C34—C35	1.379 (3)	
S11—C15	1.700 (2)	С34—Н34	0.9300	
S11—C12	1.7234 (18)	C35—C36	1.375 (3)	
C12—C13	1.360 (3)	С35—Н35	0.9300	
C13—C14	1.401 (3)	С36—Н36	0.9300	
С13—Н13	0.9300	С37—Н37А	0.9600	
C14—C15	1.354 (3)	С37—Н37В	0.9600	
C14—H14	0.9300	С37—Н37С	0.9600	
C15—C115	1.7118 (19)			
O1—C1—C12	120.29 (17)	C36—C31—C3	121.03 (18)	
O1—C1—C2	122.34 (18)	C32—C31—C3	120.03 (18)	
C12—C1—C2	117.37 (17)	C33—C32—C31	118.3 (2)	
C3—C2—C1	121.22 (18)	C33—C32—C37	119.6 (2)	
С3—С2—Н2	119.4	C31—C32—C37	122.06 (18)	
C1—C2—H2	119.4	C34—C33—C32	121.7 (2)	
C2—C3—C31	127.43 (18)	С34—С33—Н33	119.1	
С2—С3—Н3	116.3	С32—С33—Н33	119.1	
С31—С3—Н3	116.3	C33—C34—C35	120.2 (2)	
C15—S11—C12	90.70 (9)	С33—С34—Н34	119.9	
C13—C12—C1	130.82 (17)	С35—С34—Н34	119.9	
C13—C12—S11	111.20 (14)	C36—C35—C34	119.4 (2)	
C1-C12-S11	117.94 (14)	С36—С35—Н35	120.3	
C12—C13—C14	113.31 (17)	С34—С35—Н35	120.3	
C12—C13—H13	123.3	C35—C36—C31	121.4 (2)	

C14—C13—H13	123.3	С35—С36—Н36	119.3
C15—C14—C13	111.50 (18)	C31—C36—H36	119.3
C15—C14—H14	124.3	С32—С37—Н37А	109.5
C13—C14—H14	124.3	С32—С37—Н37В	109.5
C14—C15—S11	113.28 (15)	H37A—C37—H37B	109.5
C14—C15—Cl15	126.91 (17)	С32—С37—Н37С	109.5
S11—C15—Cl15	119.81 (11)	H37A—C37—H37C	109.5
C36—C31—C32	118.95 (18)	H37B—C37—H37C	109.5
O1—C1—C2—C3	4.0 (3)	C12—S11—C15—C115	-179.75 (14)
C12—C1—C2—C3	-176.9 (2)	C2—C3—C31—C36	8.6 (3)
C1—C2—C3—C31	-178.39 (19)	C2—C3—C31—C32	-171.9 (2)
O1—C1—C12—C13	175.4 (2)	C36—C31—C32—C33	0.4 (3)
C2-C1-C12-C13	-3.7 (3)	C3—C31—C32—C33	-179.11 (19)
O1-C1-C12-S11	-2.0 (3)	C36—C31—C32—C37	-177.6 (2)
C2-C1-C12-S11	178.92 (15)	C3—C31—C32—C37	2.8 (3)
C15—S11—C12—C13	0.45 (17)	C31—C32—C33—C34	-0.4 (3)
C15—S11—C12—C1	178.33 (17)	C37—C32—C33—C34	177.7 (2)
C1-C12-C13-C14	-177.9 (2)	C32—C33—C34—C35	-0.1 (4)
S11—C12—C13—C14	-0.4 (2)	C33—C34—C35—C36	0.5 (4)
C12—C13—C14—C15	0.1 (3)	C34—C35—C36—C31	-0.4 (4)
C13—C14—C15—S11	0.3 (3)	C32—C31—C36—C35	0.0 (3)
C13—C14—C15—Cl15	179.56 (17)	C3—C31—C36—C35	179.5 (2)
C12—S11—C15—C14	-0.43 (18)		

Hydrogen-bond geometry (Å, °)

D—H···A	<i>D</i> —Н	H···A	D····A	<i>D</i> —H··· <i>A</i>
C13—H13…O1 ⁱ	0.93	2.55	3.467 (2)	169

Symmetry code: (i) x, -y+1/2, z-1/2.

(II) (2*E*)-1-(Anthracen-9-yl)-3-[4-(propan-2-yl)phenyl]prop-2-en-1-one

Crystal data	
$C_{26}H_{22}O$ $M_r = 350.44$ Triclinic, $P\overline{1}$ $a = 9.0150 (4) \text{ Å}$ $b = 14.0601 (6) \text{ Å}$ $c = 16.2611 (8) \text{ Å}$ $a = 105.146 (4)^{\circ}$ $\beta = 95.967 (4)^{\circ}$	Z = 4 F(000) = 744 $D_x = 1.195 \text{ Mg m}^{-3}$ Cu K α radiation, $\lambda = 1.54184 \text{ Å}$ Cell parameters from 7095 reflections $\theta = 3.5-68.0^{\circ}$ $\mu = 0.55 \text{ mm}^{-1}$ T = 296 K
$\gamma = 98.057 (4)^{\circ}$ $V = 1948.46 (16) Å^{3}$ Data collection	Block, colourless $0.21 \times 0.14 \times 0.10 \text{ mm}$
Agilent Xcalibur Eos Gemini diffractometer Detector resolution: 16.0416 pixels mm ⁻¹ φ and ω scans	Absorption correction: multi-scan (CrysAlis PRO; Agilent, 2014) $T_{min} = 0.737, T_{max} = 0.947$ 12946 measured reflections

7079 independent reflections	$h = -10 \rightarrow 10$
5485 reflections with $I > 2\sigma(I)$	$k = -16 \rightarrow 15$
$R_{\rm int} = 0.035$	$l = -19 \rightarrow 17$
$\theta_{\max} = 68.0^{\circ}, \ \theta_{\min} = 3.3^{\circ}$	
Refinement	
Refinement on F^2	Hydrogen site location: inferred from
Least-squares matrix: full	neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.050$	H-atom parameters constrained
$wR(F^2) = 0.143$	$w = 1/[\sigma^2(F_o^2) + (0.0689P)^2 + 0.2921P]$
S = 1.04	where $P = (F_o^2 + 2F_c^2)/3$
7079 reflections	$(\Delta/\sigma)_{ m max} < 0.001$
575 parameters	$\Delta ho_{ m max} = 0.22 \ { m e} \ { m \AA}^{-3}$
72 restraints	$\Delta \rho_{\rm min} = -0.26 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters $(Å^2)$

	x	У	Ζ	$U_{ m iso}$ */ $U_{ m eq}$	Occ. (<1)
C11	0.71500 (19)	0.10252 (14)	0.34353 (11)	0.0392 (4)	
O11	0.80294 (15)	0.04914 (12)	0.31426 (9)	0.0570 (4)	
C12	0.55261 (18)	0.08287 (13)	0.30901 (11)	0.0390 (4)	
H12	0.4899	0.1246	0.3359	0.047*	
C13	0.49356 (18)	0.00672 (13)	0.24008 (10)	0.0366 (4)	
H13	0.5608	-0.0324	0.2149	0.044*	
C111	0.8247 (3)	0.08400 (15)	0.51371 (13)	0.0549 (5)	
H111	0.7926	0.0267	0.4678	0.066*	
C112	0.8752 (3)	0.07482 (18)	0.59246 (15)	0.0702 (7)	
H112	0.8780	0.0116	0.5997	0.084*	
C113	0.9235 (3)	0.16040 (19)	0.66349 (14)	0.0706 (7)	
H113	0.9579	0.1533	0.7171	0.085*	
C114	0.9198 (2)	0.25229 (17)	0.65374 (12)	0.0554 (5)	
H114	0.9512	0.3079	0.7012	0.066*	
C115	0.8165 (2)	0.47150 (15)	0.47098 (14)	0.0509 (5)	
H115	0.8472	0.5275	0.5182	0.061*	
C116	0.7706 (2)	0.48406 (17)	0.39327 (16)	0.0591 (5)	
H116	0.7697	0.5481	0.3876	0.071*	
C117	0.7240 (2)	0.39997 (17)	0.32061 (14)	0.0534 (5)	
H117	0.6926	0.4091	0.2673	0.064*	
C118	0.72458 (19)	0.30582 (15)	0.32776 (12)	0.0434 (4)	
H118	0.6952	0.2515	0.2790	0.052*	
C119	0.76971 (17)	0.19321 (13)	0.41974 (10)	0.0350 (3)	
C120	0.86656 (18)	0.36079 (13)	0.56169 (11)	0.0413 (4)	
H120	0.8978	0.4166	0.6091	0.050*	
C121	0.81986 (19)	0.17963 (13)	0.49995 (11)	0.0397 (4)	

C122	0.86911 (19)	0.26655 (14)	0.57270 (11)	0.0406 (4)	
C123	0.81887 (17)	0.37492 (13)	0.48214 (11)	0.0388 (4)	
C124	0.76959 (16)	0.28911 (13)	0.40878 (10)	0.0354 (4)	
C131	0.33555 (18)	-0.02239 (12)	0.19949 (10)	0.0334 (3)	
C132	0.22085 (19)	0.02522 (13)	0.23310 (11)	0.0388 (4)	
H132	0.2444	0.0772	0.2839	0.047*	
C133	0.07321 (19)	-0.00304(14)	0.19272 (11)	0.0415 (4)	
H133	-0.0010	0.0301	0.2168	0.050*	
C134	0.03292(18)	-0.08023(12)	0 11669 (10)	0.0346(3)	
C135	0.14636(19)	-0.12848(13)	0.08311 (10)	0.0379(4)	
H135	0.1222	-0.1806	0.0325	0.045*	
C136	0.1222	-0.10051(13)	0.12369 (10)	0.049 0.0384 (4)	
U136	0.25520 (15)	-0.1343	0.12307 (10)	0.046*	
C137	-0.12065(10)	-0.11137(14)	0.1000	0.0418(4)	
U137	0.12905 (19)	0.11137 (14)	0.0/19/(11)	0.0418 (4)	
П137 С129	-0.1203	-0.1490 0.17070 (16)	0.0124 0.11442 (17)	0.050°	
	-0.2217(2)	-0.1/9/0 (10)	0.11445 (17)	0.0379 (3)	
HI8A	-0.1/53	-0.23/4	0.1131	0.08/*	
HI8B	-0.3229	-0.2006	0.0838	0.08/*	
HI8C	-0.2247	-0.1441	0.1731	0.087*	
C139	-0.2078 (2)	-0.02305 (17)	0.06944 (15)	0.0592 (6)	
H19A	-0.2207	0.0115	0.1268	0.089*	
H19B	-0.3050	-0.0466	0.0342	0.089*	
H19C	-0.1469	0.0218	0.0457	0.089*	
C21	0.9228 (7)	0.7212 (7)	0.8184 (4)	0.0348 (4)	0.644 (3)
O21	1.039 (2)	0.767 (3)	0.8666 (11)	0.0455 (13)	0.644 (3)
C22	0.8944 (14)	0.7333 (14)	0.7316 (7)	0.0352 (12)	0.644 (3)
H22	0.9723	0.7693	0.7131	0.042*	0.644 (3)
C23	0.7644 (14)	0.6963 (12)	0.6766 (6)	0.0345 (14)	0.644 (3)
H23	0.6905	0.6548	0.6930	0.041*	0.644 (3)
C211	1.0065 (10)	0.5367 (9)	0.8411 (12)	0.0385 (13)	0.644 (3)
H211	1.0698	0.5772	0.8175	0.046*	0.644 (3)
C212	1.0495 (10)	0.4529 (8)	0.8551 (10)	0.0470 (11)	0.644 (3)
H212	1.1438	0.4383	0.8434	0.056*	0.644 (3)
C213	0.9526 (6)	0.3874 (6)	0.8874 (8)	0.0501 (13)	0.644 (3)
H213	0.9858	0.3322	0.8996	0.060*	0.644 (3)
C214	0.8124 (7)	0.4050 (6)	0.9006 (9)	0.0455 (12)	0.644 (3)
H214	0.7470	0.3587	0.9173	0.055*	0.644 (3)
C215	0.4326 (8)	0.6279(5)	0.9197 (7)	0.0391 (10)	0.644(3)
H215	0.3670	0.5831	0.9380	0.047*	0.644(3)
C216	0 3911 (8)	0.7152 (4)	0.9141(7)	0.0413(12)	0.644(3)
H216	0.2972	0.7292	0.9275	0.050*	0.644(3)
C217	0.2972 0.4909 (10)	0.7252 0.7852 (5)	0.8880 (9)	0.030 0.0407 (10)	0.044(3)
H217	0.4633	0.8461	0.8862	0.040*	0.644(3)
C218	0.6263 (12)	0.0401	0.8655 (12)	0.049	0.044(3)
U210	0.0203 (12)	0.7040 (0)	0.0033 (13)	0.0303 (10)	0.044(3)
П210 С210	0.0002	0.6104	0.0403	0.044	0.044(3)
C219	0.0100 (8)	0.048/(0)	0.040/(7)	0.0312(9)	0.044(3)
0220	0.0222 (7)	0.5165 (5)	0.9079 (7)	0.03/3 (9)	0.644(3)
H220	0.5581	0.4720	0.9275	0.045*	0.644 (3)

C221	0.0(40.(10)	0.5(2((0))	0.0(21.(12)	0.0227 (11)	0 (1)
C221	0.8648 (10)	0.5636 (9)	0.8621 (13)	0.0327(11)	0.644 (3)
C222	0.7624 (10)	0.4933 (8)	0.8892 (12)	0.0366 (8)	0.644 (3)
C223	0.5748 (10)	0.6032 (6)	0.8982 (10)	0.0339 (8)	0.644 (3)
C224	0.6750 (13)	0.6737 (9)	0.8708 (15)	0.0322 (6)	0.644 (3)
C231	0.7290 (9)	0.7158 (7)	0.5930 (4)	0.0336 (11)	0.644 (3)
C232	0.8287 (5)	0.7781 (5)	0.5618 (3)	0.0379 (7)	0.644 (3)
H232	0.9225	0.8087	0.5942	0.046*	0.644 (3)
C233	0.7891 (4)	0.7948 (3)	0.4830 (2)	0.0399 (7)	0.644 (3)
H233	0.8569	0.8372	0.4636	0.048*	0.644 (3)
C234	0.6504 (4)	0.7499 (3)	0.43184 (18)	0.0400 (7)	0.644 (3)
C235	0.5513 (4)	0.6873 (3)	0.4631 (2)	0.0434 (8)	0.644 (3)
H235	0.4581	0.6559	0.4303	0.052*	0.644 (3)
C236	0.5898 (4)	0.6713 (3)	0.5423 (2)	0.0393 (8)	0.644 (3)
H236	0.5212	0.6298	0.5622	0.047*	0.644 (3)
C237	0.6140(3)	0 7666 (3)	0.34365(19)	0.0523(7)	0.644(3)
H237	0.6886	0.8228	0 3403	0.063*	0.644(3)
C238	0.630(3)	0.6736 (15)	0.2717(5)	0.005 0.0767 (12)	0.644(3)
H28A	0.6142	0.6872	0.2717 (5)	0.115*	0.044(3)
1128A 1128A	0.0142	0.6170	0.2109	0.115*	0.044(3)
	0.5550	0.0179	0.2722	0.115*	0.044(3)
П28С	0.7289	0.03/8	0.2813	0.115°	0.044(3)
C239	0.4390 (0)	0.7932 (3)	0.3291 (4)	0.103 (2)	0.044(3)
H29A	0.4493	0.8494	0.3754	0.158*	0.644(3)
H29B	0.3832	0.7371	0.3270	0.158*	0.644 (3)
H29C	0.4460	0.8101	0.2756	0.158*	0.644 (3)
C31	0.9247 (13)	0.7242 (12)	0.8178 (7)	0.0348 (4)	0.356 (3)
O31	1.047 (4)	0.767 (5)	0.861 (2)	0.0455 (13)	0.356 (3)
C32	0.884 (3)	0.737 (3)	0.7321 (12)	0.0352 (12)	0.356 (3)
H32	0.9451	0.7860	0.7159	0.042*	0.356 (3)
C33	0.763 (3)	0.683 (2)	0.6756 (12)	0.0345 (14)	0.356 (3)
H33	0.7045	0.6333	0.6924	0.041*	0.356 (3)
C311	0.9927 (19)	0.5405 (17)	0.851 (2)	0.0385 (13)	0.356 (3)
H311	1.0682	0.5840	0.8382	0.046*	0.356 (3)
C312	1.0239 (18)	0.4537 (15)	0.8647 (19)	0.0470 (11)	0.356 (3)
H312	1.1206	0.4386	0.8608	0.056*	0.356 (3)
C313	0.9114 (16)	0.3855 (12)	0.8845 (17)	0.0501 (13)	0.356 (3)
H313	0.9311	0.3233	0.8876	0.060*	0.356 (3)
C314	0.7757 (16)	0.4111 (12)	0.8990 (17)	0.0455 (12)	0.356 (3)
H314	0.7050	0.3678	0.9157	0.055*	0.356 (3)
C315	0.4254 (15)	0.6532 (10)	0.9175 (15)	0.0391 (10)	0.356 (3)
H315	0 3529	0.6096	0.9326	0.047*	0 356 (3)
C316	0 3952 (16)	0.7435(10)	0.9131 (14)	0.0413(12)	0.356(3)
H316	0.3042	0.7624	0.9273	0.050*	0.356(3)
C317	0.5042	0.7024	0.9275	0.030	0.356(3)
U317 H317	0.3024 (19)	0.8706	0.8833	0.0407 (10)	0.350(3) 0.356(3)
C218	0.537 (2)	0.0700 0.7832 (14)	0.0033	0.079	0.350(3)
U210	0.037 (2)	0.7033 (14)	0.007(2)	0.0303 (10)	0.330(3)
r1318	0.7070	0.020/	0.0323	0.044	0.350(3)
0319	0.8138 (15)	0.0009 (12)	0.8538 (13)	0.0312 (9)	0.356 (3)
C320	0.5989 (14)	0.5309 (10)	0.9031 (15)	0.0375 (9)	0.356 (3)

H320	0.5247	0.4858	0.9155	0.045*	0.356 (3)
C321	0.845 (2)	0.5661 (16)	0.857 (3)	0.0327 (11)	0.356 (3)
C322	0.7383 (19)	0.5032 (15)	0.889(2)	0.0366 (8)	0.356 (3)
C323	0.5666 (19)	0.6235 (12)	0.899 (2)	0.0339 (8)	0.356 (3)
C324	0.671 (2)	0.6876 (16)	0.868 (3)	0.0322 (6)	0.356 (3)
C331	0.7119 (16)	0.6938 (15)	0.5904 (8)	0.0336 (11)	0.356 (3)
C332	0.7896 (11)	0.7611 (10)	0.5540 (6)	0.0379 (7)	0.356 (3)
H332	0.8793	0.8019	0.5842	0.046*	0.356 (3)
C333	0.7358 (7)	0.7684 (6)	0.4732 (4)	0.0399 (7)	0.356 (3)
H333	0.7904	0.8136	0.4500	0.048*	0.356 (3)
C334	0.6015 (7)	0.7093 (5)	0.4264 (3)	0.0400 (7)	0.356 (3)
C335	0.5251 (7)	0.6414 (5)	0.4627 (4)	0.0434 (8)	0.356 (3)
H335	0.4364	0.5998	0.4320	0.052*	0.356 (3)
C336	0.5774 (8)	0.6341 (6)	0.5432 (4)	0.0393 (8)	0.356 (3)
H336	0.5224	0.5889	0.5663	0.047*	0.356 (3)
C337	0.5407 (6)	0.7158 (4)	0.3371 (3)	0.0523 (7)	0.356 (3)
H337	0.4377	0.6773	0.3196	0.063*	0.356 (3)
C338	0.643 (5)	0.671 (3)	0.2714 (9)	0.0767 (12)	0.356 (3)
H38A	0.6069	0.6779	0.2159	0.115*	0.356 (3)
H38B	0.6394	0.6015	0.2677	0.115*	0.356 (3)
H33C	0.7450	0.7061	0.2899	0.115*	0.356 (3)
C339	0.5370 (11)	0.8232 (7)	0.3378 (7)	0.105 (2)	0.356 (3)
H39A	0.4816	0.8247	0.2846	0.158*	0.356 (3)
H39B	0.6387	0.8584	0.3443	0.158*	0.356 (3)
H39C	0.4885	0.8546	0.3849	0.158*	0.356 (3)

Atomic displacement parameters $(Å^2)$

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0341 (8)	0.0458 (10)	0.0339 (8)	0.0065 (7)	0.0038 (6)	0.0054 (7)
O11	0.0388 (7)	0.0710 (10)	0.0470 (7)	0.0172 (6)	0.0010 (5)	-0.0098 (6)
C12	0.0333 (8)	0.0402 (10)	0.0385 (8)	0.0069 (7)	0.0029 (6)	0.0030 (7)
C13	0.0340 (8)	0.0394 (9)	0.0347 (8)	0.0080 (6)	0.0055 (6)	0.0060 (7)
C111	0.0733 (14)	0.0409 (11)	0.0443 (10)	0.0019 (9)	-0.0023 (9)	0.0096 (8)
C112	0.103 (2)	0.0526 (13)	0.0551 (12)	0.0093 (12)	-0.0021 (12)	0.0232 (11)
C113	0.0945 (18)	0.0703 (16)	0.0436 (11)	0.0048 (13)	-0.0097 (11)	0.0228 (11)
C114	0.0621 (12)	0.0557 (12)	0.0372 (9)	-0.0015 (9)	-0.0069 (8)	0.0054 (9)
C115	0.0426 (10)	0.0402 (10)	0.0669 (12)	0.0042 (8)	0.0052 (9)	0.0123 (9)
C116	0.0509 (11)	0.0526 (12)	0.0834 (15)	0.0114 (9)	0.0116 (10)	0.0340 (11)
C117	0.0444 (10)	0.0693 (14)	0.0582 (11)	0.0158 (9)	0.0121 (8)	0.0333 (11)
C118	0.0328 (8)	0.0566 (11)	0.0422 (9)	0.0081 (7)	0.0073 (7)	0.0155 (8)
C119	0.0260 (7)	0.0406 (9)	0.0340 (8)	0.0025 (6)	0.0033 (6)	0.0050 (7)
C120	0.0316 (8)	0.0402 (10)	0.0410 (9)	-0.0021 (7)	-0.0001 (6)	-0.0015 (7)
C121	0.0359 (8)	0.0411 (10)	0.0375 (8)	0.0006 (7)	0.0012 (6)	0.0078 (7)
C122	0.0345 (8)	0.0436 (10)	0.0371 (8)	-0.0012 (7)	0.0000 (6)	0.0061 (7)
C123	0.0262 (7)	0.0400 (9)	0.0477 (9)	0.0020 (6)	0.0050 (6)	0.0103 (7)
C124	0.0229 (7)	0.0436 (9)	0.0396 (8)	0.0042 (6)	0.0060 (6)	0.0117 (7)
C131	0.0341 (8)	0.0324 (8)	0.0321 (8)	0.0033 (6)	0.0033 (6)	0.0084 (6)

G122	0.00(0)	0.00(0)	0.0050 (0)	0.0000 (7)	0.0050 (()	0.0001 (7)
C132	0.0360 (8)	0.0368 (9)	0.0350 (8)	0.0009 (7)	0.0053 (6)	-0.0021(7)
C133	0.0324 (8)	0.0441 (10)	0.0430 (9)	0.0065 (7)	0.0091 (7)	0.0024 (8)
C134	0.0345 (8)	0.0362 (9)	0.0323 (8)	-0.0001 (6)	0.0042 (6)	0.0120 (7)
C135	0.0407 (9)	0.0369 (9)	0.0304 (8)	0.0037 (7)	0.0020 (6)	0.0027 (7)
C136	0.0378 (9)	0.0386 (9)	0.0353 (8)	0.0097 (7)	0.0052 (6)	0.0029 (7)
C137	0.0336 (8)	0.0479 (10)	0.0382 (8)	0.0011 (7)	0.0013 (6)	0.0067 (7)
C138	0.0333 (9)	0.0473 (11)	0.0949 (16)	0.0004 (8)	0.0018 (9)	0.0294 (11)
C139	0.0498 (11)	0.0594 (13)	0.0692 (13)	0.0014 (9)	-0.0123 (10)	0.0320 (11)
C21	0.0289 (8)	0.0351 (10)	0.0393 (8)	0.0075 (6)	0.0055 (6)	0.0073 (7)
O21	0.0344 (19)	0.0502 (8)	0.046 (2)	-0.0007 (16)	-0.001 (2)	0.010 (2)
C22	0.0293 (18)	0.0388 (16)	0.0411 (9)	0.0091 (18)	0.0101 (9)	0.0138 (9)
C23	0.0330 (8)	0.036 (4)	0.0365 (8)	0.0082 (15)	0.0107 (6)	0.0093 (15)
C211	0.0384 (17)	0.0397 (13)	0.035 (4)	0.0100 (11)	0.0113 (12)	0.003 (2)
C212	0.044 (3)	0.0512 (12)	0.050 (4)	0.0248 (19)	0.018 (2)	0.0079 (15)
C213	0.055 (4)	0.0447 (12)	0.0589 (15)	0.025 (3)	0.014 (4)	0.0179 (11)
C214	0.053 (4)	0.0430 (15)	0.0487 (12)	0.017 (2)	0.018 (4)	0.0179 (11)
C215	0.0320 (11)	0.053 (4)	0.0346 (10)	0.0083 (18)	0.0064 (10)	0.015 (3)
C216	0.0313 (10)	0.055 (4)	0.0394 (10)	0.016 (2)	0.0055 (7)	0.011 (3)
C217	0.0403 (17)	0.044 (4)	0.0405 (10)	0.017 (3)	0.0026 (13)	0.012 (3)
C218	0.0358 (16)	0.039 (4)	0.0338 (11)	0.007 (2)	0.0021 (16)	0.009 (4)
C219	0.0310 (8)	0.035 (2)	0.0246 (17)	0.0045 (10)	0.0025 (8)	0.0036 (18)
C220	0.037 (2)	0.043 (2)	0.0362 (16)	0.0079 (12)	0.008 (2)	0.0168 (17)
C221	0.031 (2)	0.0375 (11)	0.027 (2)	0.0063 (14)	0.000 (3)	0.0055 (7)
C222	0.040 (3)	0.040 (2)	0.0331 (8)	0.0126 (13)	0.010 (3)	0.0109 (19)
C223	0.0322 (12)	0.039 (4)	0.0300 (9)	0.0087 (16)	0.0046 (10)	0.008 (3)
C224	0.0304 (9)	0.039 (3)	0.0270 (15)	0.0079 (14)	0.0006 (10)	0.010 (3)
C231	0.0298 (19)	0.037 (5)	0.0360 (9)	0.0092 (12)	0.0104 (9)	0.0099 (18)
C232	0.025 (3)	0.046 (3)	0.0436 (13)	0.0042 (16)	0.0064 (13)	0.0144 (15)
C233	0.032 (2)	0.044 (2)	0.0473 (13)	0.0029 (14)	0.0112 (14)	0.0195 (15)
C234	0.0310 (19)	0.047 (2)	0.0451 (12)	0.0090 (13)	0.0074 (13)	0.0159 (15)
C235	0.0305 (14)	0.056 (3)	0.0450 (11)	0.0028 (17)	0.0042 (10)	0.0187 (18)
C236	0.0304 (11)	0.048 (3)	0.0421 (10)	0.0039 (17)	0.0100 (8)	0.0164 (16)
C237	0.0462 (17)	0.062 (2)	0.0557 (15)	0.0058 (12)	0.0034 (13)	0.0323 (15)
C238	0.095 (4)	0.092(2)	0.0423 (11)	0.0142 (16)	0.0016 (12)	0.0234 (12)
C239	0.083 (4)	0.170(5)	0.116 (3)	0.066 (4)	0.028 (4)	0.105 (4)
C31	0.0289 (8)	0.0351 (10)	0.0393 (8)	0.0075 (6)	0.0055 (6)	0.0073 (7)
031	0.0209(0)	0.0502(8)	0.0353(0)	-0.0007(16)	-0.001(2)	0.0072(7)
C32	0.0293(18)	0.0382(0)	0.0411(9)	0.0091 (18)	0.001(2)	0.013(2)
C33	0.0230(8)	0.036(4)	0.0365(8)	0.0091(10) 0.0082(15)	0.0107(6)	0.0190(9)
C311	0.0384(17)	0.030(1) 0.0397(13)	0.035(4)	0.0100(11)	0.0107(0)	0.003(2)
C312	0.0304(17)	0.0577(13)	0.050(4)	0.0100(11) 0.0248(19)	0.0113(12)	0.003(2)
C312	0.044(3)	0.0312(12) 0.0447(12)	0.050(4)	0.0240(1))	0.013(2)	0.0079(13)
C314	0.053(4)	0.0447(12) 0.0430(15)	0.0387(13)	0.023(3)	0.014(4)	0.0179(11)
C315	0.033(+) 0.0320(11)	0.053 (4)	0.0707 (12)	0.017(2) 0.0083(18)	0.016(-1)	0.0179(11) 0.015(3)
C316	0.0320(11) 0.0313(10)	0.055(4)	0.0340(10) 0.0304(10)	0.0003(10)	0.000+(10)	0.013(3)
C317	0.0313(10) 0.0403(17)	0.033(4)	0.0394(10) 0.0405(10)	0.010(2) 0.017(3)	0.0035(7)	0.011(3) 0.012(3)
C318	0.0703(17) 0.0358(16)	0.039(4)	0.0703(10) 0.0338(11)	0.017(3)	0.0020(13) 0.0021(16)	0.012(3)
C310	0.0330(10)	0.037(4)	0.0330(11) 0.0246(17)	0.007(2)	0.0021(10)	0.009(4)
0319	0.0310(8)	0.035 (2)	0.0240 (17)	0.0045 (10)	0.0025 (8)	0.0030 (18)

C320	0.037 (2)	0.043 (2)	0.0362 (16)	0.0079 (12)	0.008 (2)	0.0168 (17)
C321	0.031 (2)	0.0375 (11)	0.027 (2)	0.0063 (14)	0.000 (3)	0.0055 (7)
C322	0.040 (3)	0.040 (2)	0.0331 (8)	0.0126 (13)	0.010 (3)	0.0109 (19)
C323	0.0322 (12)	0.039 (4)	0.0300 (9)	0.0087 (16)	0.0046 (10)	0.008 (3)
C324	0.0304 (9)	0.039 (3)	0.0270 (15)	0.0079 (14)	0.0006 (10)	0.010 (3)
C331	0.0298 (19)	0.037 (5)	0.0360 (9)	0.0092 (12)	0.0104 (9)	0.0099 (18)
C332	0.025 (3)	0.046 (3)	0.0436 (13)	0.0042 (16)	0.0064 (13)	0.0144 (15)
C333	0.032 (2)	0.044 (2)	0.0473 (13)	0.0029 (14)	0.0112 (14)	0.0195 (15)
C334	0.0310 (19)	0.047 (2)	0.0451 (12)	0.0090 (13)	0.0074 (13)	0.0159 (15)
C335	0.0305 (14)	0.056 (3)	0.0450 (11)	0.0028 (17)	0.0042 (10)	0.0187 (18)
C336	0.0304 (11)	0.048 (3)	0.0421 (10)	0.0039 (17)	0.0100 (8)	0.0164 (16)
C337	0.0462 (17)	0.062 (2)	0.0557 (15)	0.0058 (12)	0.0034 (13)	0.0323 (15)
C338	0.095 (4)	0.092 (2)	0.0423 (11)	0.0142 (16)	0.0016 (12)	0.0234 (12)
C339	0.083 (4)	0.170 (5)	0.116 (3)	0.066 (4)	0.028 (4)	0.105 (4)

Geometric parameters (Å, °)

C11—011	1.215 (2)	C219—C221	1.413 (3)
C11—C12	1.473 (2)	C220—C223	1.390 (3)
C11—C119	1.509 (2)	C220—C222	1.392 (3)
C12—C13	1.332 (2)	C220—H220	0.9300
С12—Н12	0.9300	C221—C222	1.441 (3)
C13—C131	1.460 (2)	C223—C224	1.435 (3)
С13—Н13	0.9300	C231—C232	1.393 (4)
C111—C112	1.358 (3)	C231—C236	1.394 (4)
C111—C121	1.425 (3)	C232—C233	1.383 (3)
C111—H111	0.9300	С232—Н232	0.9300
C112—C113	1.412 (3)	C233—C234	1.394 (4)
С112—Н112	0.9300	С233—Н233	0.9300
C113—C114	1.347 (3)	C234—C235	1.392 (4)
С113—Н113	0.9300	C234—C237	1.523 (3)
C114—C122	1.423 (3)	C235—C236	1.382 (3)
C114—H114	0.9300	С235—Н235	0.9300
C115—C116	1.352 (3)	С236—Н236	0.9300
C115—C123	1.419 (3)	C237—C239	1.507 (5)
С115—Н115	0.9300	C237—C238	1.545 (11)
C116—C117	1.413 (3)	С237—Н237	0.9800
C116—H116	0.9300	C238—H28A	0.9600
C117—C118	1.359 (3)	C238—H28B	0.9600
С117—Н117	0.9300	C238—H28C	0.9600
C118—C124	1.426 (2)	С239—Н29А	0.9600
C118—H118	0.9300	C239—H29B	0.9600
C119—C121	1.405 (2)	С239—Н29С	0.9600
C119—C124	1.406 (2)	C31—O31	1.221 (4)
C120—C122	1.386 (3)	C31—C32	1.468 (4)
C120—C123	1.393 (3)	C31—C319	1.510 (4)
С120—Н120	0.9300	C32—C33	1.345 (6)
C121—C122	1.440 (2)	C32—H32	0.9300

C123—C124	1.436 (2)	C33—C331	1.465 (4)
C131—C132	1.392 (2)	С33—Н33	0.9300
C131—C136	1.395 (2)	C311—C312	1.356 (4)
C132—C133	1.376 (2)	C311—C321	1.433 (5)
С132—Н132	0.9300	С311—Н311	0.9300
C133—C134	1.390 (2)	C312—C313	1.417 (5)
С133—Н133	0.9300	С312—Н312	0.9300
C134—C135	1.386 (2)	C313—C314	1.349 (5)
C134—C137	1.517 (2)	С313—Н313	0.9300
C135—C136	1.387 (2)	C314—C322	1.427 (4)
С135—Н135	0.9300	C314—H314	0.9300
С136—Н136	0.9300	C315—C316	1.353 (5)
C137—C139	1.518 (3)	C315—C323	1.430 (4)
C137—C138	1.526 (3)	С315—Н315	0.9300
C137—H137	0.9800	C316—C317	1.416 (5)
C138—H18A	0.9600	С316—Н316	0.9300
C138—H18B	0.9600	C317—C318	1.358 (4)
C138—H18C	0.9600	С317—Н317	0.9300
C139—H19A	0.9600	C318—C324	1.429 (5)
C139—H19B	0.9600	C318—H318	0.9300
С139—Н19С	0.9600	C319—C324	1.412 (4)
C21—O21	1.222 (3)	C319—C321	1.414 (5)
C21—C22	1.466 (3)	C320—C323	1.390 (4)
C21—C219	1.509 (3)	C320—C322	1.390 (4)
C22—C23	1.346 (5)	С320—Н320	0.9300
С22—Н22	0.9300	C321—C322	1.443 (4)
C23—C231	1.467 (3)	C323—C324	1.436 (4)
С23—Н23	0.9300	C331—C332	1.390 (5)
C211—C212	1.356 (3)	C331—C336	1.396 (5)
C211—C221	1.432 (4)	C332—C333	1.387 (4)
C211—H211	0.9300	С332—Н332	0.9300
C212—C213	1.417 (4)	C333—C334	1.390 (5)
C212—H212	0.9300	С333—Н333	0.9300
C213—C214	1.350 (4)	C334—C335	1.389 (5)
С213—Н213	0.9300	C334—C337	1.528 (4)
C214—C222	1.428 (3)	C335—C336	1.380 (5)
C214—H214	0.9300	С335—Н335	0.9300
C215—C216	1.354 (4)	С336—Н336	0.9300
C215—C223	1.430 (3)	C337—C339	1.511 (6)
С215—Н215	0.9300	C337—C338	1.549 (13)
C216—C217	1.417 (4)	С337—Н337	0.9800
C216—H216	0.9300	С338—Н38А	0.9600
C217—C218	1.358 (3)	C338—H38B	0.9600
C217—H217	0.9300	С338—Н33С	0.9600
C218—C224	1.428 (3)	С339—Н39А	0.9600
C218—H218	0.9300	С339—Н39В	0.9600
C219—C224	1.411 (3)	С339—Н39С	0.9600

O11—C11—C12	123.20 (16)	C214—C222—C221	118.5 (2)
O11—C11—C119	120.18 (15)	C220—C223—C215	122.1 (3)
C12—C11—C119	116.61 (14)	C220—C223—C224	119.4 (3)
C13—C12—C11	121.47 (16)	C215—C223—C224	118.5 (2)
C13—C12—H12	119.3	C219—C224—C218	122.9 (3)
C11—C12—H12	119.3	C219—C224—C223	119.0 (2)
C12—C13—C131	127.53 (16)	C218—C224—C223	118.1 (2)
C12—C13—H13	116.2	C232—C231—C236	117.9 (2)
C131—C13—H13	116.2	C232—C231—C23	122.7 (3)
C112—C111—C121	121.36 (19)	C236—C231—C23	119.4 (3)
C112—C111—H111	119.3	C233—C232—C231	120.4 (3)
C121—C111—H111	119.3	C233—C232—H232	119.8
C111—C112—C113	120.6 (2)	C231—C232—H232	119.8
C111—C112—H112	119.7	$C_{232} - C_{233} - C_{234}$	121.8(3)
C113—C112—H112	119.7	$C_{232} = C_{233} = H_{233}$	119.1
C114 - C113 - C112	120 14 (19)	C234—C233—H233	119.1
C114—C113—H113	119.9	$C_{235} = C_{234} = C_{233}$	117.6(2)
$C_{112} = C_{113} = H_{113}$	110.0	$C_{235} = C_{234} = C_{235}$	117.0(2) 122.2(3)
$C_{112} - C_{113} - C_{113}$	119.9	$C_{233} = C_{234} = C_{237}$	122.2(3) 120.2(3)
$C_{113} = C_{114} = C_{122}$	121.74 (10)	$C_{235} - C_{234} - C_{234}$	120.2(3)
C122 C114 H114	119.1	$C_{230} - C_{235} - C_{234}$	120.8 (5)
C122 - C114 - H114	119.1	$C_{230} - C_{235} - H_{235}$	119.0
C116—C115—C125	121.34 (19)	$C_{234} = C_{235} = H_{235}$	119.0
	119.2	$C_{235} - C_{236} - C_{231}$	121.5 (5)
C123—C115—H115	119.2	C235—C236—H236	119.3
C115—C116—C117	120.09 (19)	C231—C236—H236	119.3
C115—C116—H116	120.0	C239—C237—C234	112.7 (3)
C117—C116—H116	120.0	C239—C237—C238	110.2 (9)
C118—C117—C116	120.72 (18)	C234—C237—C238	110.5 (4)
C118—C117—H117	119.6	С239—С237—Н237	107.7
C116—C117—H117	119.6	С234—С237—Н237	107.7
C117—C118—C124	121.02 (18)	С238—С237—Н237	107.7
C117—C118—H118	119.5	С237—С238—Н28А	109.5
C124—C118—H118	119.5	С237—С238—Н28В	109.5
C121—C119—C124	121.64 (15)	H28A—C238—H28B	109.5
C121—C119—C11	119.09 (15)	С237—С238—Н28С	109.5
C124—C119—C11	119.28 (14)	H28A—C238—H28C	109.5
C122—C120—C123	122.30 (15)	H28B—C238—H28C	109.5
C122—C120—H120	118.9	С237—С239—Н29А	109.5
C123—C120—H120	118.9	С237—С239—Н29В	109.5
C119—C121—C111	123.58 (16)	H29A—C239—H29B	109.5
C119—C121—C122	118.70 (16)	C237—C239—H29C	109.5
C111—C121—C122	117.71 (16)	H29A—C239—H29C	109.5
C120-C122-C114	122.24 (16)	H29B-C239-H29C	109 5
$C_{120} - C_{122} - C_{121}$	119 33 (16)	031 - C31 - C32	120.0(5)
C114 $C122$ $C121$	118 43 (17)	031 - C31 - C319	120.0(3)
C120 - C123 - C115	122 21 (17)	C_{32} C_{31} C_{319}	119.8(5)
C120 - C123 - C124	119 17 (16)	C_{33} C_{32} C_{31} C_{31}	1741(7)
$C_{120} = C_{120} = C_{1$	118.62 (16)	$C_{33} = C_{32} = C_{31}$	118.0
0113-0123-0124	110.02 (10)	033-032-032	110.0

C119—C124—C118	123.16 (16)	C31—C32—H32	118.0
C119—C124—C123	118.85 (15)	C32—C33—C331	127.1 (7)
C118—C124—C123	117.99 (16)	С32—С33—Н33	116.4
C132—C131—C136	117.53 (15)	С331—С33—Н33	116.4
C132—C131—C13	122.65 (15)	C312—C311—C321	120.9 (5)
C136—C131—C13	119.81 (15)	С312—С311—Н311	119.5
C133—C132—C131	121.31 (15)	С321—С311—Н311	119.5
C133—C132—H132	119.3	C311—C312—C313	121.0 (5)
C131—C132—H132	119.3	С311—С312—Н312	119.5
C132—C133—C134	121.27 (16)	C313—C312—H312	119.5
С132—С133—Н133	119.4	C314—C313—C312	119.9 (5)
C134—C133—H133	119.4	C314—C313—H313	120.0
C135—C134—C133	117.76 (15)	С312—С313—Н313	120.0
C135—C134—C137	120.87 (15)	C313—C314—C322	121.4 (5)
C133—C134—C137	121.37 (15)	C313—C314—H314	119.3
C134—C135—C136	121.19 (15)	C322—C314—H314	119.3
С134—С135—Н135	119.4	C316—C315—C323	121.4 (5)
C136—C135—H135	119.4	C316—C315—H315	119.3
C135—C136—C131	120.92 (16)	С323—С315—Н315	119.3
C135—C136—H136	119.5	C315—C316—C317	119.8 (5)
C131—C136—H136	119.5	C315—C316—H316	120.1
C134—C137—C139	113.05 (15)	C317—C316—H316	120.1
C134—C137—C138	110.88 (15)	C318—C317—C316	121.1 (5)
C139—C137—C138	109.53 (16)	C318—C317—H317	119.5
C134—C137—H137	107.7	C316—C317—H317	119.5
С139—С137—Н137	107.7	C317—C318—C324	121.0 (5)
С138—С137—Н137	107.7	C317—C318—H318	119.5
C137—C138—H18A	109.5	C324—C318—H318	119.5
C137—C138—H18B	109.5	C324—C319—C321	121.0 (5)
H18A—C138—H18B	109.5	C324—C319—C31	120.4 (5)
C137—C138—H18C	109.5	C321—C319—C31	117.7 (5)
H18A—C138—H18C	109.5	C323—C320—C322	122.3 (5)
H18B—C138—H18C	109.5	С323—С320—Н320	118.9
С137—С139—Н19А	109.5	С322—С320—Н320	118.9
С137—С139—Н19В	109.5	C319—C321—C311	122.2 (8)
H19A—C139—H19B	109.5	C319—C321—C322	118.6 (5)
С137—С139—Н19С	109.5	C311—C321—C322	117.3 (6)
H19A—C139—H19C	109.5	C320—C322—C314	122.4 (5)
H19B—C139—H19C	109.5	C320—C322—C321	118.9 (4)
O21—C21—C22	119.9 (3)	C314—C322—C321	118.5 (4)
O21—C21—C219	119.8 (3)	C320—C323—C315	122.0 (5)
C22—C21—C219	120.2 (3)	C320—C323—C324	119.2 (4)
C23—C22—C21	125.1 (5)	C315—C323—C324	118.6 (4)
C23—C22—H22	117.4	C319—C324—C318	122.6 (6)
C21—C22—H22	117.4	C319—C324—C323	118.8 (5)
C22—C23—C231	125.7 (4)	C318—C324—C323	117.7 (5)
С22—С23—Н23	117.1	C332—C331—C336	117.8 (4)
C231—C23—H23	117.1	C332—C331—C33	123.7 (5)
			· · ·

C212—C211—C221	120.8 (3)	C336—C331—C33	118.5 (5)
C212—C211—H211	119.6	C333—C332—C331	121.2 (5)
C221—C211—H211	119.6	С333—С332—Н332	119.4
C211—C212—C213	120.9 (3)	C331—C332—H332	119.4
C211—C212—H212	119.5	C332—C333—C334	121.1 (5)
C213—C212—H212	119.5	C332—C333—H333	119.5
C214—C213—C212	120.2 (3)	C334—C333—H333	119.5
C214—C213—H213	119.9	C335—C334—C333	117.5 (4)
C212—C213—H213	119.9	C335—C334—C337	120.6 (5)
C_{213} C_{214} C_{222}	121 3 (3)	$C_{333} - C_{334} - C_{337}$	121.9(5)
C213—C214—H214	119.4	$C_{336} - C_{335} - C_{334}$	121.9(0) 121.8(4)
C_{222} C_{214} H_{214}	119.4	C336—C335—H335	119.1
C_{216} C_{215} C_{223}	121 5 (3)	C_{334} C_{335} H_{335}	119.1
$C_{216} = C_{215} = H_{215}$	119.3	$C_{335} - C_{336} - C_{331}$	120.7(5)
C_{223} C_{215} H_{215}	119.3	C335_C336_H336	119.7
$C_{215} = C_{216} = C_{217}$	120.0(3)	C331—C336—H336	119.7
$C_{215} = C_{216} = C_{217}$	120.0 (3)	$C_{330} = C_{337} = C_{334}$	110.9 (5)
$C_{213} = C_{216} = H_{216}$	120.0	$C_{339} = C_{337} = C_{334}$	110.9(3)
$C_{217} = C_{210} = H_{210}$	120.0 120.8(3)	$C_{334} = C_{337} = C_{338}$	109.2(12)
$C_{218} = C_{217} = C_{210}$	120.8 (3)	$C_{334} = C_{337} = C_{338}$	109.0 (0)
$C_{216} = C_{217} = H_{217}$	119.0	$C_{339} = C_{337} = H_{337}$	109.0
$C_{210} = C_{217} = H_{217}$	119.0	$C_{228} = C_{227} = H_{227}$	109.0
$C_{217} = C_{218} = C_{224}$	121.1 (5)	$C_{330} = C_{337} = H_{337}$	109.0
$C_{21} = C_{218} = H_{218}$	119.4	$C_{33}/-C_{338}$ -H38A	109.5
C224—C218—H218	119.4	U28A C228 U28D	109.5
$C_{224} = C_{219} = C_{221}$	120.7(3)	H38A—C338—H38B	109.5
$C_{224} = C_{219} = C_{21}$	120.0(3)		109.5
$C_{221} = C_{219} = C_{21}$	117.9 (3)	H38A—C338—H33C	109.5
$C_{223} = C_{220} = C_{222}$	122.2 (3)	H38B—C338—H33C	109.5
C223—C220—H220	118.9	C337—C339—H39A	109.5
C222—C220—H220	118.9	C337—C339—H39B	109.5
C219—C221—C211	122.8 (3)	H39A—C339—H39B	109.5
C219—C221—C222	118.8 (3)	С337—С339—Н39С	109.5
C211—C221—C222	117.8 (3)	H39A—C339—H39C	109.5
C220—C222—C214	122.3 (3)	Н39В—С339—Н39С	109.5
C220—C222—C221	119.1 (3)		
			176 6 (12)
011-011-012-013	-3.2(3)	C216—C215—C223—C220	1/6.6 (13)
C119 - C11 - C12 - C13	177.72 (16)	$C_{216} - C_{215} - C_{223} - C_{224}$	0(2)
CII_CI2_CI3_CI3I	1/9.25 (1/)	C221—C219—C224—C218	172.8 (19)
C121—C111—C112—C113	0.5 (4)	C21—C219—C224—C218	4 (3)
C111—C112—C113—C114	0.0 (4)	C221—C219—C224—C223	-10 (3)
C112—C113—C114—C122	-0.5 (4)	C21—C219—C224—C223	-179.4 (14)
C123—C115—C116—C117	-0.2(3)	C217—C218—C224—C219	179.0 (18)
C115—C116—C117—C118	0.0 (3)	C217—C218—C224—C223	2 (3)
C116—C117—C118—C124	1.1 (3)	C220—C223—C224—C219	6 (3)
O11—C11—C119—C121	-66.5(2)	C215—C223—C224—C219	-178.0 (16)
C12—C11—C119—C121	112.60 (18)	C220—C223—C224—C218	-177.0 (18)
O11—C11—C119—C124	113.3 (2)	C215—C223—C224—C218	-1 (3)

C12—C11—C119—C124	-67.5 (2)	C22—C23—C231—C232	-1 (3)
C124—C119—C121—C111	-178.78 (18)	C22—C23—C231—C236	179.3 (19)
C11—C119—C121—C111	1.1 (3)	C236—C231—C232—C233	0.2 (15)
C124—C119—C121—C122	1.3 (2)	C23—C231—C232—C233	-179.3 (13)
C11—C119—C121—C122	-178.81 (15)	C231—C232—C233—C234	-0.6 (12)
C112—C111—C121—C119	179.5 (2)	C232—C233—C234—C235	0.3 (7)
C112—C111—C121—C122	-0.6 (3)	C232—C233—C234—C237	-177.3(5)
C123—C120—C122—C114	179.13 (18)	C233—C234—C235—C236	0.4 (6)
C123—C120—C122—C121	-0.5 (3)	C237—C234—C235—C236	178.0 (3)
C113—C114—C122—C120	-179.2(2)	C234—C235—C236—C231	-0.9(8)
C113—C114—C122—C121	0.4 (3)	C_{232} C_{231} C_{236} C_{235}	0.6 (14)
$C_{119} - C_{121} - C_{122} - C_{120}$	-0.3(2)	C_{23} C_{231} C_{236} C_{235}	-180.0(12)
$C_{111} - C_{121} - C_{122} - C_{120}$	179 76 (18)	C_{235} C_{234} C_{237} C_{239}	50.0 (5)
$C_{119} - C_{121} - C_{122} - C_{114}$	-17999(17)	$C_{233} = C_{234} = C_{237} = C_{239}$	-1325(5)
$C_{111} - C_{121} - C_{122} - C_{114}$	01(3)	$C_{235} = C_{234} = C_{237} = C_{238}$	-73.9(11)
C_{122} C_{120} C_{123} C_{115}	-179.93(17)	C_{233} C_{234} C_{237} C_{238}	103.7(12)
$C_{122} = C_{120} = C_{123} = C_{124}$	04(3)	031 - C31 - C32 - C33	170 (6)
$C_{116} - C_{115} - C_{123} - C_{120}$	179 68 (18)	$C_{319} - C_{31} - C_{32} - C_{33}$	-12(5)
$C_{116} - C_{115} - C_{123} - C_{124}$	-0.6(3)	$C_{31} - C_{32} - C_{33} - C_{331}$	12(3)
$C_{121} - C_{119} - C_{124} - C_{118}$	17772(15)	$C_{321} = C_{311} = C_{312} = C_{313}$	0(5)
C11-C119-C124-C118	-21(2)	$C_{311} = C_{312} = C_{313} = C_{314}$	-7(5)
$C_{121} - C_{119} - C_{124} - C_{123}$	-14(2)	C_{312} C_{313} C_{314} C_{322}	4(5)
C11-C119-C124-C123	178 69 (14)	$C_{323} = C_{315} = C_{316} = C_{317}$	2(4)
$C_{117} - C_{118} - C_{124} - C_{119}$	178.90 (16)	$C_{315} = C_{316} = C_{317} = C_{318}$	-1(4)
C_{117} C_{118} C_{124} C_{123}	-19(2)	$C_{316} = C_{317} = C_{318} = C_{324}$	3(5)
C_{120} C_{123} C_{124} C_{119}	0.6(2)	031 - C31 - C319 - C324	116(5)
$C_{115} - C_{123} - C_{124} - C_{119}$	-17911(15)	C_{32} C_{31} C_{319} C_{324}	-61(4)
C_{120} C_{123} C_{124} C_{118}	-178.63(15)	031 - C31 - C319 - C321	-75(5)
$C_{115} - C_{123} - C_{124} - C_{118}$	17(2)	C_{32} C_{31} C_{319} C_{321}	107(3)
C_{12} C_{13} C_{131} C_{132}	-41(3)	$C_{324} = C_{319} = C_{321} = C_{311}$	-174(3)
C_{12} C_{13} C_{131} C_{136}	175 54 (17)	C_{31} C_{319} C_{321} C_{311}	18 (4)
C_{136} C_{131} C_{132} C_{133}	-0.5(3)	$C_{324} C_{319} C_{321} C_{322}$	-10(5)
C_{13} $-C_{131}$ $-C_{132}$ $-C_{133}$	179 11 (17)	C_{31} C_{319} C_{321} C_{322} C_{322} C_{323} C_{333} C_{33	-178(3)
$C_{131} - C_{132} - C_{133} - C_{134}$	-0.1(3)	C_{312} C_{311} C_{321} C_{319} C_{319}	172(3)
C_{132} C_{133} C_{134} C_{135}	0.1(3)	$C_{312} = C_{311} = C_{321} = C_{322}$	8(5)
C_{132} C_{133} C_{134} C_{137}	-179.87(17)	$C_{323} = C_{320} = C_{322} = C_{314}$	175(3)
$C_{133} - C_{134} - C_{135} - C_{136}$	-0.3(3)	$C_{323} = C_{320} = C_{322} = C_{321}$	-9(5)
C_{137} C_{134} C_{135} C_{136}	-179.92(16)	C_{313} C_{314} C_{322} C_{320} C_{320}	180(3)
C_{134} C_{135} C_{136} C_{131}	-0.3(3)	C_{313} C_{314} C_{322} C_{321}	4(5)
C_{132} C_{131} C_{136} C_{135}	0.8(3)	$C_{319} = C_{321} = C_{322} = C_{320}$	9 (5)
C_{13} C_{131} C_{136} C_{135}	-178.92(16)	$C_{311} = C_{321} = C_{322} = C_{320}$	174(3)
$C_{135} - C_{134} - C_{137} - C_{139}$	-137.63(18)	$C_{319} C_{321} C_{322} C_{314}$	-175(3)
C_{133} C_{134} C_{137} C_{139}	42.8(2)	$C_{311} - C_{321} - C_{322} - C_{314}$	-10(5)
C_{135} C_{134} C_{137} C_{138}	98.9 (2)	C_{322} C_{320} C_{323} C_{315}	-177(3)
C133—C134—C137—C138	-80.7 (2)	C322—C320—C323—C324	9 (5)
021—C21—C22—C23	-172 (3)	C316—C315—C323—C320	-180(3)
C219—C21—C22—C23	11 (3)	C316—C315—C323—C324	-6 (5)
C21—C22—C23—C231	173.8 (18)	C321—C319—C324—C318	179 (4)
	···· 、 ···		

C221—C211—C212—C213	3 (3)	C31—C319—C324—C318	-13 (5)
C211—C212—C213—C214	3 (2)	C321—C319—C324—C323	10 (5)
C212—C213—C214—C222	-5 (2)	C31—C319—C324—C323	178 (3)
C223—C215—C216—C217	-1 (2)	C317—C318—C324—C319	-175 (3)
C215—C216—C217—C218	2 (2)	C317—C318—C324—C323	-6 (6)
C216—C217—C218—C224	-3 (3)	C320—C323—C324—C319	-9 (5)
O21—C21—C219—C224	106 (3)	C315—C323—C324—C319	177 (3)
C22—C21—C219—C224	-77.3 (19)	C320—C323—C324—C318	-178 (3)
O21—C21—C219—C221	-64 (3)	C315—C323—C324—C318	7 (5)
C22—C21—C219—C221	113.1 (16)	C32—C33—C331—C332	3 (6)
C224—C219—C221—C211 C214—C219—C221—C211 C224—C219—C221—C212 C214—C219—C221—C222 C212—C211—C221—C222 C212—C211—C221—C222 C223—C220—C222—C214 C223—C220—C222—C221	$\begin{array}{c} -178.3 (18) \\ -9 (2) \\ 10 (2) \\ -180.0 (13) \\ -178.4 (17) \\ -7 (3) \\ 178.9 (15) \\ 3 (3) \end{array}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{c} -176 (4) \\ 0 (3) \\ -180 (3) \\ 1 (2) \\ -1.3 (14) \\ -179.7 (10) \\ 1.7 (12) \\ -179.9 (7) \end{array}$
C213—C214—C222—C220 C213—C214—C222—C221 C219—C221—C222—C220 C211—C221—C222—C220 C219—C221—C222—C214 C211—C221—C222—C214 C222—C220—C223—C215 C222—C220—C223—C224	$\begin{array}{c} -175.6 (15) \\ 1 (3) \\ -7 (3) \\ -178.3 (17) \\ 177.0 (16) \\ 5 (3) \\ -178.2 (15) \\ -2 (3) \end{array}$	C334—C335—C336—C331 C332—C331—C336—C335 C33—C331—C336—C335 C335—C334—C337—C339 C333—C334—C337—C339 C335—C334—C337—C338 C333—C334—C337—C338	$\begin{array}{c} -1.4 \ (18) \\ 1 \ (3) \\ -180 \ (2) \\ 131.4 \ (7) \\ -50.3 \ (9) \\ -108 \ (2) \\ 70 \ (2) \end{array}$

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 are the centroids of rings (C111–C114/C122/C111) and (C114–C118/C124/C113), respectively.

D—H···A	D—H	$H \cdots A$	$D \cdots A$	D—H··· A
C133—H133…O11 ⁱ	0.93	2.49	3.336 (2)	151
C216—H216…O21 ⁱ	0.93	2.61	3.41 (2)	144
C216—H216…O31 ⁱ	0.93	2.58	3.36 (4)	142
C316—H316…O21 ⁱ	0.93	2.51	3.30 (3)	144
C316—H316…O31 ⁱ	0.93	2.47	3.25 (4)	142
C233—H233…Cg1 ⁱⁱ	0.93	2.64	3.355 (4)	134
C236—H236…Cg2 ⁱⁱⁱ	0.93	2.75	3.519 (4)	140
C336—H336…Cg2 ⁱⁱⁱ	0.93	2.80	3.354 (7)	119

Symmetry codes: (i) *x*-1, *y*, *z*; (ii) -*x*+2, -*y*+1, -*z*+1; (iii) -*x*+1, -*y*+1, -*z*+1.