

An infinite two-dimensional hybrid water–chloride network in a 4'-(furan-2-yl)-2,2':6',2''-terpyridine nickel(II) matrix

Wei-Wei Fu,* Ya-Qian Li, Yang Liu, Man-Sheng Chen, Wei Li and Ying-Qun Yang

Received 26 April 2017

Accepted 13 May 2017

Edited by A. J. Lough, University of Toronto, Canada

Keywords: crystal structure; terpyridine; Ni^{II}; water cluster; water–chloride network; hydrogen bonds.

CCDC reference: 1498201


Supporting information: this article has supporting information at journals.iucr.org/e

Key Laboratory of Functional Organometallic Materials of General Colleges and Universities in Hunan Province, Department of Chemistry and Materials Science, Hengyang Normal University, Hengyang 421008, People's Republic of China. *Correspondence e-mail: w.w.fu@hotmail.com

A new complex, namely bis[4'-(furan-2-yl)-2,2':6',2''-terpyridine]nickel(II) dichloride decahydrate, [Ni(C₁₉H₁₃N₃O)₂]Cl₂·10H₂O, has been crystallized by solvent evaporation and characterized by single-crystal X-ray diffraction. The coordination environment of the Ni^{II} cation is distorted octahedral with slight deviations from an idealized geometry. The most intriguing structural feature is an infinite two-dimensional hybrid water–chloride network parallel to (011) constructed by O–H···O and O–H···Cl hydrogen bonds involving two independent chloride ions and ten independent solvent water molecules with an L-shaped pattern. One of the furyl rings is disordered with a refined occupancy ratio of 0.786 (13):0.214 (13)

1. Chemical context

Water has received much scientific interest as it is a major chemical constituent on the earth's surface and it is also the source of life. Many discrete water clusters and polymeric water aggregates, with different types of hydrogen bonds and in diverse sizes and shapes, captured in the crystal lattice of an organic or metal coordination complex during crystallization have been found and investigated experimentally and theoretically (Dutta *et al.*, 2015; Ganguly & Mondal, 2015; Han *et al.*, 2014; Hundal *et al.*, 2014; Pati *et al.*, 2014).



Hybrid water–chloride associates incorporated in various crystal matrixes are one of the most interesting combinations in water clusters research due to their fundamental importance for understanding water–halide interactions in the atmosphere, the ocean and in biological systems (Inumaru *et al.*, 2008; Kumar *et al.*, 2011; Lakshminarayanan *et al.*, 2006; Li *et al.*, 2008). According to a search of the Cambridge Structural Database (CSD Version 5.37, May 2016; Groom *et al.*,




Table 1
 Selected geometric parameters (Å, °).

Ni1—N5	1.974 (3)	Ni1—N3	2.096 (3)
Ni1—N2	1.977 (3)	Ni1—N1	2.098 (3)
Ni1—N6	2.093 (3)	Ni1—N4	2.099 (3)
N5—Ni1—N2	178.36 (10)	N2—Ni1—N1	77.77 (11)
N5—Ni1—N6	77.81 (11)	N6—Ni1—N1	93.13 (11)
N2—Ni1—N6	102.65 (11)	N3—Ni1—N1	155.38 (11)
N5—Ni1—N3	100.71 (11)	N5—Ni1—N4	78.10 (11)
N2—Ni1—N3	77.74 (11)	N2—Ni1—N4	101.46 (12)
N6—Ni1—N3	89.84 (11)	N6—Ni1—N4	155.89 (11)
N5—Ni1—N1	103.80 (10)	N3—Ni1—N4	95.46 (11)

2016), there are about nine examples with water–chloride hydrogen bonds forming one-dimensional tapes (Boyer *et al.*, 2011; van Holst *et al.*, 2008; Kepert *et al.*, 1999; Jitsukawa *et al.*, 1994), two-dimensional (Kepert *et al.*, 1994; Chowdhury *et al.*, 2011; Duan *et al.*, 2016) and three-dimensional (Figgis *et al.*, 1983; Pruchnik *et al.*, 1996) networks from 2,2':6',2''-terpyridine ligands. When 4'-substituted terpyridines with phenyl, pyridyl, imidazolyl rings were considered, two-dimensional and three-dimensional water–chloride networks with two chloride ions and at least six water molecules were found (Constable *et al.*, 1990; Kou *et al.*, 2008; Chen *et al.*, 2013; Fernandes *et al.*, 2008; McMurtrie & Dance, 2010; Padhi *et al.*, 2010; Indumathy *et al.*, 2008; Mahendiran *et al.*, 2016). The hydrophobic and hydrophilic layers are further linked by two kinds of C—H...O hydrogen bonds into three-dimensional networks. In this context, a ftpy–Ni^{II} complex [ftpy = 4'-(furan-2-yl)-2,2':6',2''-terpyridine] (Fig. 1) with two chlorides as counter-ions and ten solvent water molecules (**1**) is described herein.

2. Structural commentary

The asymmetric unit of **1** is composed of a cationic [Ni(ftpy)₂]²⁺ part, two chloride anions, and ten water molecules of crystallization. The distances between Ni1 and the N atoms of the central pyridyl rings [1.974 (3) and 1.977 (3) Å] are slightly shorter than those between Ni1 and the N atoms of outer pyridyl rings [2.093 (3)–2.099 (3) Å; Table 1]. The angles involving Ni1 can be divided into two sets, *viz.* three




Figure 1
 The molecular structure of [Ni(ftpy)₂]²⁺ in **1**, with displacement ellipsoids drawn at the 30% probability level.




Figure 2
 View of the hydrophobic (represented by wireframes) and hydrophilic (represented by spheres) layers in **1**.

transoid angles [178.36 (10), 155.38 (11) and 155.89 (11)°] and 12 cisoid angles, which range from 77.74 (11) to 103.80 (10)°. The differences in the bond lengths and angles indicate a distorted octahedral geometry (Constable *et al.*, 1990; Logacheva *et al.*, 2009; Padhi *et al.*, 2010; Fu *et al.*, 2013). The terpyridyl ring systems [maximum deviations of ±0.058 (4) Å for C27/C31 and 0.192 (4) Å for C17] are almost perpendicular to each other, subtending a dihedral angle of 87.35 (6)°. The furyl rings are almost coplanar with the terpyridyl ring systems, making dihedral angles of 8.1 (2) and 3.2 (3)° for the O1- and O2-containing rings, respectively.

3. Supramolecular features

In the crystal, there are hydrophobic layers composed of [Ni(ftpy)₂]²⁺ dications and hydrophilic layers composed of water molecules and chloride anions (Fig. 2). In the hydrophobic layers, shown in Fig. 3, [Ni(ftpy)₂]²⁺ dications are linked by two kinds of face-to-face π – π interactions with centroid–centroid distances of 3.530 (4) and 3.760 (4) Å between the furyl and outer pyridyl rings, forming one-





Figure 3
 A view of the two-dimensional undulating sheet of hydrophobic layers, with π – π interactions highlighted by dashed lines [purple for 3.533 (5) and 3.761 (4) Å, and green for 4.338 (14) and 4.405 (4) Å].


Figure 4

A view of the hybrid water–chloride hydrogen-bonded assemblies in **1**, with water molecules and chloride anions shown as coloured balls and hydrogen bonds as dashed lines.

dimensional (1D) chains. These 1D chains are linked by further π – π interactions with centroid distances of 4.367 (4) Å between furyl rings and 4.405 (4) Å between furyl and central pyridyl rings, forming two-dimensional networks. The water molecules and chloride anions form a two-dimensional network parallel to (011) *via* O–H···O and O–H···Cl hydrogen bonds (Table 2), as shown in Fig. 4.

The multicyclic $\{[(\text{H}_2\text{O})_{10}\text{Cl}_2]^{2-}\}_n$ fragments in the hydrophilic layers are constructed by means of 11 non-equivalent O–H···O hydrogen bonds with O···O distances ranging from 2.756 (6) to 3.134 (7) Å and nine O–H···Cl hydrogen bonds with O···Cl distances ranging from 3.079 (4) to 3.225 (4) Å (Table 2, Fig. 4). Both the O···O and O···Cl distances are comparable with those found in various types of water clusters and water–chloride associates (Safin *et al.*, 2015; Bhat & Revankar, 2016; Ris *et al.*, 2016). The resulting two-dimensional network can be considered as a set of alternating cyclic fragments with three tetranuclear, three pentanuclear, one hexanuclear and two octanuclear fragments, as shown in Fig. 5*a*. Two of these fragments are composed only of water molecules, whereas the other seven rings are water–chloride hybrids with one or two Cl[−] anions. Most of the rings are non-planar, contributing to the formation of an intricate relief geometry of the water–chloride layer. Using the method described by Infantes and co-workers (Infantes & Motherwell, 2002; Infantes *et al.*, 2003), this two-dimensional water–chloride network can be described as having an L4(6)4(6)4(6)5(5)5(6)5(6)6(8)8(8)8(10) pattern.

4. Comparison with other terpyridine complexes possessing 10 solvent water molecules

It is interesting to make a comparison of the two-dimensional water–chloride networks in **1** and those found in other terpyridine complexes possessing 10 solvent water molecules, *viz.* [Fe(phtpy)₂]Cl₂·10H₂O (**2**; refcode: VOBKON; Fernandes *et al.*, 2008), [Ni(phtpy)₂]Cl₂·10H₂O, (**3**; refcode: SIXLIU01; Chen *et al.*, 2013), [Ru(phtpy)₂]Cl₂·10H₂O (**4**; refcode: FAFFID; McMurtrie & Dance, 2010) and [Ru(pytpy)₂]Cl₂·10H₂O (**5**; refcode: TUXGUP; Padhi *et al.*, 2010) [phtpy = 4′-phenyl-2,2′:6′,2′′-terpyridine and pytpy = 4′-(2-pyridyl)-

Table 2

Hydrogen-bond geometry (Å, °).

D–H···A	D–H	H···A	D···A	D–H···A
O1W–H1WA···Cl1	0.87	2.25	3.113 (4)	169
O1W–H1WB···O9W ⁱ	0.87	2.06	2.923 (6)	175
O2W–H2WB···O5W ⁱⁱ	0.83	1.99	2.813 (7)	172
O2W–H2WA···Cl1	0.84	2.39	3.215 (4)	168
O3W–H3WC···O4W	0.86	2.05	2.760 (9)	140
O3W–H3WA···O6W ⁱⁱⁱ	0.88	2.35	3.134 (7)	148
O4W–H4WB···Cl2	0.88	2.58	3.107 (5)	119
O4W–H4WA···Cl2	0.87	2.56	3.107 (5)	122
O5W–H5WA···Cl2	0.87	2.37	3.079 (4)	138
O5W–H5WB···O9W	0.89	2.16	2.991 (6)	156
O6W–H6WC···O2W ⁱⁱ	0.83	2.11	2.929 (6)	167
O6W–H6WA···O7W	0.83	2.18	2.838 (6)	136
O7W–H7WA···Cl2	0.87	2.34	3.190 (4)	167
O7W–H7WB···O4W ⁱⁱ	0.87	1.93	2.798 (5)	172
O8W–H8WC···O3W ⁱⁱ	0.85	2.06	2.856 (8)	155
O8W–H8WD···Cl2 ^{iv}	0.85	2.40	3.204 (6)	157
O9W–H9WA···O10W ^v	0.86	1.93	2.756 (6)	159
O9W–H9WB···O1W ^{vi}	0.86	2.11	2.878 (5)	147
O10W–H10A···Cl1 ^{vii}	0.88	2.27	3.141 (4)	171
O10W–H10B···Cl1 ^{viii}	0.87	2.38	3.225 (4)	165

Symmetry codes: (i) $-x, -y + 1, -z + 1$; (ii) $-x + 1, -y + 1, -z + 1$; (iii) $x - 1, y, z$; (iv) $x + 1, y, z$; (v) $-x + 1, -y + 1, -z$; (vi) $x, y + 1, z - 1$; (vii) $x, y, z - 1$; (viii) $-x + 1, -y, -z + 1$.

2,2′:6′,2′′-terpyridine]. In spite of the differences in the metal ions and terpyridine ligands, the crystal parameters are almost the same for compounds **2–5**. Where a five-membered furyl ring is involved instead of a six-membered phenyl or pyridyl ring, the size of the crystal cell decreases with reduction in the cell volume of about 4.5% from 2200 to 2100 Å³. Considering the O···O and O···Cl distances within the two-dimensional water–chloride networks, a different number of trinuclear, tetranuclear, pentanuclear, hexanuclear and octanuclear rings have been determined, giving an L4(6)4(6)4(6)4(6)4(6)5(6)5(6)5(6)6(8)8(12) pattern for **2**, an L4(6)4(6)4(6)5(7)5(7)5(8)5(8)6(7)6(9)6(9)8(12) pattern for **3**, an L4(6)4(6)4(6)4(6)4(6)4(6)5(6)5(6)5(7)6(7)8(12) pattern for **4** and an L3(6)4(6)5(5)5(6)5(6)6(8)6(8)8(8)8(10) pattern for **5** (Fig. 5*b–e*). These results illustrate how a water–chloride assembly could be fine-tuned by adopting diverse ligands and different metal ions. It is potentially useful for future studies of water–water or water–chloride interactions for chemists as well as theoreticians.

5. Synthesis and crystallization

4′-Furyl-2,2′:6′,2′′-terpyridine was prepared by a literature method (Wang & Hanan, 2005). Other reagents and solvents used in reactions were purchased from Aladdin Chemical and used without purification, unless otherwise indicated.

NiCl₂·6H₂O (0.1 mmol, 0.024g) and ftpy (0.2 mmol, 0.060 g) were dissolved in 10 ml distilled water and 10 ml methanol. The solution was left alone for slow evaporation without disturbance for about one month and reddish brown crystals of (**1**) suitable for X-ray analysis were obtained.




Figure 5
Multicyclic $\{[(\text{H}_2\text{O})_{10}\text{Cl}_2]^{2-}\}_n$ fragments with repeating units of two-dimensional water–chloride networks in (a) **1**, (b) **2**, (c) **3**, (d) **4** and (e) **5**.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. All hydrogen atoms except those of water molecules were generated geometrically and refined isotropically using a riding model, with $\text{C}–\text{H} = 0.93 \text{ \AA}$ and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The hydrogen atoms of solvent water molecules were located in difference-Fourier maps, refined with DFIX restraints of O–H distances and finally fixed at those positions using AFIX 3 in *SHELXL* (Sheldrick, 2015b). Atoms C36, C37, C38 and O2 were found to be disordered over two sets of sites with a refined occupancy ratio of 0.786 (13):0.214 (13) for C36/C36A, C37/C37A, C38/C38A, and O2/O2A. In order to model the disorder of this furyl ring, various restraints (DFIX, FLAT, ISOR, DELU, EADP) were applied in the refinement.

Acknowledgements

Financial support by the Key Discipline Project of Hunan Province, the Open Fund of the Key Laboratory of Functional

Organometallic Materials of Hunan Province College (GN14K02), the Aid program for Science and Technology Innovative Research Team in Higher Educational Institutions of Hunan Province and the Scientific Research Fund of Hunan Provincial Education Department (16B037) and Key Laboratory of Catalysis and Materials Science of the State Ethnic Affairs Commission & Ministry of Education (CHCL16002) are gratefully acknowledged.

References

- Bhat, S. S. & Revankar, V. K. (2016). *J. Chem. Crystallogr.* **46**, 9–14.
 Boyer, J. L., Polyansky, D. E., Szalda, D. J., Zong, R., Thummel, R. P. & Fujita, E. (2011). *Angew. Chem. Int. Ed.* **50**, 12600–12604.
 Brandenburg, K. & Putz, H. (2008). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Bruker (2012). *SMART, SAINT and SADABS*. Bruker AXS inc., Madison, Wisconsin, USA.
 Chen, G.-J., Wang, Z.-G., Kou, Y.-Y., Tian, J.-L. & Yan, S.-P. (2013). *J. Inorg. Biochem.* **122**, 49–56.
 Chowdhury, A. D., Das, A. K. I., Mobin, S. M. & Lahiri, G. K. (2011). *Inorg. Chem.* **50**, 1775–1785.

Table 3

Experimental details.

Crystal data	
Chemical formula	[Ni(C ₁₉ H ₁₃ N ₃ O ₂) ₂]Cl ₂ ·10H ₂ O
<i>M_r</i>	908.42
Crystal system, space group	Triclinic, <i>P</i> $\bar{1}$
Temperature (K)	296
<i>a</i> , <i>b</i> , <i>c</i> (Å)	10.351 (7), 11.894 (8), 19.070 (13)
α , β , γ (°)	76.33 (1), 88.582 (12), 67.077 (11)
<i>V</i> (Å ³)	2095 (2)
<i>Z</i>	2
Radiation type	Mo <i>K</i> α
μ (mm ⁻¹)	0.66
Crystal size (mm)	0.23 × 0.18 × 0.15
Data collection	
Diffractometer	Bruker SMART CCD area-detector
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2012)
<i>T</i> _{min} , <i>T</i> _{max}	0.864, 0.908
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	10779, 7382, 5322
<i>R</i> _{int}	0.029
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.597
Refinement	
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.050, 0.144, 1.07
No. of reflections	7382
No. of parameters	546
No. of restraints	75
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}$, $\Delta\rho_{\text{min}}$ (e Å ⁻³)	0.42, -0.53

Computer programs: *SMART* and *SAINT* (Bruker, 2012), *SHELXT* (Sheldrick, 2015a); *SHELXL2014* (Sheldrick, 2015b), *OLEX2* (Dolomanov *et al.*, 2009), *DIAMOND* (Brandenburg & Putz, 2008) and *publCIF* (Westrip, 2010).

- Constable, E. C., Lewis, J., Liptrot, M. C. & Raithby, P. R. (1990). *Inorg. Chim. Acta*, **178**, 47–54.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Duan, L., Manbeck, G. F., Kowalczyk, M., Szalda, D. J., Muckerman, J. T., Himeda, Y. & Fujita, E. (2016). *Inorg. Chem.* **55**, 4582–4594.
- Dutta, R., Akhuli, B. & Ghosh, P. (2015). *Dalton Trans.* **44**, 15075–15078.
- Fernandes, R. R., Kirillov, A. M., da Silva, M. F. C. G., Ma, Z., da Silva, J. A. L., da Silva, J. J. R. F. & Pombeiro, A. J. L. (2008). *Cryst. Growth Des.* **8**, 782–785.
- Figgis, B., Kucharski, E. & White, A. (1983). *Aust. J. Chem.* **36**, 1563–1571.
- Fu, W.-W., Kuang, D.-Z., Zhang, F.-X., Liu, Y., Li, W. & Kuang, Y.-F. (2013). *Chin. J. Inorg. Chem.* **29**, 654–658.
- Ganguly, S. & Mondal, R. (2015). *Cryst. Growth Des.* **15**, 2211–2222.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst. B* **72**, 171–179.
- Han, L.-L., Hu, T.-P., Chen, J.-S., Li, Z.-H., Wang, X.-P., Zhao, Y.-Q., Li, X.-Y. & Sun, D. (2014). *Dalton Trans.* **43**, 8774–8780.
- Holst, M. van, Le Pevelen, D. & Aldrich-Wright, J. (2008). *Eur. J. Inorg. Chem.* pp. 4608–4615.
- Hundal, G., Hundal, M. S., Hwang, Y. K. & Chang, J. S. (2014). *Cryst. Growth Des.* **14**, 172–176.
- Indumathy, R., Kanthimathi, M., Weyhermuller, T. & Nair, B. U. (2008). *Polyhedron*, **27**, 3443–3450.
- Infantes, L., Chisholm, J. & Motherwell, S. (2003). *CrystEngComm*, **5**, 480–486.
- Infantes, L. & Motherwell, S. (2002). *CrystEngComm*, **4**, 454–461.
- Inumaru, K., Kikudome, T., Fukuoka, H. & Yamanaka, S. (2008). *J. Am. Chem. Soc.* **130**, 10038–10039.
- Jitsukawa, K., Hata, T., Yamamoto, T., Kano, K., Masuda, H. & Einaga, H. (1994). *Chem. Lett.* **23**, 1169–1172.
- Keper, C., Lu, W., Skelton, B. & White, A. (1994). *Aust. J. Chem.* **47**, 365–384.
- Keper, C. J., Semenova, L. I., Wei-Min, L., Skelton, B. W. & White, A. H. (1999). *Aust. J. Chem.* **52**, 481–496.
- Kou, Y.-Y., Gu, W., Liu, H., Ma, X.-F., Li, D.-D. & Yan, S.-P. (2008). *Acta Sci. Nat. Univ. Nankaiensis*, **41**, 19–25.
- Kumar, R., Pandey, A. K., Sharma, M. K., Panicker, L. V., Sodaye, S., Suresh, G., Ramagiri, S. V., Bellare, J. R. & Goswami, A. (2011). *J. Phys. Chem. B*, **115**, 5856–5867.
- Lakshminarayanan, P. S., Suresh, E. & Ghosh, P. (2006). *Angew. Chem. Int. Ed.* **45**, 3807–3811.
- Li, Y., Jiang, L., Feng, X.-L. & Lu, T.-B. (2008). *Cryst. Growth Des.* **8**, 3689–3694.
- Logacheva, N. M., Baulin, V. E., Tsvadze, A. Y., Pyatova, E. N., Ivanova, I. S., Velikodny, Y. A. & Chernyshev, V. V. (2009). *Dalton Trans.* pp. 2482–2489.
- Mahendiran, D., Kumar, R. S., Viswanathan, V., Velmurugan, D. & Rahiman, A. K. (2016). *Dalton Trans.* **45**, 7794–7814.
- McMurtrie, J. & Dance, I. (2010). *CrystEngComm*, **12**, 3207–3217.
- Padhi, S. K., Sahu, R. & Manivannan, V. (2010). *Polyhedron*, **29**, 709–714.
- Pati, A., Athilakshmi, J., Ramkumar, V. & Chand, D. K. (2014). *CrystEngComm*, **16**, 6827–6830.
- Pruchnik, F. P., Robert, F., Jeannin, Y. & Jeannin, S. (1996). *Inorg. Chem.* **35**, 4261–4263.
- Ris, D. P., Schneider, G. E., Ertl, C. D., Kohler, E., Müntener, T., Neuburger, M., Constable, E. C. & Housecroft, C. E. (2016). *J. Organomet. Chem.* **812**, 272–279.
- Safin, D. A., Szell, P. M. J., Keller, A., Korobkov, I., Bryce, D. L. & Murugesu, M. (2015). *New J. Chem.* **39**, 7147–7152.
- Sheldrick, G. M. (2015a). *Acta Cryst. A* **71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst. C* **71**, 3–8.
- Wang, J. & Hanan, G. S. (2005). *Synlett*, pp. 1251–1254.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2017). E73, 871-875 [https://doi.org/10.1107/S2056989017007095]

An infinite two-dimensional hybrid water–chloride network in a 4'-(furan-2-yl)-2,2':6',2''-terpyridine nickel(II) matrix

Wei-Wei Fu, Ya-Qian Li, Yang Liu, Man-Sheng Chen, Wei Li and Ying-Qun Yang

Computing details

Data collection: *SMART* (Bruker, 2012); cell refinement: *SMART* (Bruker, 2012); data reduction: *SAINTE* (Bruker, 2012); program(s) used to solve structure: *SHELXT* (Sheldrick, 2015a) and *OLEX2* (Dolomanov *et al.*, 2009); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015b) and *OLEX2* (Dolomanov *et al.*, 2009); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2008); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Bis[4'-(furan-2-yl)-2,2':6',2''-terpyridine]nickel(II) dichloride decahydrate

Crystal data

[Ni(C₁₉H₁₃N₃O)₂]Cl₂·10H₂O

$M_r = 908.42$

Triclinic, *P1*

$a = 10.351$ (7) Å

$b = 11.894$ (8) Å

$c = 19.070$ (13) Å

$\alpha = 76.33$ (1)°

$\beta = 88.582$ (12)°

$\gamma = 67.077$ (11)°

$V = 2095$ (2) Å³

$Z = 2$

$F(000) = 948$

$D_x = 1.440$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3615 reflections

$\theta = 2.2$ – 24.0 °

$\mu = 0.66$ mm⁻¹

$T = 296$ K

Block, brown

$0.23 \times 0.18 \times 0.15$ mm

Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube
phi and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2012)

$T_{\min} = 0.864$, $T_{\max} = 0.908$

10779 measured reflections

7382 independent reflections

5322 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\max} = 25.1$ °, $\theta_{\min} = 1.9$ °

$h = -12 \rightarrow 8$

$k = -14 \rightarrow 13$

$l = -22 \rightarrow 19$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.050$

$wR(F^2) = 0.144$

$S = 1.07$

7382 reflections

546 parameters

75 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: mixed

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0689P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.42 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.53 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ni1	0.97894 (4)	0.78241 (4)	0.74747 (2)	0.03697 (15)	
Cl1	0.30308 (10)	0.17463 (10)	0.94851 (6)	0.0626 (3)	
Cl2	0.28593 (15)	0.70315 (14)	0.37352 (7)	0.0975 (4)	
N1	0.9435 (3)	0.6158 (2)	0.77056 (14)	0.0395 (6)	
N2	1.0808 (3)	0.6936 (2)	0.84370 (14)	0.0371 (6)	
N3	1.0524 (3)	0.9124 (2)	0.76977 (14)	0.0402 (6)	
N4	1.1361 (3)	0.7137 (2)	0.67860 (15)	0.0404 (6)	
N5	0.8802 (3)	0.8743 (2)	0.65098 (14)	0.0363 (6)	
N6	0.7778 (3)	0.8893 (3)	0.77339 (14)	0.0399 (6)	
O1W	0.0399 (3)	0.1312 (3)	1.0094 (2)	0.1017 (12)	
H1WA	0.1069	0.1479	0.9865	0.153*	
H1WB	-0.0184	0.1283	0.9785	0.153*	
O1	1.3786 (3)	0.5497 (3)	1.07455 (16)	0.0771 (8)	
O2W	0.3405 (4)	0.3554 (3)	0.8020 (2)	0.1191 (14)	
H2WB	0.4256	0.3390	0.7997	0.179*	
H2WA	0.3280	0.3179	0.8431	0.179*	
O2	0.7323 (4)	1.0325 (5)	0.38766 (18)	0.0652 (12)	0.786 (13)
O2A	0.5318 (13)	1.1546 (17)	0.4459 (8)	0.0652 (12)	0.214 (13)
O3W	0.0344 (6)	0.4187 (5)	0.4364 (3)	0.161 (2)	
H3WC	0.1242	0.3849	0.4377	0.241*	
H3WA	-0.0004	0.4792	0.3967	0.241*	
O4W	0.2936 (5)	0.4343 (4)	0.4358 (2)	0.1321 (15)	
H4WB	0.2366	0.5098	0.4401	0.198*	
H4WA	0.3534	0.4658	0.4168	0.198*	
O5W	0.3714 (4)	0.7092 (4)	0.21731 (19)	0.1075 (12)	
H5WA	0.3229	0.7484	0.2484	0.161*	
H5WB	0.3079	0.7751	0.1866	0.161*	
O6W	0.8002 (4)	0.6442 (4)	0.3295 (2)	0.1187 (14)	
H6WC	0.7594	0.6559	0.2898	0.178*	
H6WA	0.7339	0.6982	0.3442	0.178*	
O7W	0.5809 (3)	0.6957 (3)	0.42459 (18)	0.0892 (10)	
H7WA	0.5069	0.6842	0.4137	0.134*	
H7WB	0.6169	0.6500	0.4680	0.134*	
O8W	0.9848 (5)	0.7051 (5)	0.4183 (3)	0.165 (2)	
H8WC	0.9741	0.6910	0.4634	0.247*	
H8WD	1.0541	0.7266	0.4086	0.247*	

O9W	0.1697 (4)	0.8720 (3)	0.08918 (18)	0.0972 (11)
H9WA	0.2328	0.8596	0.0583	0.146*
H9WB	0.1381	0.9527	0.0824	0.146*
O10W	0.6127 (3)	0.1247 (3)	-0.00361 (17)	0.0790 (9)
H10A	0.5244	0.1474	-0.0182	0.119*
H10B	0.6177	0.0479	0.0111	0.119*
C1	1.4491 (5)	0.4603 (6)	1.1365 (3)	0.0893 (16)
H1A	1.4970	0.4732	1.1724	0.107*
C2	1.4381 (5)	0.3551 (6)	1.1366 (2)	0.0892 (16)
H2A	1.4785	0.2800	1.1719	0.107*
C3	1.3555 (4)	0.3735 (4)	1.0750 (2)	0.0629 (10)
H3A	1.3284	0.3149	1.0618	0.075*
C4	1.3234 (4)	0.4928 (4)	1.03888 (19)	0.0502 (8)
C5	1.2401 (3)	0.5626 (3)	0.97208 (18)	0.0439 (8)
C6	1.2338 (3)	0.6820 (3)	0.93761 (18)	0.0450 (8)
H6	1.2842	0.7184	0.9575	0.054*
C7	1.1634 (3)	0.5118 (3)	0.93943 (18)	0.0435 (8)
H7	1.1657	0.4322	0.9610	0.052*
C8	1.0858 (3)	0.5796 (3)	0.87613 (17)	0.0400 (7)
C9	1.1521 (3)	0.7449 (3)	0.87409 (17)	0.0386 (7)
C10	1.1305 (3)	0.8734 (3)	0.83215 (18)	0.0405 (8)
C11	1.1802 (4)	0.9497 (4)	0.8570 (2)	0.0519 (9)
H11	1.2359	0.9201	0.9003	0.062*
C12	1.1457 (4)	1.0711 (4)	0.8163 (2)	0.0591 (10)
H12	1.1772	1.1250	0.8319	0.071*
C13	1.0652 (4)	1.1115 (4)	0.7531 (2)	0.0574 (10)
H13	1.0410	1.1932	0.7251	0.069*
C14	1.0200 (4)	1.0298 (3)	0.7311 (2)	0.0506 (9)
H14	0.9649	1.0577	0.6877	0.061*
C15	0.9995 (3)	0.5383 (3)	0.83507 (18)	0.0384 (7)
C16	0.9774 (4)	0.4297 (3)	0.8607 (2)	0.0485 (9)
H16	1.0166	0.3774	0.9060	0.058*
C17	0.8971 (4)	0.3996 (4)	0.8187 (2)	0.0568 (10)
H17	0.8794	0.3274	0.8356	0.068*
C18	0.8430 (4)	0.4757 (4)	0.7518 (2)	0.0557 (10)
H18	0.7900	0.4550	0.7223	0.067*
C19	0.8678 (4)	0.5830 (3)	0.7287 (2)	0.0491 (9)
H19	0.8315	0.6347	0.6828	0.059*
C20	1.2646 (4)	0.6268 (3)	0.6978 (2)	0.0507 (9)
H20	1.2924	0.5921	0.7468	0.061*
C21	1.3586 (4)	0.5861 (4)	0.6483 (2)	0.0624 (11)
H21	1.4483	0.5246	0.6635	0.075*
C22	1.3193 (4)	0.6362 (4)	0.5779 (2)	0.0605 (10)
H22	1.3821	0.6104	0.5435	0.073*
C23	1.1853 (4)	0.7262 (4)	0.5561 (2)	0.0515 (9)
H23	1.1565	0.7615	0.5073	0.062*
C24	1.0956 (3)	0.7623 (3)	0.60840 (18)	0.0389 (7)
C25	0.9482 (3)	0.8536 (3)	0.59157 (17)	0.0383 (7)

C26	0.8800 (4)	0.9120 (3)	0.52461 (18)	0.0438 (8)	
H26	0.9271	0.8958	0.4836	0.053*	
C27	0.7407 (4)	0.9953 (3)	0.51755 (18)	0.0421 (8)	
C28	0.6734 (4)	1.0179 (3)	0.57974 (18)	0.0423 (8)	
H28	0.5803	1.0746	0.5767	0.051*	
C29	0.7472 (3)	0.9549 (3)	0.64582 (17)	0.0367 (7)	
C30	0.6892 (3)	0.9652 (3)	0.71648 (17)	0.0378 (7)	
C31	0.5555 (4)	1.0467 (3)	0.7236 (2)	0.0488 (8)	
H31	0.4960	1.0997	0.6831	0.059*	
C32	0.5115 (4)	1.0483 (4)	0.7922 (2)	0.0551 (10)	
H32	0.4211	1.1021	0.7987	0.066*	
C33	0.6021 (4)	0.9698 (4)	0.8509 (2)	0.0549 (10)	
H33	0.5743	0.9694	0.8977	0.066*	
C34	0.7340 (4)	0.8921 (3)	0.83933 (19)	0.0499 (9)	
H34	0.7956	0.8391	0.8791	0.060*	
C35	0.6651 (4)	1.0581 (3)	0.44799 (17)	0.0473 (8)	
C36	0.5368 (6)	1.1455 (5)	0.4263 (4)	0.0530 (14)	0.786 (13)
H36	0.4693	1.1807	0.4566	0.064*	0.786 (13)
C36A	0.695 (2)	1.046 (2)	0.3801 (7)	0.0530 (14)	0.214 (13)
H36A	0.7779	0.9894	0.3672	0.064*	0.214 (13)
C37	0.5193 (7)	1.1758 (6)	0.3518 (4)	0.0582 (16)	0.786 (13)
H37	0.4390	1.2338	0.3231	0.070*	0.786 (13)
C37A	0.585 (3)	1.130 (2)	0.3344 (7)	0.0582 (16)	0.214 (13)
H37A	0.5793	1.1393	0.2846	0.070*	0.214 (13)
C38	0.6383 (8)	1.1068 (6)	0.3293 (2)	0.0593 (15)	0.786 (13)
H38	0.6561	1.1081	0.2811	0.071*	0.786 (13)
C38A	0.4838 (19)	1.198 (2)	0.3714 (10)	0.0593 (15)	0.214 (13)
H38A	0.3980	1.2620	0.3516	0.071*	0.214 (13)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0399 (3)	0.0386 (3)	0.0332 (2)	-0.01552 (19)	0.00434 (16)	-0.01032 (18)
Cl1	0.0538 (6)	0.0650 (6)	0.0640 (6)	-0.0216 (5)	-0.0007 (4)	-0.0095 (5)
Cl2	0.1043 (10)	0.1146 (11)	0.0714 (9)	-0.0447 (9)	0.0106 (7)	-0.0168 (8)
N1	0.0398 (15)	0.0438 (16)	0.0373 (16)	-0.0172 (13)	0.0044 (12)	-0.0131 (13)
N2	0.0400 (15)	0.0369 (15)	0.0358 (15)	-0.0158 (12)	0.0047 (11)	-0.0110 (12)
N3	0.0430 (16)	0.0411 (16)	0.0405 (16)	-0.0189 (13)	0.0082 (12)	-0.0138 (13)
N4	0.0399 (16)	0.0407 (15)	0.0411 (17)	-0.0147 (13)	0.0069 (12)	-0.0136 (13)
N5	0.0383 (15)	0.0357 (14)	0.0361 (15)	-0.0151 (12)	0.0085 (11)	-0.0106 (12)
N6	0.0434 (15)	0.0457 (16)	0.0337 (15)	-0.0192 (13)	0.0088 (12)	-0.0132 (13)
O1W	0.075 (2)	0.106 (3)	0.121 (3)	-0.032 (2)	0.034 (2)	-0.033 (2)
O1	0.080 (2)	0.091 (2)	0.0604 (19)	-0.0284 (18)	-0.0039 (15)	-0.0255 (17)
O2W	0.153 (4)	0.072 (2)	0.096 (3)	-0.015 (2)	0.047 (3)	-0.008 (2)
O2	0.054 (2)	0.083 (3)	0.0378 (19)	-0.010 (2)	-0.0021 (15)	-0.0050 (18)
O2A	0.054 (2)	0.083 (3)	0.0378 (19)	-0.010 (2)	-0.0021 (15)	-0.0050 (18)
O3W	0.202 (5)	0.220 (6)	0.127 (4)	-0.143 (5)	0.047 (4)	-0.068 (4)
O4W	0.120 (3)	0.150 (4)	0.113 (4)	-0.051 (3)	-0.009 (3)	-0.011 (3)

O5W	0.149 (4)	0.128 (3)	0.073 (2)	-0.083 (3)	0.012 (2)	-0.025 (2)
O6W	0.114 (3)	0.157 (4)	0.141 (4)	-0.090 (3)	0.056 (3)	-0.079 (3)
O7W	0.085 (2)	0.081 (2)	0.084 (2)	-0.0156 (18)	-0.0003 (17)	-0.0184 (18)
O8W	0.163 (5)	0.214 (6)	0.121 (4)	-0.081 (4)	0.011 (3)	-0.034 (4)
O9W	0.097 (2)	0.104 (3)	0.092 (3)	-0.049 (2)	0.031 (2)	-0.015 (2)
O10W	0.0620 (18)	0.094 (2)	0.085 (2)	-0.0325 (17)	-0.0038 (15)	-0.0244 (18)
C1	0.064 (3)	0.129 (5)	0.055 (3)	-0.010 (3)	-0.018 (2)	-0.034 (3)
C2	0.089 (4)	0.093 (4)	0.042 (2)	0.003 (3)	-0.0113 (18)	-0.003 (2)
C3	0.075 (3)	0.0587 (19)	0.050 (2)	-0.022 (2)	0.0019 (17)	-0.0116 (17)
C4	0.047 (2)	0.0585 (18)	0.0374 (19)	-0.0128 (18)	0.0000 (14)	-0.0124 (14)
C5	0.0400 (19)	0.048 (2)	0.0362 (19)	-0.0093 (16)	0.0054 (14)	-0.0112 (16)
C6	0.0435 (19)	0.051 (2)	0.044 (2)	-0.0188 (17)	0.0034 (15)	-0.0183 (17)
C7	0.050 (2)	0.0385 (19)	0.0375 (19)	-0.0151 (16)	0.0043 (15)	-0.0055 (15)
C8	0.0401 (18)	0.0429 (19)	0.0377 (19)	-0.0152 (16)	0.0080 (14)	-0.0135 (16)
C9	0.0388 (18)	0.0428 (19)	0.0337 (18)	-0.0132 (15)	0.0066 (13)	-0.0141 (15)
C10	0.0414 (18)	0.0447 (19)	0.042 (2)	-0.0201 (16)	0.0089 (14)	-0.0181 (16)
C11	0.055 (2)	0.058 (2)	0.052 (2)	-0.0278 (19)	0.0027 (17)	-0.0200 (19)
C12	0.064 (3)	0.056 (2)	0.073 (3)	-0.034 (2)	0.014 (2)	-0.028 (2)
C13	0.071 (3)	0.043 (2)	0.063 (3)	-0.027 (2)	0.007 (2)	-0.0150 (19)
C14	0.055 (2)	0.045 (2)	0.048 (2)	-0.0179 (18)	0.0037 (16)	-0.0072 (17)
C15	0.0359 (17)	0.0386 (18)	0.0415 (19)	-0.0134 (15)	0.0070 (14)	-0.0139 (15)
C16	0.048 (2)	0.041 (2)	0.054 (2)	-0.0163 (17)	0.0054 (16)	-0.0071 (17)
C17	0.059 (2)	0.046 (2)	0.073 (3)	-0.0288 (19)	0.007 (2)	-0.016 (2)
C18	0.054 (2)	0.055 (2)	0.068 (3)	-0.029 (2)	0.0026 (19)	-0.022 (2)
C19	0.051 (2)	0.053 (2)	0.047 (2)	-0.0222 (18)	-0.0010 (16)	-0.0158 (18)
C20	0.044 (2)	0.049 (2)	0.056 (2)	-0.0134 (18)	0.0049 (17)	-0.0156 (18)
C21	0.038 (2)	0.064 (3)	0.077 (3)	-0.0086 (19)	0.0091 (19)	-0.023 (2)
C22	0.050 (2)	0.067 (3)	0.066 (3)	-0.018 (2)	0.024 (2)	-0.030 (2)
C23	0.050 (2)	0.059 (2)	0.047 (2)	-0.0196 (19)	0.0147 (16)	-0.0205 (18)
C24	0.0402 (18)	0.0378 (18)	0.043 (2)	-0.0171 (15)	0.0103 (14)	-0.0145 (15)
C25	0.0433 (19)	0.0414 (18)	0.0358 (18)	-0.0197 (16)	0.0114 (14)	-0.0154 (15)
C26	0.051 (2)	0.049 (2)	0.0360 (19)	-0.0231 (18)	0.0153 (15)	-0.0148 (16)
C27	0.050 (2)	0.0430 (19)	0.0353 (19)	-0.0212 (17)	0.0044 (14)	-0.0087 (15)
C28	0.0429 (19)	0.0409 (19)	0.040 (2)	-0.0133 (16)	0.0039 (14)	-0.0102 (15)
C29	0.0415 (19)	0.0349 (17)	0.0358 (18)	-0.0157 (15)	0.0086 (14)	-0.0121 (14)
C30	0.0422 (19)	0.0366 (18)	0.0377 (19)	-0.0169 (15)	0.0076 (14)	-0.0128 (15)
C31	0.047 (2)	0.047 (2)	0.048 (2)	-0.0144 (17)	0.0088 (16)	-0.0121 (17)
C32	0.051 (2)	0.059 (2)	0.060 (3)	-0.0197 (19)	0.0225 (19)	-0.027 (2)
C33	0.063 (2)	0.063 (2)	0.044 (2)	-0.025 (2)	0.0233 (18)	-0.023 (2)
C34	0.061 (2)	0.056 (2)	0.0343 (19)	-0.0238 (19)	0.0103 (16)	-0.0136 (17)
C35	0.058 (2)	0.057 (2)	0.0349 (19)	-0.029 (2)	0.0046 (16)	-0.0142 (17)
C36	0.057 (3)	0.050 (3)	0.049 (3)	-0.016 (2)	0.016 (3)	-0.019 (3)
C36A	0.057 (3)	0.050 (3)	0.049 (3)	-0.016 (2)	0.016 (3)	-0.019 (3)
C37	0.045 (3)	0.057 (3)	0.060 (3)	-0.010 (3)	-0.003 (3)	-0.010 (3)
C37A	0.045 (3)	0.057 (3)	0.060 (3)	-0.010 (3)	-0.003 (3)	-0.010 (3)
C38	0.056 (3)	0.074 (4)	0.033 (2)	-0.014 (3)	-0.004 (2)	-0.005 (2)
C38A	0.056 (3)	0.074 (4)	0.033 (2)	-0.014 (3)	-0.004 (2)	-0.005 (2)

Geometric parameters (Å, °)

Ni1—N5	1.974 (3)	C7—H7	0.9300
Ni1—N2	1.977 (3)	C8—C15	1.487 (4)
Ni1—N6	2.093 (3)	C9—C10	1.480 (4)
Ni1—N3	2.096 (3)	C10—C11	1.376 (5)
Ni1—N1	2.098 (3)	C11—C12	1.378 (5)
Ni1—N4	2.099 (3)	C11—H11	0.9300
N1—C15	1.335 (4)	C12—C13	1.356 (6)
N1—C19	1.350 (4)	C12—H12	0.9300
N2—C8	1.331 (4)	C13—C14	1.377 (5)
N2—C9	1.339 (4)	C13—H13	0.9300
N3—C14	1.329 (4)	C14—H14	0.9300
N3—C10	1.333 (4)	C15—C16	1.373 (5)
N4—C20	1.322 (4)	C16—C17	1.364 (5)
N4—C24	1.332 (4)	C16—H16	0.9300
N5—C29	1.326 (4)	C17—C18	1.361 (5)
N5—C25	1.340 (4)	C17—H17	0.9300
N6—C34	1.329 (4)	C18—C19	1.368 (5)
N6—C30	1.332 (4)	C18—H18	0.9300
O1W—H1WA	0.8732	C19—H19	0.9300
O1W—H1WB	0.8701	C20—C21	1.370 (5)
O1—C4	1.336 (4)	C20—H20	0.9300
O1—C1	1.379 (6)	C21—C22	1.336 (6)
O2W—H2WB	0.8275	C21—H21	0.9300
O2W—H2WA	0.8393	C22—C23	1.379 (5)
O2—C35	1.364 (4)	C22—H22	0.9300
O2—C38	1.373 (5)	C23—C24	1.373 (4)
O2A—C35	1.404 (9)	C23—H23	0.9300
O2A—C38A	1.421 (9)	C24—C25	1.476 (5)
O3W—H3WC	0.8556	C25—C26	1.362 (5)
O3W—H3WA	0.8797	C26—C27	1.383 (5)
O4W—H4WB	0.8839	C26—H26	0.9300
O4W—H4WA	0.8729	C27—C28	1.389 (4)
O5W—H5WA	0.8692	C27—C35	1.435 (5)
O5W—H5WB	0.8890	C28—C29	1.374 (5)
O6W—H6WC	0.8315	C28—H28	0.9300
O6W—H6WA	0.8339	C29—C30	1.474 (4)
O7W—H7WA	0.8667	C30—C31	1.372 (5)
O7W—H7WB	0.8744	C31—C32	1.375 (5)
O8W—H8WC	0.8502	C31—H31	0.9300
O8W—H8WD	0.8528	C32—C33	1.372 (5)
O9W—H9WA	0.8616	C32—H32	0.9300
O9W—H9WB	0.8629	C33—C34	1.365 (5)
O10W—H10A	0.8785	C33—H33	0.9300
O10W—H10B	0.8705	C34—H34	0.9300
C1—C2	1.299 (7)	C35—C36	1.328 (6)
C1—H1A	0.9300	C35—C36A	1.352 (9)

C2—C3	1.394 (6)	C36—C37	1.378 (6)
C2—H2A	0.9300	C36—H36	0.9300
C3—C4	1.334 (5)	C36A—C37A	1.344 (10)
C3—H3A	0.9300	C36A—H36A	0.9300
C4—C5	1.432 (5)	C37—C38	1.315 (6)
C5—C6	1.394 (5)	C37—H37	0.9300
C5—C7	1.400 (5)	C37A—C38A	1.351 (9)
C6—C9	1.368 (5)	C37A—H37A	0.9300
C6—H6	0.9300	C38—H38	0.9300
C7—C8	1.355 (5)	C38A—H38A	0.9300
N5—Ni1—N2	178.36 (10)	N3—C14—H14	118.9
N5—Ni1—N6	77.81 (11)	C13—C14—H14	118.9
N2—Ni1—N6	102.65 (11)	N1—C15—C16	121.9 (3)
N5—Ni1—N3	100.71 (11)	N1—C15—C8	114.7 (3)
N2—Ni1—N3	77.74 (11)	C16—C15—C8	123.5 (3)
N6—Ni1—N3	89.84 (11)	C17—C16—C15	119.0 (4)
N5—Ni1—N1	103.80 (10)	C17—C16—H16	120.5
N2—Ni1—N1	77.77 (11)	C15—C16—H16	120.5
N6—Ni1—N1	93.13 (11)	C18—C17—C16	119.8 (3)
N3—Ni1—N1	155.38 (11)	C18—C17—H17	120.1
N5—Ni1—N4	78.10 (11)	C16—C17—H17	120.1
N2—Ni1—N4	101.46 (12)	C17—C18—C19	119.1 (3)
N6—Ni1—N4	155.89 (11)	C17—C18—H18	120.5
N3—Ni1—N4	95.46 (11)	C19—C18—H18	120.5
N1—Ni1—N4	91.74 (10)	N1—C19—C18	121.7 (4)
C15—N1—C19	118.5 (3)	N1—C19—H19	119.1
C15—N1—Ni1	114.5 (2)	C18—C19—H19	119.1
C19—N1—Ni1	126.9 (2)	N4—C20—C21	122.5 (4)
C8—N2—C9	120.0 (3)	N4—C20—H20	118.8
C8—N2—Ni1	120.1 (2)	C21—C20—H20	118.8
C9—N2—Ni1	119.7 (2)	C22—C21—C20	118.9 (4)
C14—N3—C10	118.6 (3)	C22—C21—H21	120.6
C14—N3—Ni1	126.4 (2)	C20—C21—H21	120.6
C10—N3—Ni1	114.8 (2)	C21—C22—C23	119.9 (4)
C20—N4—C24	118.8 (3)	C21—C22—H22	120.0
C20—N4—Ni1	127.0 (3)	C23—C22—H22	120.0
C24—N4—Ni1	114.1 (2)	C24—C23—C22	118.4 (4)
C29—N5—C25	120.9 (3)	C24—C23—H23	120.8
C29—N5—Ni1	119.5 (2)	C22—C23—H23	120.8
C25—N5—Ni1	119.6 (2)	N4—C24—C23	121.5 (3)
C34—N6—C30	118.6 (3)	N4—C24—C25	115.4 (3)
C34—N6—Ni1	126.8 (2)	C23—C24—C25	123.1 (3)
C30—N6—Ni1	114.5 (2)	N5—C25—C26	120.4 (3)
H1WA—O1W—H1WB	109.3	N5—C25—C24	112.8 (3)
C4—O1—C1	105.9 (4)	C26—C25—C24	126.8 (3)
H2WB—O2W—H2WA	108.4	C25—C26—C27	120.0 (3)
C35—O2—C38	106.7 (4)	C25—C26—H26	120.0

C35—O2A—C38A	104.3 (8)	C27—C26—H26	120.0
H3WC—O3W—H3WA	110.8	C26—C27—C28	118.7 (3)
H4WB—O4W—H4WA	89.3	C26—C27—C35	121.7 (3)
H5WA—O5W—H5WB	81.1	C28—C27—C35	119.7 (3)
H6WC—O6W—H6WA	96.0	C29—C28—C27	118.6 (3)
H7WA—O7W—H7WB	110.0	C29—C28—H28	120.7
H8WC—O8W—H8WD	111.3	C27—C28—H28	120.7
H9WA—O9W—H9WB	101.0	N5—C29—C28	121.4 (3)
H10A—O10W—H10B	87.7	N5—C29—C30	113.5 (3)
C2—C1—O1	109.2 (4)	C28—C29—C30	125.1 (3)
C2—C1—H1A	125.4	N6—C30—C31	122.3 (3)
O1—C1—H1A	125.4	N6—C30—C29	114.5 (3)
C1—C2—C3	108.6 (5)	C31—C30—C29	123.1 (3)
C1—C2—H2A	125.7	C30—C31—C32	118.4 (3)
C3—C2—H2A	125.7	C30—C31—H31	120.8
C4—C3—C2	105.5 (4)	C32—C31—H31	120.8
C4—C3—H3A	127.3	C33—C32—C31	119.4 (3)
C2—C3—H3A	127.3	C33—C32—H32	120.3
C3—C4—O1	110.8 (3)	C31—C32—H32	120.3
C3—C4—C5	129.7 (4)	C34—C33—C32	118.7 (3)
O1—C4—C5	119.5 (3)	C34—C33—H33	120.7
C6—C5—C7	118.2 (3)	C32—C33—H33	120.7
C6—C5—C4	121.2 (3)	N6—C34—C33	122.6 (3)
C7—C5—C4	120.6 (3)	N6—C34—H34	118.7
C9—C6—C5	118.8 (3)	C33—C34—H34	118.7
C9—C6—H6	120.6	C36—C35—O2	107.6 (4)
C5—C6—H6	120.6	C36A—C35—O2A	109.5 (8)
C8—C7—C5	119.4 (3)	C36—C35—C27	133.6 (4)
C8—C7—H7	120.3	C36A—C35—C27	133.4 (8)
C5—C7—H7	120.3	O2—C35—C27	118.8 (3)
N2—C8—C7	121.8 (3)	O2A—C35—C27	117.1 (7)
N2—C8—C15	112.6 (3)	C35—C36—C37	109.4 (4)
C7—C8—C15	125.6 (3)	C35—C36—H36	125.3
N2—C9—C6	121.7 (3)	C37—C36—H36	125.3
N2—C9—C10	112.7 (3)	C37A—C36A—C35	108.2 (9)
C6—C9—C10	125.5 (3)	C37A—C36A—H36A	125.9
N3—C10—C11	122.2 (3)	C35—C36A—H36A	125.9
N3—C10—C9	114.7 (3)	C38—C37—C36	106.7 (4)
C11—C10—C9	122.9 (3)	C38—C37—H37	126.7
C10—C11—C12	118.4 (4)	C36—C37—H37	126.7
C10—C11—H11	120.8	C36A—C37A—C38A	110.1 (10)
C12—C11—H11	120.8	C36A—C37A—H37A	124.9
C13—C12—C11	119.5 (3)	C38A—C37A—H37A	124.9
C13—C12—H12	120.3	C37—C38—O2	109.6 (4)
C11—C12—H12	120.3	C37—C38—H38	125.2
C12—C13—C14	119.1 (4)	O2—C38—H38	125.2
C12—C13—H13	120.4	C37A—C38A—O2A	107.8 (9)
C14—C13—H13	120.4	C37A—C38A—H38A	126.1

N3—C14—C13

122.1 (4)

O2A—C38A—H38A

126.1

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1 <i>W</i> —H1 <i>WA</i> ···C11	0.87	2.25	3.113 (4)	169
O1 <i>W</i> —H1 <i>WB</i> ···O9 <i>W</i> ⁱ	0.87	2.06	2.923 (6)	175
O2 <i>W</i> —H2 <i>WB</i> ···O5 <i>W</i> ⁱⁱ	0.83	1.99	2.813 (7)	172
O2 <i>W</i> —H2 <i>WA</i> ···C11	0.84	2.39	3.215 (4)	168
O3 <i>W</i> —H3 <i>WC</i> ···O4 <i>W</i>	0.86	2.05	2.760 (9)	140
O3 <i>W</i> —H3 <i>WA</i> ···O6 <i>W</i> ⁱⁱⁱ	0.88	2.35	3.134 (7)	148
O4 <i>W</i> —H4 <i>WB</i> ···C12	0.88	2.58	3.107 (5)	119
O4 <i>W</i> —H4 <i>WA</i> ···C12	0.87	2.56	3.107 (5)	122
O5 <i>W</i> —H5 <i>WA</i> ···C12	0.87	2.37	3.079 (4)	138
O5 <i>W</i> —H5 <i>WB</i> ···O9 <i>W</i>	0.89	2.16	2.991 (6)	156
O6 <i>W</i> —H6 <i>WC</i> ···O2 <i>W</i> ⁱⁱ	0.83	2.11	2.929 (6)	167
O6 <i>W</i> —H6 <i>WA</i> ···O7 <i>W</i>	0.83	2.18	2.838 (6)	136
O7 <i>W</i> —H7 <i>WA</i> ···C12	0.87	2.34	3.190 (4)	167
O7 <i>W</i> —H7 <i>WB</i> ···O4 <i>W</i> ⁱⁱ	0.87	1.93	2.798 (5)	172
O8 <i>W</i> —H8 <i>WC</i> ···O3 <i>W</i> ⁱⁱ	0.85	2.06	2.856 (8)	155
O8 <i>W</i> —H8 <i>WD</i> ···C12 ^{iv}	0.85	2.40	3.204 (6)	157
O9 <i>W</i> —H9 <i>WA</i> ···O10 <i>W</i> ^v	0.86	1.93	2.756 (6)	159
O9 <i>W</i> —H9 <i>WB</i> ···O1 <i>W</i> ^{vi}	0.86	2.11	2.878 (5)	147
O10 <i>W</i> —H10 <i>A</i> ···C11 ^{vii}	0.88	2.27	3.141 (4)	171
O10 <i>W</i> —H10 <i>B</i> ···C11 ^{viii}	0.87	2.38	3.225 (4)	165

Symmetry codes: (i) $-x, -y+1, -z+1$; (ii) $-x+1, -y+1, -z+1$; (iii) $x-1, y, z$; (iv) $x+1, y, z$; (v) $-x+1, -y+1, -z$; (vi) $x, y+1, z-1$; (vii) $x, y, z-1$; (viii) $-x+1, -y, -z+1$.