



Aquachlorido(2-[[6-(dimethylamino)pyrimidin-4-yl]sulfanyl]pyrimidine-4,6-diamine)copper(II) chloride hydrate

Tristen E. Moyaert,^a Christina Paul,^a Weibin Chen,^a Amy A. Sarjeant^b and Louise N. Dawe^{a*}

Received 11 September 2017
Accepted 19 September 2017

Edited by M. Zeller, Purdue University, USA

Keywords: crystal structure; copper(II); non-symmetric sulfanyl ligand; disorder.

CCDC reference: 1575371



Supporting information: this article has supporting information at journals.iucr.org/e

^aDepartment of Chemistry and Biochemistry, Faculty of Science, Wilfrid Laurier University, 75 University Ave. W., Waterloo, ON, N2L 3C5, Canada, and ^bCambridge Crystallographic Data Centre, Center for Integrative Proteomics Research, 174 Frelinghuysen Road, Piscataway, NJ 08854 USA. *Correspondence e-mail: ldawe@wlu.ca

A copper(II) complex of the non-symmetric bidentate ligand 2-[[6-(dimethylamino)pyrimidin-4-yl]sulfanyl]pyrimidine-4,6-diamine (*L1*) is reported. The single-crystal X-ray structure of aqua[aqua/chlorido(0.49/0.51)](2-[[6-(dimethylamino)pyrimidin-4-yl]sulfanyl]pyrimidine-4,6-diamine)copper(II) 0.49-chloride 1.51-hydrate, $[\text{CuCl}_{1.51}(\text{C}_{10}\text{H}_{13}\text{N}_7\text{S})(\text{H}_2\text{O})_{1.49}]\text{Cl}_{0.49}\cdot 1.51\text{H}_2\text{O}$ or $[(L1)\text{Cl}_{1.51}(\text{H}_2\text{O})_{1.49}\text{Cu}]0.49\text{Cl}\cdot 1.51\text{H}_2\text{O}$, exhibits distorted square-pyramidal geometry around the metal centre, with disorder in the axial position, occupied by chloride or water. The six-membered metal–chelate ring is in a boat conformation, and short intermolecular S··S interactions are observed. In addition to its capacity for bidentate metal coordination, the ligand has the ability to engage in further supramolecular interactions as both a hydrogen-bond donor and acceptor, and multiple interactions with lattice solvent water molecules are present in the reported structure.

1. Chemical context

Non-symmetric ligand–metal complexes have been explored for their applications in chiral synthesis (Asay & Morales-Morales, 2015; Pfaltz & Drury, 2004), or for their potential to yield new multimetallic topologies which combine homo- and heteroleptic sites into a single molecule (Dawe *et al.*, 2006). Non-symmetric thio-bis-(pyridin-2-yl) or bis-(pyrimidin-2-yl) ligands are known, and upon bidentate coordination with transition metal cations, these form six-membered chelate rings, which adopt a boat-shaped conformation (Fig. 1). Some reported transition metal complexes resulting from this class of ligands have been employed as possible alternatives to traditional chemotherapy drugs (Ray *et al.*, 1994; Mandal *et al.*, 2007), as a step en route to new thrombin inhibitors (Chung *et al.*, 2003), and have led to the formation of one Cu^I 30-nuclear cluster (Li *et al.*, 2012).






Figure 1
Boat-shaped chelate ring for bidentate coordination of thio-bis-(pyridin-2-yl) or bis-(pyrimidin-2-yl) ligands.

In the interest of exploring simultaneous coordination chemistry and anion–ligand affinity *via* hydrogen-bonding interactions, the non-symmetric ligand 2-[[6-(dimethylamino)pyrimidin-4-yl]sulfanyl]pyrimidine-4,6-diamine ($C_{10}H_{13}N_7S$; *L1*), was synthesized, and its metal complex with copper(II) chloride, is reported here. Even upon metal coordination, the ligand can still serve as a hydrogen-bond donor to anions *via* the amine moieties. Alternatively, these free amines could also act as possible anchors for surface attachment, with a view towards future device applications.

2. Structural commentary

The title compound crystallizes in the monoclinic space group $P2_1/c$ with one bidentate ligand bound to a copper(II) cation (*via* N1 and N4; Fig. 2). The copper(II) cation is five-coordinate, with the remaining coordination sites occupied by a chloride anion (Cl1), a water molecule (O1), and a disordered site, with either chloride (Cl2; Fig. 2*a*) or water (O2; Fig. 2*b*)

Table 1
Selected geometric parameters (\AA , $^\circ$).

Cu1—Cl1	2.2689 (7)	Cu1—N4	1.996 (2)
Cu1—Cl2	2.5273 (19)	S1—C4	1.777 (2)
Cu1—O1	2.0158 (19)	S1—C5	1.774 (3)
Cu1—O2	2.229 (6)	N1—C4	1.332 (3)
Cu1—N1	2.034 (2)	N4—C5	1.353 (3)
Cl1—Cu1—Cl2	92.89 (5)	N1—Cu1—O2	101.55 (18)
O1—Cu1—Cl1	91.94 (6)	N4—Cu1—Cl1	171.10 (6)
O1—Cu1—Cl2	92.69 (7)	N4—Cu1—Cl2	95.20 (8)
O1—Cu1—O2	100.64 (18)	N4—Cu1—O1	83.98 (8)
O1—Cu1—N1	157.01 (8)	N4—Cu1—O2	95.5 (2)
O2—Cu1—Cl1	93.1 (2)	N4—Cu1—N1	87.99 (8)
N1—Cu1—Cl1	92.81 (6)	C5—S1—C4	104.53 (12)
N1—Cu1—Cl2	109.51 (8)		

with occupancies of 0.511 (5) and 0.489 (5), respectively. The two largest ligand–metal–ligand bond angles (Table 1) are N4—Cu1—Cl1 and O1—Cu1—N1 [171.10 (6) and 157.01 (8) $^\circ$, respectively] giving a τ value of 0.23 (where $\tau = 0$ is ideal square-pyramidal geometry, and $\tau = 1$ is ideal trigonal-bipyramidal geometry; Addison *et al.*, 1984), indicating that the geometry is distorted square pyramidal. Examination of the bond lengths (Table 1), is also consistent with the disordered Cl/O as the axial site for this geometry. An intramolecular hydrogen bond is present between the amine group (*via* N5—H5A) and the apical ligand (Fig. 2; Table 2). The six-membered chelate ring adopts a boat conformation. The angle between the distorted square plane defined by N1/N4/C4/C5 (r.m.s. deviation from the plane is 0.032 \AA) and the flap defined by C4/S1/C5 (θ_1) is 34.51 (17) $^\circ$, while the angle between the square plane and the flap defined by N1/Cu1/N4 (θ_2) is 46.93 (14) $^\circ$. The boat-shaped configuration accommodates the C—S and N—Cu bonds, making up the flaps,




Figure 2
Asymmetric unit for $[(C_{10}H_{13}N_7S)Cl_{1.51}(H_2O)_{1.49}Cu]0.49Cl \cdot 1.51H_2O$, with 50% displacement ellipsoids. (a) Disordered atoms with 0.51-occupancy; (b) disordered atoms with 0.49-occupancy. All atoms in (a) and (b) are identical, except those labelled in (b). Hydrogen bonds are represented by dashed lines.

Table 2
 Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1–H1A \cdots N2 ⁱ	0.92 (1)	1.99 (2)	2.897 (3)	170 (3)
O1–H1B \cdots O4 ⁱⁱ	0.91 (2)	1.82 (6)	2.71 (6)	167 (4)
O1–H1B \cdots O5 ⁱⁱ	0.91 (2)	1.80 (6)	2.69 (6)	162 (4)
O3–H3A \cdots Cl2	0.92	2.28	3.200 (7)	176
O3–H3B \cdots O4	0.93	2.17	2.94 (6)	139
O4–H4A \cdots O3 ⁱⁱ	0.91	2.25	3.07 (6)	150
O2–H2B \cdots Cl3	0.91	2.33	3.177 (7)	154
O5–H5C \cdots Cl3	0.93	2.02	2.83 (6)	145
N5–H5A \cdots Cl2	0.84 (3)	2.46 (3)	3.295 (3)	172 (3)
N5–H5A \cdots O2	0.84 (3)	2.07 (3)	2.903 (7)	167 (3)
N5–H5B \cdots Cl1 ⁱⁱⁱ	0.79 (3)	2.56 (4)	3.353 (3)	173 (3)
N6–H6A \cdots Cl3 ⁱⁱⁱ	0.80 (4)	2.41 (4)	3.187 (4)	165 (3)
N6–H6A \cdots O3 ⁱⁱⁱ	0.80 (4)	2.32 (4)	3.094 (8)	163 (3)

Symmetry codes: (i) $-x+1, y-\frac{1}{2}, -z+\frac{3}{2}$; (ii) $-x+2, -y+1, -z+1$; (iii) $-x+1, -y+2, -z+1$.

which are significantly longer than the C–N bonds in the square plane (Table 1.)

A simpler, symmetric bidentate ligand, di(pyridin-2-yl)sulfide (DPS), has been reported to exhibit a very similar metal coordination environment to the major component reported here, upon reaction with $\text{CuCl}_2\cdot\text{H}_2\text{O}$, to yield $[\text{Cu}(\text{DPS})(\text{H}_2\text{O})\text{Cl}_2]\cdot\text{H}_2\text{O}$ (Teles *et al.*, 2006). In this complex, the authors report $\tau = 0.06$, with the square plane formed by the two nitrogen atoms from DPS, a coordinating water molecule, and one chloride ion (with the second chloride occupying the axial position). Similar to the reported structure here, the six-membered chelate ring adopts a boat conformation, which is characteristic for transition metal complexes with this class of ligands upon bidentate coordination (*vide infra*).




Figure 3
 Packed unit cell for $[(\text{L1})\text{Cl}_{1.51}(\text{H}_2\text{O})_{1.49}\text{Cu}]_{0.49}\text{Cl}_{1.51}\text{H}_2\text{O}$. Only atoms in the major occupancy component are shown. All solvent water molecules and hydrogen atoms have been omitted for clarity.

3. Supramolecular features


In the crystal, molecules of the title complex pack in columns, parallel to the crystallographic b axis (Fig. 3), with short $\text{S}\cdots\text{S}^i$ intermolecular distances [3.7327 (3) Å; symmetry code: (i) $-x+1, y+\frac{1}{2}, -z+\frac{3}{2}$]. Note that each chelated ‘boat’ points in the same direction within a column, and the opposite direction is observed in adjacent columns.

4. Database survey

A survey was performed of the Cambridge Structural Database (version 5.38 with May 2017 updates; Groom *et al.*, 2016), using *ConQuest* (version 1.19; Bruno *et al.*, 2002), for six-membered transition metal chelate rings resulting from bidentate ligand coordination, where the metal was any transition metal, and the other ring components were N–C–S–C–N. Further, within the ligand, each C–N was required to be part of a six-membered ring, where the remaining four atoms could be any non-metal, and the bond type within the ring was unspecified (allowed to be ‘any’ bond type). This resulted in 74 hits, which were then manually sorted to omit systems where the ligand exhibited anything greater than bidenticity, leaving 68 structures for further analysis using *Mercury* (version 3.9; Macrae *et al.*, 2006). All of these exhibited boat-shaped puckering of the chelate ring, with mean values for $\theta_1 = 43$ (7) and $\theta_2 = 37$ (5)°. While the larger angle for the title complex is θ_2 , both θ_1 and θ_2 are within two standard deviations of comparable structures from the database.

5. Synthesis and crystallization

2-[[6-(Dimethylamino)pyrimidin-4-yl]sulfanyl]pyrimidine-4,6-diamine (C₁₀H₁₃N₇S; L1): 0.972 g (7.03 mmol) of potassium carbonate and 1.000 g (6.24 mmol) of 4,6-diamino-2-mercaptopyrimidine hydrate were combined in 20 mL of dimethylformamide, and stirred at 333 K for 20 min, prior to the addition of 0.524 g (3.51 mmol) of 4,6-dichloropyrimidine (see reaction scheme). The resulting cloudy orange solution was refluxed for 24 h. It was then filtered, and the brown filtrate was reduced *in vacuo* to yield 0.387 g (1.47 mmol) of orange solid, after washing with ethanol (42% yield).



Aquachlorido(2-[[6-(dimethylamino)pyrimidin-4-yl]sulfanyl]pyrimidine-4,6-diamine)copper(II) chloride hydrate $[\text{CuCl}_{1.51}(\text{C}_{10}\text{H}_{13}\text{N}_7\text{S})(\text{H}_2\text{O})_{1.49}]\text{Cl}_{0.49}\cdot 1.51\text{H}_2\text{O}$: 0.050 g (0.19 mmol) of L1 and 0.048 g (0.28 mmol) of $\text{CuCl}_2\cdot 2\text{H}_2\text{O}$ were separately dissolved in 5 mL of 1:1 methanol/acetonitrile. The solution of CuCl_2 was added dropwise to the solution of L1. The resulting cloudy brown solution was stirred vigorously with heating (333 K) for 20 min. This was filtered, yielding 0.007 g of brown, amorphous powder, and a clear green filtrate

that was left for slow evaporation. Green, prismatic X-ray quality crystals grew from the filtrate over the course of six weeks. 3.6 mg (0.0080 mmol) of analytically pure crystals were harvested as soon as they formed, though the mother liquor was still highly coloured, accounting for the low (4.2%) yield. These crystals were analyzed *via* small molecule X-ray diffraction, and elemental analysis. Analysis calculated for $[(C_{10}H_{13}N_7S)CuCl_2 \cdot 3H_2O]$: C, 26.58; H, 4.24; N, 21.7. Found: C, 26.26; H, 4.13; N, 21.31. Presence of copper confirmed *via* graphite furnace atomic absorption spectroscopy: calculated: $25 \mu\text{g L}^{-1}$; found: $31.0 \pm 0.14 \mu\text{g L}^{-1}$ ($n = 8$).

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. Hydrogen atoms were introduced in calculated positions and refined using a riding model, except those bonded to oxygen or nitrogen atoms, which were introduced in difference-map positions. N–H hydrogen atoms were refined isotropically, with no restraints. All O–H hydrogen atoms (all associated with water molecules) were refined with $U_{\text{iso}}(\text{H})$ 1.5 times that of the parent atoms and rotating geometry constraints (AFIX 7). Similar distance restraints (SADI, esd 0.02) were applied for all water molecules.

The structure exhibited significant disorder. This included main fragment disorder in the coordination sphere around Cu1. As such, similar distance restraints (SADI, esd 0.02) were applied to the Cu–OH₂ and Cu–Cl bonds; for each, one O atom (O1) and one Cl atom (Cl1) were fully occupied, while the other (O2 and Cl2) were at partial occupancy, occupying the same coordination site on Cu1, with a sum of their occupancy equal to one. Identical anisotropic displacement parameter (EADP) constraints were applied to Cl2 and O2. Finally, EADP constraints were also applied to a disordered water molecule (O4 and O5), with a sum occupancy of one.

While the structure does exhibit significant disorder, careful consideration was given to ensure that: (i) charge balance was established; (ii) the model was consistent with a reasonable hydrogen-bonding network; and (iii) the next highest residual electron density peak was associated along a C–S bond.

Acknowledgements

Dr Paul Boyle, Department of Chemistry, X-Ray Facility, University of Western Ontario, Canada, is sincerely thanked for X-ray data collections. Dr Ken Maly, Department of Chemistry and Biochemistry, Wilfrid Laurier University, is thanked for helpful discussions on ligand synthesis. Dr Scott Smith, Department of Chemistry and Biochemistry, Wilfrid Laurier University, is thanked for helpful discussions on metal confirmation *via* GFAA spectroscopy.

Funding information

Funding for this research was provided by: Natural Sciences and Engineering Research Council of Canada, Canada

Table 3
Experimental details.

Crystal data	
Chemical formula	$[CuCl_{1.51}(C_{10}H_{13}N_7S)(H_2O)_{1.49}] \cdot Cl_{0.49} \cdot 1.51H_2O$
M_r	451.82
Crystal system, space group	Monoclinic, $P2_1/c$
Temperature (K)	110
a, b, c (Å)	11.42069 (19), 7.23911 (12), 21.6990 (3)
β (°)	103.1543 (16)
V (Å ³)	1746.91 (5)
Z	4
Radiation type	Cu $K\alpha$
μ (mm ⁻¹)	5.94
Crystal size (mm)	0.27 × 0.19 × 0.17
Data collection	
Diffractometer	Bruker APEXII CCD with <i>CrysAlis PRO</i> imported SAXI images
Absorption correction	Multi-scan (<i>CrysAlis PRO</i> ; Rigaku Oxford Diffraction, 2015)
$T_{\text{min}}, T_{\text{max}}$	0.034, 0.115
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	22331, 3000, 2523
R_{int}	0.051
$(\sin \theta/\lambda)_{\text{max}}$ (Å ⁻¹)	0.594
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.030, 0.079, 1.04
No. of reflections	3000
No. of parameters	261
No. of restraints	47
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ⁻³)	0.32, -0.32

Computer programs: *APEX2* (Bruker, 2012), *CrysAlis PRO* (Rigaku OD, 2015), *SHELXT* (Sheldrick, 2015a), *SHELXL2016* (Sheldrick, 2015b) and *OLEX2* (Dolomanov *et al.*, 2009).

Foundation for Innovation, Wilfrid Laurier University Research Office, The Research Support Fund (Wilfrid Laurier University), Laurier Faculty of Science Student Association (award to T. Moyaert).

References

- Addison, A. W., Rao, T. N., Reedijk, J., van Rijn, J. & Verschoor, G. C. (1984). *J. Chem. Soc. Dalton Trans.* pp. 1349–1356.
- Asay, M. & Morales-Morales, D. (2015). *Dalton Trans.* **44**, 17432–17447.
- Bruker (2012). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruno, I. J., Cole, J. C., Edgington, P. R., Kessler, M., Macrae, C. F., McCabe, P., Pearson, J. & Taylor, R. (2002). *Acta Cryst.* **B58**, 389–397.
- Chung, J. Y. L., Cvetovich, R. J., Tsay, F.-R., Dormer, P. G., DiMichele, L., Mathre, D. J., Chilenski, J. R., Mao, B. & Wenslow, R. (2003). *J. Org. Chem.* **68**, 8838–8846.
- Dawe, L., Abedin, T., Kelly, T., Thompson, L., Miller, D., Zhao, L., Wilson, C., Leech, M. & Howard, J. (2006). *J. Mater. Chem.* **16**, 2645–2659.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.

- Li, H.-X., Zhao, W., Li, H.-Y., Xu, Z.-L., Wang, W.-X. & Lang, J.-P. (2012). *Chem. Commun.* pp. 4259–4261.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Mandal, S., Bérubé, G., Asselin, É., Richardson, V. J., Church, J. G., Bridson, J., Pham, T. N. Q., Pramanik, S. K. & Mandal, S. K. (2007). *Bioorg. Med. Chem. Lett.* **17**, 2139–2145.
- Pfaltz, A. & Drury, W. J. (2004). *Proc. Natl Acad. Sci. USA*, **101**, 5723–5726.
- Ray, S., Smith, F. R., Bridson, J. N., Hong, Q., Richardson, V. J. & Mandal, S. K. (1994). *Inorg. Chim. Acta*, **227**, 175–179.
- Rigaku OD (2015). *CrysAlis PRO*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **C71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **A71**, 3–8.
- Teles, W. M., Marinho, M. V., Yoshida, M. I., Speziali, N. L., Krambrock, K., Pinheiro, C. B., Pinhal, N. M., Leitão, A. A. & Machado, F. C. (2006). *Inorg. Chim. Acta*, **359**, 4613–4618.

supporting information

Acta Cryst. (2017). E73, 1534-1538 [https://doi.org/10.1107/S205698901701338X]

Aquachlorido(2-[[6-(dimethylamino)pyrimidin-4-yl]sulfanyl]pyrimidine-4,6-diamine)copper(II) chloride hydrate

Tristen E. Moyaert, Christina Paul, Weibin Chen, Amy A. Sarjeant and Louise N. Dawe

Computing details

Data collection: *APEX2* (Bruker, 2012); cell refinement: *CrysAlis PRO* (Rigaku OD, 2015); data reduction: *CrysAlis PRO* (Rigaku OD, 2015); program(s) used to solve structure: *SHELXT* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2016* (Sheldrick, 2015b); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2* (Dolomanov *et al.*, 2009).

Aqua[aqua/chlorido(0.49/0.51)](2-[[6-(dimethylamino)pyrimidin-4-yl]sulfanyl]pyrimidine-4,6-diamine)copper(II) 0.49-chloride 1.51-hydrate

Crystal data

[CuCl_{1.51}(C₁₀H₁₃N₇S)(H₂O)_{1.49}].0.49Cl·1.51H₂O
M_r = 451.82
 Monoclinic, *P2₁/c*
a = 11.42069 (19) Å
b = 7.23911 (12) Å
c = 21.6990 (3) Å
 β = 103.1543 (16)°
V = 1746.91 (5) Å³
Z = 4

F(000) = 924
D_x = 1.718 Mg m⁻³
 Cu *K*α radiation, λ = 1.54178 Å
 Cell parameters from 13698 reflections
 θ = 4.0–66.3°
 μ = 5.94 mm⁻¹
T = 110 K
 Prism, green
 0.27 × 0.19 × 0.17 mm

Data collection

Bruker APEXII CCD with *CrysAlis PRO*
 imported SAXI images
 diffractometer
 Radiation source: fine-focus sealed X-ray tube,
 Enhance (Cu) X-ray Source
 Graphite monochromator
 ω and φ scans
 Absorption correction: multi-scan
 (*CrysAlis PRO*; Rigaku Oxford Diffraction,
 2015)

T_{min} = 0.034, *T_{max}* = 0.115
 22331 measured reflections
 3000 independent reflections
 2523 reflections with *I* > 2σ(*I*)
R_{int} = 0.051
 θ_{\max} = 66.4°, θ_{\min} = 4.0°
h = -13→13
k = -8→8
l = -25→22

Refinement

Refinement on *F*²
 Least-squares matrix: full
R[*F*² > 2σ(*F*²)] = 0.030
wR(*F*²) = 0.079
S = 1.04
 3000 reflections

261 parameters
 47 restraints
 Hydrogen site location: mixed
 H atoms treated by a mixture of independent
 and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0438P)^2 + 0.8086P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$

$$\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.32 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cu1	0.65886 (3)	0.87641 (5)	0.36726 (2)	0.02056 (12)	
Cl1	0.59471 (5)	0.68609 (9)	0.43587 (3)	0.02412 (16)	
Cl2	0.81961 (17)	1.0193 (3)	0.45364 (9)	0.0250 (6)	0.511 (5)
Cl3	0.9316 (3)	0.7674 (6)	0.55061 (13)	0.0648 (12)	0.489 (5)
S1	0.47778 (6)	0.94220 (9)	0.22968 (3)	0.02244 (16)	
O1	0.77219 (16)	0.6829 (3)	0.34783 (8)	0.0256 (4)	
H1A	0.743 (3)	0.635 (4)	0.3084 (9)	0.051 (11)*	
H1B	0.790 (3)	0.586 (4)	0.3750 (15)	0.062 (12)*	
O3	0.9002 (6)	0.6924 (9)	0.5536 (4)	0.0305 (15)	0.511 (5)
H3A	0.873556	0.784224	0.524170	0.046*	0.511 (5)
H3B	0.982596	0.704284	0.569168	0.046*	0.511 (5)
O4	1.148 (5)	0.571 (8)	0.560 (3)	0.056 (3)	0.511 (5)
H4A	1.164621	0.489914	0.531128	0.084*	0.511 (5)
H4B	1.218311	0.634714	0.578705	0.084*	0.511 (5)
O2	0.7814 (6)	1.0316 (10)	0.4446 (3)	0.0250 (6)	0.489 (5)
H2A	0.816574	1.128467	0.429387	0.038*	0.489 (5)
H2B	0.842833	0.962117	0.467637	0.038*	0.489 (5)
O5	1.141 (5)	0.553 (9)	0.557 (3)	0.056 (3)	0.489 (5)
H5C	1.094042	0.658743	0.549583	0.084*	0.489 (5)
H5D	1.164982	0.487073	0.526489	0.084*	0.489 (5)
N1	0.50737 (19)	1.0333 (3)	0.35292 (9)	0.0214 (5)	
N2	0.31504 (19)	1.0711 (3)	0.28275 (10)	0.0229 (5)	
N3	0.86159 (19)	1.1243 (3)	0.25035 (10)	0.0238 (5)	
N4	0.70468 (19)	1.0147 (3)	0.29669 (9)	0.0213 (5)	
N5	0.5433 (2)	1.1471 (4)	0.45509 (11)	0.0286 (5)	
H5A	0.614 (3)	1.106 (5)	0.4577 (15)	0.034*	
H5B	0.517 (3)	1.185 (5)	0.4834 (16)	0.034*	
N6	0.1509 (2)	1.1695 (4)	0.32017 (14)	0.0321 (6)	
H6A	0.122 (3)	1.202 (5)	0.3488 (17)	0.033 (9)*	
H6B	0.109 (3)	1.134 (5)	0.2867 (17)	0.037 (10)*	
N7	0.8262 (2)	1.1817 (3)	0.14302 (10)	0.0266 (5)	
C1	0.4646 (2)	1.1146 (4)	0.40076 (12)	0.0232 (6)	
C2	0.3434 (2)	1.1595 (4)	0.39165 (12)	0.0250 (6)	
H2	0.312207	1.208633	0.425222	0.030*	
C3	0.2696 (2)	1.1307 (4)	0.33244 (13)	0.0246 (6)	
C4	0.4299 (2)	1.0264 (3)	0.29687 (11)	0.0204 (5)	

C5	0.6270 (2)	1.0255 (3)	0.23952 (11)	0.0205 (5)
C6	0.6605 (2)	1.0850 (3)	0.18649 (11)	0.0203 (5)
H6	0.604065	1.095192	0.147050	0.024*
C7	0.7828 (2)	1.1307 (4)	0.19266 (12)	0.0229 (6)
C8	0.8173 (2)	1.0684 (4)	0.29833 (12)	0.0235 (6)
H8	0.871669	1.066177	0.338604	0.028*
C9	0.7483 (3)	1.1897 (4)	0.07954 (13)	0.0342 (7)
H9A	0.729605	1.063940	0.063603	0.051*
H9B	0.789517	1.256260	0.051290	0.051*
H9C	0.673663	1.254206	0.081154	0.051*
C10	0.9543 (3)	1.2111 (5)	0.14952 (14)	0.0360 (7)
H10A	0.984405	1.295187	0.184969	0.054*
H10B	0.968239	1.265239	0.110432	0.054*
H10C	0.996500	1.092607	0.157550	0.054*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0206 (2)	0.0275 (2)	0.0147 (2)	0.00178 (16)	0.00640 (14)	0.00253 (14)
Cl1	0.0240 (3)	0.0325 (3)	0.0171 (3)	0.0027 (3)	0.0072 (2)	0.0045 (2)
Cl2	0.0180 (12)	0.0367 (7)	0.0183 (8)	0.0039 (8)	-0.0002 (7)	0.0004 (5)
Cl3	0.055 (2)	0.110 (3)	0.0285 (11)	0.0397 (19)	0.0078 (11)	-0.0013 (16)
S1	0.0219 (3)	0.0293 (3)	0.0168 (3)	-0.0034 (3)	0.0060 (2)	-0.0018 (2)
O1	0.0215 (10)	0.0351 (11)	0.0208 (10)	0.0028 (8)	0.0059 (8)	0.0012 (8)
O3	0.021 (3)	0.038 (3)	0.035 (3)	0.007 (2)	0.011 (2)	0.012 (2)
O4	0.080 (5)	0.054 (8)	0.034 (4)	0.022 (4)	0.013 (3)	0.009 (4)
O2	0.0180 (12)	0.0367 (7)	0.0183 (8)	0.0039 (8)	-0.0002 (7)	0.0004 (5)
O5	0.080 (5)	0.054 (8)	0.034 (4)	0.022 (4)	0.013 (3)	0.009 (4)
N1	0.0239 (11)	0.0268 (12)	0.0149 (10)	0.0019 (9)	0.0075 (9)	0.0032 (9)
N2	0.0216 (11)	0.0266 (12)	0.0209 (11)	-0.0009 (9)	0.0059 (9)	0.0051 (9)
N3	0.0212 (11)	0.0319 (12)	0.0188 (11)	-0.0002 (10)	0.0055 (9)	0.0019 (9)
N4	0.0212 (11)	0.0276 (12)	0.0156 (10)	0.0006 (9)	0.0051 (8)	-0.0001 (8)
N5	0.0288 (13)	0.0421 (15)	0.0162 (11)	0.0084 (11)	0.0080 (10)	-0.0024 (10)
N6	0.0233 (13)	0.0439 (16)	0.0306 (15)	0.0030 (11)	0.0092 (12)	0.0048 (12)
N7	0.0251 (12)	0.0394 (14)	0.0175 (11)	-0.0014 (10)	0.0092 (9)	0.0033 (9)
C1	0.0276 (14)	0.0244 (14)	0.0199 (13)	0.0024 (11)	0.0100 (11)	0.0068 (10)
C2	0.0285 (15)	0.0282 (14)	0.0213 (13)	0.0046 (11)	0.0117 (11)	0.0054 (10)
C3	0.0253 (14)	0.0234 (14)	0.0277 (14)	0.0016 (11)	0.0113 (11)	0.0101 (11)
C4	0.0236 (14)	0.0200 (13)	0.0191 (12)	-0.0006 (11)	0.0083 (10)	0.0044 (10)
C5	0.0222 (13)	0.0215 (13)	0.0189 (12)	0.0023 (10)	0.0068 (10)	-0.0024 (10)
C6	0.0222 (13)	0.0251 (14)	0.0144 (12)	0.0013 (11)	0.0057 (10)	0.0001 (10)
C7	0.0275 (14)	0.0234 (13)	0.0196 (13)	0.0005 (11)	0.0091 (11)	-0.0029 (10)
C8	0.0238 (14)	0.0292 (14)	0.0174 (12)	0.0023 (11)	0.0047 (10)	0.0013 (10)
C9	0.0341 (16)	0.0502 (18)	0.0193 (14)	-0.0019 (14)	0.0079 (12)	0.0053 (12)
C10	0.0254 (15)	0.057 (2)	0.0291 (15)	-0.0059 (14)	0.0138 (12)	0.0002 (14)

Geometric parameters (Å, °)

Cu1—C11	2.2689 (7)	N4—C5	1.353 (3)
Cu1—C12	2.5273 (19)	N4—C8	1.336 (3)
Cu1—O1	2.0158 (19)	N5—H5A	0.84 (3)
Cu1—O2	2.229 (6)	N5—H5B	0.79 (3)
Cu1—N1	2.034 (2)	N5—C1	1.331 (4)
Cu1—N4	1.996 (2)	N6—H6A	0.80 (4)
S1—C4	1.777 (2)	N6—H6B	0.81 (4)
S1—C5	1.774 (3)	N6—C3	1.351 (4)
O1—H1A	0.915 (14)	N7—C7	1.337 (3)
O1—H1B	0.911 (15)	N7—C9	1.461 (3)
O3—H3A	0.9233	N7—C10	1.452 (3)
O3—H3B	0.9290	C1—C2	1.392 (4)
O4—H4A	0.9144	C2—H2	0.9500
O4—H4B	0.9323	C2—C3	1.383 (4)
O2—H2A	0.9063	C5—C6	1.363 (4)
O2—H2B	0.9141	C6—H6	0.9500
O5—H5C	0.9293	C6—C7	1.411 (4)
O5—H5D	0.9107	C8—H8	0.9500
N1—C1	1.376 (3)	C9—H9A	0.9800
N1—C4	1.332 (3)	C9—H9B	0.9800
N2—C3	1.368 (4)	C9—H9C	0.9800
N2—C4	1.318 (3)	C10—H10A	0.9800
N3—C7	1.366 (3)	C10—H10B	0.9800
N3—C8	1.320 (3)	C10—H10C	0.9800
C11—Cu1—C12	92.89 (5)	C7—N7—C10	120.8 (2)
O1—Cu1—C11	91.94 (6)	C10—N7—C9	118.0 (2)
O1—Cu1—C12	92.69 (7)	N1—C1—C2	120.3 (2)
O1—Cu1—O2	100.64 (18)	N5—C1—N1	117.4 (2)
O1—Cu1—N1	157.01 (8)	N5—C1—C2	122.3 (2)
O2—Cu1—C11	93.1 (2)	C1—C2—H2	120.9
N1—Cu1—C11	92.81 (6)	C3—C2—C1	118.2 (2)
N1—Cu1—C12	109.51 (8)	C3—C2—H2	120.9
N1—Cu1—O2	101.55 (18)	N2—C3—C2	121.3 (2)
N4—Cu1—C11	171.10 (6)	N6—C3—N2	117.0 (3)
N4—Cu1—C12	95.20 (8)	N6—C3—C2	121.6 (3)
N4—Cu1—O1	83.98 (8)	N1—C4—S1	119.87 (19)
N4—Cu1—O2	95.5 (2)	N2—C4—S1	111.66 (18)
N4—Cu1—N1	87.99 (8)	N2—C4—N1	128.4 (2)
C5—S1—C4	104.53 (12)	N4—C5—S1	120.15 (18)
Cu1—O1—H1A	110 (2)	N4—C5—C6	122.7 (2)
Cu1—O1—H1B	118 (3)	C6—C5—S1	116.92 (19)
H1A—O1—H1B	107 (3)	C5—C6—H6	121.4
H3A—O3—H3B	109.4	C5—C6—C7	117.1 (2)
H4A—O4—H4B	108.7	C7—C6—H6	121.4
Cu1—O2—H2A	111.7	N3—C7—C6	120.7 (2)

Cu1—O2—H2B	114.0	N7—C7—N3	117.4 (2)
H2A—O2—H2B	106.1	N7—C7—C6	122.0 (2)
H5C—O5—H5D	123.9	N3—C8—N4	127.4 (2)
C1—N1—Cu1	124.00 (17)	N3—C8—H8	116.3
C4—N1—Cu1	118.91 (17)	N4—C8—H8	116.3
C4—N1—C1	115.4 (2)	N7—C9—H9A	109.5
C4—N2—C3	115.5 (2)	N7—C9—H9B	109.5
C8—N3—C7	116.2 (2)	N7—C9—H9C	109.5
C5—N4—Cu1	120.02 (17)	H9A—C9—H9B	109.5
C8—N4—Cu1	122.97 (17)	H9A—C9—H9C	109.5
C8—N4—C5	115.7 (2)	H9B—C9—H9C	109.5
H5A—N5—H5B	126 (3)	N7—C10—H10A	109.5
C1—N5—H5A	116 (2)	N7—C10—H10B	109.5
C1—N5—H5B	116 (2)	N7—C10—H10C	109.5
H6A—N6—H6B	122 (4)	H10A—C10—H10B	109.5
C3—N6—H6A	119 (2)	H10A—C10—H10C	109.5
C3—N6—H6B	118 (2)	H10B—C10—H10C	109.5
C7—N7—C9	120.8 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1A...S1 ⁱ	0.92 (1)	2.83 (3)	3.439 (2)	125 (3)
O1—H1A...N2 ⁱ	0.92 (1)	1.99 (2)	2.897 (3)	170 (3)
O1—H1B...O4 ⁱⁱ	0.91 (2)	1.82 (6)	2.71 (6)	167 (4)
O1—H1B...O5 ⁱⁱ	0.91 (2)	1.80 (6)	2.69 (6)	162 (4)
O3—H3A...Cl2	0.92	2.28	3.200 (7)	176
O3—H3B...O4	0.93	2.17	2.94 (6)	139
O4—H4A...Cl1 ⁱⁱ	0.91	2.97	3.46 (6)	116
O4—H4A...O3 ⁱⁱ	0.91	2.25	3.07 (6)	150
O4—H4B...Cl2 ⁱⁱⁱ	0.93	2.61	3.01 (6)	107
O2—H2A...O5 ⁱⁱⁱ	0.91	2.36	3.13 (6)	144
O2—H2B...Cl3	0.91	2.33	3.177 (7)	154
O5—H5C...Cl3	0.93	2.02	2.83 (6)	145
O5—H5C...O2 ⁱⁱⁱ	0.93	2.64	3.13 (6)	114
O5—H5D...Cl1 ⁱⁱ	0.91	2.96	3.45 (6)	116
O5—H5D...Cl3 ⁱⁱ	0.91	2.56	3.27 (6)	135
N5—H5A...Cl1	0.84 (3)	3.08 (3)	3.430 (3)	108 (2)
N5—H5A...Cl2	0.84 (3)	2.46 (3)	3.295 (3)	172 (3)
N5—H5A...O2	0.84 (3)	2.07 (3)	2.903 (7)	167 (3)
N5—H5B...Cl1 ^{iv}	0.79 (3)	2.56 (4)	3.353 (3)	173 (3)
N6—H6A...Cl3 ^{iv}	0.80 (4)	2.41 (4)	3.187 (4)	165 (3)
N6—H6A...O3 ^{iv}	0.80 (4)	2.32 (4)	3.094 (8)	163 (3)
N6—H6B...N3 ^v	0.81 (4)	2.76 (4)	3.323 (3)	128 (3)

Symmetry codes: (i) $-x+1, y-1/2, -z+1/2$; (ii) $-x+2, -y+1, -z+1$; (iii) $-x+2, -y+2, -z+1$; (iv) $-x+1, -y+2, -z+1$; (v) $x-1, y, z$.