

Crystal structure of 4,6-dimethyl-2-[[3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2H-pyran-2-yl]sulfanyl]nicotinonitrile

Doaa M. Masoud,^a Sherif F. Hammad,^b Galal H. Elgemeie^a and Peter G. Jones^{c*}

Received 18 October 2017

Accepted 18 October 2017

Edited by P. McArdle, National University of Ireland, Ireland

Keywords: crystal structure; pyridine; thio-glucose.**CCDC reference:** 1580696**Supporting information:** this article has supporting information at journals.iucr.org/e


^aChemistry Department, Faculty of Science, Helwan University, Cairo, Egypt, ^bPharmaceutical Chemistry Department, Faculty of Pharmacy, Helwan University, Cairo, Egypt, and ^cInstitut für Anorganische und Analytische Chemie, Technische Universität Braunschweig, Hagenring 30, D-38106 Braunschweig, Germany. *Correspondence e-mail: p.jones@tu-bs.de


In the title compound, C₁₄H₁₈N₂O₅S, the C–S bond lengths are unequal, with S–C_{glucose} = 1.8016 (15) Å and S–C_{pyridyl} = 1.7723 (13) Å. The hydrophilic glucose residues lie in the regions $z \simeq 0.25$ and 0.75 . Four classical hydrogen bonds link the molecules to form layers parallel to the *ab* plane, from which the pyridyl rings project; pyridyl ring stacking parallel to the *a* axis links adjacent layers.

1. Chemical context

The search for new anticancer chemotherapeutic agents continues to be an active area of research (Elgemeie, 2003; Elgemeie & Jones, 2004). In recent years nucleoside analogs have occupied a significant position in the search for effective chemotherapeutic agents, because many non-natural nucleoside derivatives have been shown to possess bioactivity (Elgemeie & Abou-Zeid, 2015). In the last few decades, pyridine derivatives have received considerable attention because of their wide-ranging applications as antimetabolic agents (Elgemeie *et al.*, 2009). Recently, we reported that many pyridine thioglycosides showed strong cytotoxicity against several human cancer cell lines and block proliferation of various cancer cell lines (Elgemeie, Abou-Zeid *et al.*, 2015). We also showed that thioglycosides involving pyridine and dihydropyridine groups exerted inhibitory effects on both DNA- and RNA-containing viruses and inhibitors of protein glycosylation, respectively (Elgemeie *et al.*, 2010). In view of these observations and with the aim of identifying new anticancer agents with improved pharmacokinetic and safety profiles, we have synthesized some new non-classical nucleoside analogs incorporating pyridine thioglycosides.

We report here a novel one-step synthesis of a pyridine-2-thioglycoside derivative by reaction of the pyridine-2(1H)-thione derivative (1) with 2,3,4,6-tetra-*O*-acetyl- α -D-glucopyranosyl bromide (2). Thus, (1) reacted with (2) in KOH/acetone to give a product for which two isomeric structures, (3) and (4), seemed possible, corresponding to two possible modes of glucosylation. After deprotection of the product (see Scheme), the final free sugar pyridinethione *N*-glucoside (5) or its regioisomer pyridine-2-thioglycoside (6) was obtained. Spectroscopic data cannot differentiate between these structures.





2. Structural commentary

The X-ray structure determination indicated unambiguously the formation of the pyridine-2-thioglucoside (**6**) as the product in the solid state. The molecule is shown in Fig. 1 and geometrical parameters are given in Table 1. The C—S bond lengths are markedly unequal, with S—C_{glucose} 1.8016 (15),

Table 1
Selected geometric parameters (Å, °).

S1—C12	1.7723 (13)	S1—C1	1.8016 (15)
C12—S1—C1	100.43 (6)		
C12—S1—C1—C2	152.98 (9)	C1—S1—C12—N11	−2.08 (14)

Table 2
Hydrogen-bond geometry (Å, °).

D—H...A	D—H	H...A	D...A	D—H...A
O2—H02...O3 ⁱ	0.80 (2)	2.01 (2)	2.7909 (16)	165 (3)
O3—H03...O5 ⁱⁱ	0.81 (2)	1.93 (2)	2.7394 (14)	174 (2)
O4—H04...O1 ⁱⁱⁱ	0.80 (2)	2.06 (2)	2.7490 (15)	144 (2)
O5—H05...O4 ^{iv}	0.79 (2)	1.96 (2)	2.7324 (18)	165 (3)
C2—H2...O5 ^{iv}	1.00	2.47	3.3326 (19)	144
C5—H5...N12 ^v	1.00	2.64	3.638 (2)	179

Symmetry codes: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x - 1, y, z$; (iii) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (v) $x, y - 1, z$.

S—C_{pyridyl} 1.7723 (13) Å. The main torsional degrees of freedom are between the rings, as defined by the torsion angles N11—C12—S1—C1 = −2.08 (14) and C2—C1—S1—C12 = 152.98 (9)°.

3. Supramolecular features

The glucose moieties of (**6**) occupy the regions at $z \approx 0.25$ and 0.75 . The layer structure of (**6**) is shown in Fig. 2. Each O—H donor forms one two-centre hydrogen bond (Table 2); the two most linear C—H...X interactions also lie within the layer, but are not drawn explicitly in Fig. 2. The same applies to the short contact S1...O5 ($\frac{1}{2} - x, \frac{1}{2} + y, \frac{1}{2} - z$) = 3.1417 (10) Å.

Adjacent layers are connected *via* the pyridyl rings, which project into the spaces between the hydrophilic layers and form π stacks parallel to the *a* axis. Adjacent rings in the stack are related by the twofold axis (operators $1 - x, y, 1 - z$ and $2 - x, y, 1 - z$). The interplanar angles are 4.33 (5)°, the centroid-to-centroid distances are 3.96 and 3.72 Å, and the ring offsets are *ca* 1.26 and 0.94 Å; these cannot be expressed exactly because neighbouring rings are not exactly parallel.




Figure 1
The structure of the title compound in the crystal. Displacement ellipsoids represent 50% probability levels.




Figure 2
Packing diagram of the title compound, viewed perpendicular to the *ab* plane. Classical hydrogen bonds are indicated by dashed lines. Methyl and nitrile substituents of the pyridine rings have been omitted for clarity.

4. Database survey

Perhaps surprisingly, a database search revealed only one other example of a pyridine ring with a thioglucose substituent at the 2-position, namely pyridyl thioglucose monohydrate (Nordenson & Jeffrey, 1980; refcode PYSGPR). This compound also shows a marked inequality between the S—C bond lengths (*cf.* Table 1); S—C_{glucose} is 1.793 (3), S—C_{pyridyl} is 1.759 (3) Å.

5. Synthesis and crystallization

To a solution of the pyridine-2-(1*H*)-thione (**1**) (1.64 gm, 0.01 mol) in aqueous potassium hydroxide (6 ml, 0.56 g, 0.01 mol) was added a solution of 2,3,4,6-tetra-*O*-acetyl- α -D-glucopyranosyl bromide (**2**) (4.52 g, 0.011 mol) in acetone (30 ml). The reaction mixture was stirred at room temperature until the reaction was judged complete by TLC (30 min to 2 h). The mixture was evaporated under reduced pressure at 313 K and the residue was washed with distilled water to remove the potassium bromide. The solid was collected by filtration and crystallized from ethanol to give compound (**3**) in 85% yield (m.p. 468 K). Dry gaseous ammonia was then passed through a solution of the protected thioglycoside (**3**) (0.5 g) in dry methanol (20 ml) at 273 K for 0.5 h, then the mixture was stirred at 273 K until completion of the reaction (TLC, 2–6 h). The mixture was evaporated at 313 K to give a solid residue, which was recrystallized from ethanol to give compound (**6**) in 85% yield (m.p. 482–483 K).

IR (KBr): 3600–3258 (OH), 2222 (CN) cm⁻¹. ¹H NMR (DMSO-*d*₆): δ 2.22 (*s*, 3H, CH₃), 2.31 (*s*, 3H, CH₃), 3.15–3.80

Table 3
Experimental details.

Crystal data	
Chemical formula	C ₁₄ H ₁₈ N ₂ O ₅ S
<i>M</i> _r	326.36
Crystal system, space group	Monoclinic, <i>I</i> 2
Temperature (K)	100
<i>a</i> , <i>b</i> , <i>c</i> (Å)	7.66978 (18), 8.72860 (13), 23.7524 (4)
β (°)	98.7356 (16)
<i>V</i> (Å ³)	1571.69 (5)
<i>Z</i>	4
Radiation type	Mo <i>K</i> α
μ (mm ⁻¹)	0.23
Crystal size (mm)	0.40 × 0.35 × 0.20
Data collection	
Diffractometer	Oxford Diffraction Xcalibur Eos
Absorption correction	Multi-scan (<i>CrysAlis PRO</i> ; Rigaku Oxford Diffraction, 2015)
<i>T</i> _{min} , <i>T</i> _{max}	0.980, 1.000
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	84844, 4741, 4590
<i>R</i> _{int}	0.027
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.729
Refinement	
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.024, 0.063, 1.05
No. of reflections	4741
No. of parameters	217
No. of restraints	7
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	0.30, -0.17
Absolute structure	Flack <i>x</i> determined using 2066 quotients [(<i>I</i> ⁺) - (<i>I</i> ⁻)] / [(<i>I</i> ⁺) + (<i>I</i> ⁻)] (Parsons <i>et al.</i> , 2013)
Absolute structure parameter	-0.008 (8)

Computer programs: *CrysAlis PRO* (Rigaku Oxford Diffraction, 2015), *SHELXS97* (Sheldrick, 2008), *SHELXL2017* (Sheldrick, 2015) and *XP* (Siemens, 1994).

(*m*, 6H, 2H-6', H-5', H-4', H-3', H-2'), 4.40 (*d*, *J* = 9.55 Hz, 2H, HO-2' and HO-3'), 4.90 (*s*, 1H, HO-4'), 5.30 (*s*, 1H, HO-6'), 5.59 (*d*, *J*_{1,2} = 9.86 Hz, 1H, H-1'), 7.19 (*s*, 1H, pyridine H-5) ppm. ¹³C NMR: δ 20.7 (CH₃), 22.9 (CH₃), 61.0 (C6'), 68.9 (C4'), 72.7 (C2'), 75.9 (C3'), 80.7 (C5'), 83.7 (C1'), 103.2 (C3), 116.0 (CN), 119.7 (C5), 149.2 (C4), 159.0 (C6), 164.0 (C2) ppm.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. The space group as initially found by the diffractometer program was *P*2₁, with two independent but virtually identical molecules in the asymmetric unit. It became apparent that the two molecules formed layer structures independent of each other, and were related by a translation vector (0.5, 0.5, 0.5). The *checkCIF* program also indicated that the true space group should be centred, with a 100% fit and a small deviation. The same cell was retained for ease of checking, and the structure determination and refinement repeated in space group *I*2. The refinement was entirely satisfactory, and corresponds to the structure presented here. However, the reflections with (*h* + *k* + *l*) odd, which are required to be systematically absent in *I*2, seemed to

be quite definitely present. We are unable to explain this anomaly. The HKL file appended to the CIF contains these reflections.

Crystal data, data collection and structure refinement details are summarized in Table 2. OH hydrogen atoms were refined freely but with an O—H distance restraint (SADI). Methyl groups were refined as idealized rigid groups allowed to rotate but not tip (AFIX 137), with C—H 0.98 Å and H—C—H 109.5°. Other hydrogen atoms were included using a riding model starting from calculated positions (C—H_{aromatic} 0.95, C—H_{methylene} 0.99, C—H_{methine} 1.00 Å).

The absolute configuration was determined by the unambiguous Flack parameter of $-0.008(8)$ (Parsons *et al.*, 2013).

References

Elgemeie, G. H. (2003). *Curr. Pharm. Design*, **9**, 2627–2642.

- Elgemeie, G. H., Abou-Zeid, M., Alsaïd, S., Hebshy, A. & Essa, H. (2015). *Nucleosides Nucleotides Nucleic Acids*, **34**, 659–673.
- Elgemeie, G. H. & Abu-Zaïed, M. (2015). *Nucleosides Nucleotides Nucleic Acids*, **34**, 834–847.
- Elgemeie, G. H., Eltamny, E. H., Elgawad, I. I. & Mahmoud, N. M. (2009). *Synth. Commun.* **39**, 443–458.
- Elgemeie, G. H. & Jones, P. G. (2004). *Acta Cryst.* **E60**, o2107–o2109.
- Elgemeie, G. H., Mahdy, E. M., Elgawish, M. A., Ahmed, M. M., Shousha, W. G. & Eldin, M. E. (2010). *Z. Naturforsch.* **65c**, 577–587.
- Nordenson, S. & Jeffrey, G. A. (1980). *Acta Cryst.* **B36**, 1214–1216.
- Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
- Rigaku Oxford Diffraction (2015). *CrysAlis PRO*. Rigaku Oxford Diffraction Ltd, Yarnton, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Siemens (1994). *XP*. Siemens Analytical X-ray Instruments, Madison, Wisconsin, USA.

supporting information

Acta Cryst. (2017). E73, 1751-1754 [https://doi.org/10.1107/S2056989017015213]

Crystal structure of 4,6-dimethyl-2-[[3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2H-pyran-2-yl]sulfonyl]nicotinonitrile

Doaa M. Masoud, Sherif F. Hammad, Galal H. Elgemeie and Peter G. Jones

Computing details

Data collection: *CrysAlis PRO* (Rigaku Oxford Diffraction, 2015); cell refinement: *CrysAlis PRO* (Rigaku Oxford Diffraction, 2015); data reduction: *CrysAlis PRO* (Rigaku Oxford Diffraction, 2015); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2017* (Sheldrick, 2015); molecular graphics: *XP* (Siemens, 1994); software used to prepare material for publication: *SHELXL2017* (Sheldrick, 2015).

4,6-Dimethyl-2-[[3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2H-pyran-2-yl]sulfonyl]nicotinonitrile

Crystal data

$C_{14}H_{18}N_2O_5S$

$M_r = 326.36$

Monoclinic, *I2*

$a = 7.66978$ (18) Å

$b = 8.72860$ (13) Å

$c = 23.7524$ (4) Å

$\beta = 98.7356$ (16)°

$V = 1571.69$ (5) Å³

$Z = 4$

$F(000) = 688$

$D_x = 1.379$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 46975 reflections

$\theta = 2.7\text{--}30.9^\circ$

$\mu = 0.23$ mm⁻¹

$T = 100$ K

Block, colourless

0.40 × 0.35 × 0.20 mm

Data collection

Oxford Diffraction Xcalibur Eos
diffractometer

Radiation source: Enhance (Mo) X-ray Source

Detector resolution: 16.1419 pixels mm⁻¹

ω -scan

Absorption correction: multi-scan

(*CrysAlis PRO*; Rigaku Oxford Diffraction,
2015)

$T_{\min} = 0.980$, $T_{\max} = 1.000$

84844 measured reflections

4741 independent reflections

4590 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.027$

$\theta_{\max} = 31.2^\circ$, $\theta_{\min} = 2.5^\circ$

$h = -10 \rightarrow 11$

$k = -12 \rightarrow 12$

$l = -34 \rightarrow 33$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.024$

$wR(F^2) = 0.063$

$S = 1.05$

4741 reflections

217 parameters

7 restraints

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0354P)^2 + 0.4674P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.30$ e Å⁻³

$\Delta\rho_{\min} = -0.17$ e Å⁻³

Absolute structure: Flack x determined using
2066 quotients $[(F^+)-(F^-)]/[(F^+)+(F^-)]$ (Parsons *et al.*, 2013)
Absolute structure parameter: -0.008 (8)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.59062 (4)	0.54939 (4)	0.37504 (2)	0.01953 (8)
C1	0.57772 (17)	0.35296 (17)	0.35150 (5)	0.0175 (2)
H1	0.557606	0.284848	0.383749	0.021*
O1	0.74310 (12)	0.31545 (12)	0.33376 (4)	0.0183 (2)
C2	0.42750 (17)	0.33167 (17)	0.30152 (5)	0.0177 (3)
H2	0.444291	0.404110	0.270164	0.021*
O2	0.26172 (13)	0.35885 (13)	0.31878 (4)	0.0200 (2)
H02	0.223 (4)	0.438 (3)	0.3050 (12)	0.054 (8)*
C3	0.42986 (17)	0.16706 (17)	0.27975 (5)	0.0184 (2)
H3	0.399828	0.095244	0.309645	0.022*
O3	0.30583 (13)	0.14881 (14)	0.22910 (4)	0.0226 (2)
H03	0.207 (3)	0.161 (3)	0.2365 (9)	0.033 (6)*
C4	0.61090 (18)	0.12809 (18)	0.26570 (6)	0.0185 (3)
H4	0.636616	0.194695	0.233704	0.022*
O4	0.61422 (15)	-0.02842 (14)	0.24885 (5)	0.0250 (2)
H04	0.672 (3)	-0.035 (3)	0.2237 (9)	0.039 (6)*
C5	0.74991 (17)	0.15622 (18)	0.31794 (5)	0.0186 (2)
H5	0.722628	0.091101	0.350129	0.022*
C6	0.93568 (18)	0.12113 (19)	0.30688 (5)	0.0210 (3)
H6A	1.021081	0.174340	0.335809	0.025*
H6B	0.957132	0.009636	0.311368	0.025*
O5	0.96641 (13)	0.16663 (14)	0.25148 (4)	0.0208 (2)
H05	0.957 (3)	0.256 (2)	0.2480 (11)	0.043 (7)*
N11	0.71744 (19)	0.38506 (16)	0.46739 (5)	0.0247 (3)
C12	0.68518 (18)	0.52490 (16)	0.44732 (5)	0.0193 (3)
C13	0.7223 (2)	0.65733 (19)	0.48057 (6)	0.0219 (3)
C14	0.8011 (2)	0.64238 (19)	0.53789 (6)	0.0235 (3)
C15	0.8355 (2)	0.4955 (2)	0.55786 (6)	0.0274 (3)
H15	0.889755	0.479840	0.596114	0.033*
C16	0.7915 (2)	0.3696 (2)	0.52236 (6)	0.0290 (3)
C17	0.6799 (3)	0.8053 (2)	0.45591 (7)	0.0313 (4)
N12	0.6448 (3)	0.9219 (2)	0.43492 (7)	0.0491 (5)
C18	0.8435 (3)	0.7795 (2)	0.57556 (7)	0.0357 (4)
H18A	0.876528	0.745852	0.615104	0.054*
H18B	0.739865	0.846339	0.572717	0.054*

H18C	0.941903	0.835962	0.563468	0.054*
C19	0.8230 (4)	0.2094 (2)	0.54374 (8)	0.0472 (5)
H19A	0.779315	0.137031	0.513317	0.071*
H19B	0.760624	0.192842	0.576301	0.071*
H19C	0.949717	0.193415	0.555621	0.071*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.01731 (15)	0.02917 (15)	0.01144 (12)	-0.00026 (13)	0.00001 (10)	0.00067 (12)
C1	0.0108 (6)	0.0311 (7)	0.0110 (5)	-0.0011 (5)	0.0029 (4)	-0.0017 (5)
O1	0.0094 (4)	0.0321 (5)	0.0138 (4)	-0.0017 (4)	0.0028 (3)	-0.0016 (4)
C2	0.0091 (6)	0.0334 (7)	0.0107 (5)	-0.0015 (5)	0.0023 (4)	-0.0006 (5)
O2	0.0097 (4)	0.0329 (5)	0.0181 (4)	0.0011 (4)	0.0045 (3)	0.0002 (4)
C3	0.0094 (6)	0.0328 (7)	0.0132 (5)	-0.0023 (5)	0.0029 (4)	-0.0023 (5)
O3	0.0084 (4)	0.0429 (6)	0.0162 (4)	-0.0026 (4)	0.0014 (3)	-0.0066 (4)
C4	0.0093 (6)	0.0329 (7)	0.0136 (5)	-0.0026 (5)	0.0029 (4)	-0.0033 (5)
O4	0.0159 (5)	0.0371 (6)	0.0239 (5)	-0.0036 (4)	0.0095 (4)	-0.0100 (5)
C5	0.0114 (6)	0.0322 (6)	0.0123 (5)	-0.0006 (5)	0.0025 (4)	-0.0015 (5)
C6	0.0104 (6)	0.0404 (8)	0.0124 (5)	0.0011 (5)	0.0018 (4)	-0.0012 (5)
O5	0.0113 (4)	0.0378 (6)	0.0138 (4)	-0.0013 (4)	0.0040 (3)	-0.0025 (4)
N11	0.0303 (7)	0.0304 (6)	0.0123 (5)	-0.0068 (5)	-0.0003 (5)	0.0028 (5)
C12	0.0153 (6)	0.0318 (8)	0.0107 (5)	-0.0027 (5)	0.0022 (4)	0.0005 (5)
C13	0.0213 (7)	0.0316 (7)	0.0131 (6)	-0.0011 (6)	0.0039 (5)	-0.0001 (5)
C14	0.0249 (7)	0.0330 (8)	0.0127 (6)	-0.0042 (6)	0.0030 (5)	-0.0021 (5)
C15	0.0331 (8)	0.0353 (7)	0.0124 (6)	-0.0069 (6)	-0.0010 (5)	0.0010 (5)
C16	0.0397 (9)	0.0319 (8)	0.0139 (6)	-0.0065 (7)	-0.0009 (6)	0.0042 (6)
C17	0.0441 (10)	0.0340 (8)	0.0151 (6)	0.0055 (7)	0.0029 (6)	-0.0042 (6)
N12	0.0849 (15)	0.0372 (9)	0.0232 (7)	0.0143 (9)	0.0022 (8)	-0.0021 (6)
C18	0.0511 (11)	0.0371 (9)	0.0175 (7)	-0.0039 (8)	0.0005 (7)	-0.0059 (6)
C19	0.0847 (17)	0.0335 (9)	0.0182 (7)	-0.0065 (10)	-0.0093 (8)	0.0069 (7)

Geometric parameters (Å, °)

S1—C12	1.7723 (13)	C6—H6A	0.9900
S1—C1	1.8016 (15)	C6—H6B	0.9900
C1—O1	1.4338 (16)	O5—H05	0.79 (2)
C1—C2	1.5348 (17)	N11—C12	1.3201 (19)
C1—H1	1.0000	N11—C16	1.3492 (19)
O1—C5	1.4428 (18)	C12—C13	1.405 (2)
C2—O2	1.4141 (16)	C13—C14	1.4092 (19)
C2—C3	1.528 (2)	C13—C17	1.435 (2)
C2—H2	1.0000	C14—C15	1.379 (2)
O2—H02	0.80 (2)	C14—C18	1.500 (2)
C3—O3	1.4247 (16)	C15—C16	1.394 (2)
C3—C4	1.5152 (18)	C15—H15	0.9500
C3—H3	1.0000	C16—C19	1.495 (2)
O3—H03	0.809 (19)	C17—N12	1.147 (2)

C4—O4	1.4249 (18)	C18—H18A	0.9800
C4—C5	1.5280 (18)	C18—H18B	0.9800
C4—H4	1.0000	C18—H18C	0.9800
O4—H04	0.800 (19)	C19—H19A	0.9800
C5—C6	1.5187 (18)	C19—H19B	0.9800
C5—H5	1.0000	C19—H19C	0.9800
C6—O5	1.4279 (16)		
C12—S1—C1	100.43 (6)	O5—C6—H6A	108.9
O1—C1—C2	109.80 (10)	C5—C6—H6A	108.9
O1—C1—S1	107.40 (9)	O5—C6—H6B	108.9
C2—C1—S1	110.77 (10)	C5—C6—H6B	108.9
O1—C1—H1	109.6	H6A—C6—H6B	107.8
C2—C1—H1	109.6	C6—O5—H05	110.3 (19)
S1—C1—H1	109.6	C12—N11—C16	118.04 (14)
C1—O1—C5	111.41 (10)	N11—C12—C13	123.16 (13)
O2—C2—C3	108.25 (11)	N11—C12—S1	119.22 (10)
O2—C2—C1	110.96 (10)	C13—C12—S1	117.62 (11)
C3—C2—C1	109.24 (11)	C12—C13—C14	119.18 (14)
O2—C2—H2	109.5	C12—C13—C17	119.82 (13)
C3—C2—H2	109.5	C14—C13—C17	121.00 (15)
C1—C2—H2	109.5	C15—C14—C13	116.75 (14)
C2—O2—H02	109 (2)	C15—C14—C18	121.60 (13)
O3—C3—C4	107.80 (10)	C13—C14—C18	121.64 (15)
O3—C3—C2	110.49 (11)	C14—C15—C16	120.56 (14)
C4—C3—C2	110.12 (11)	C14—C15—H15	119.7
O3—C3—H3	109.5	C16—C15—H15	119.7
C4—C3—H3	109.5	N11—C16—C15	122.29 (15)
C2—C3—H3	109.5	N11—C16—C19	116.40 (15)
C3—O3—H03	109.1 (16)	C15—C16—C19	121.31 (14)
O4—C4—C3	109.46 (12)	N12—C17—C13	178.31 (17)
O4—C4—C5	110.01 (12)	C14—C18—H18A	109.5
C3—C4—C5	109.57 (11)	C14—C18—H18B	109.5
O4—C4—H4	109.3	H18A—C18—H18B	109.5
C3—C4—H4	109.3	C14—C18—H18C	109.5
C5—C4—H4	109.3	H18A—C18—H18C	109.5
C4—O4—H04	108.6 (19)	H18B—C18—H18C	109.5
O1—C5—C6	108.14 (11)	C16—C19—H19A	109.5
O1—C5—C4	108.53 (11)	C16—C19—H19B	109.5
C6—C5—C4	112.61 (11)	H19A—C19—H19B	109.5
O1—C5—H5	109.2	C16—C19—H19C	109.5
C6—C5—H5	109.2	H19A—C19—H19C	109.5
C4—C5—H5	109.2	H19B—C19—H19C	109.5
O5—C6—C5	113.19 (11)		
C12—S1—C1—O1	-87.12 (9)	C3—C4—C5—C6	-179.39 (13)
C12—S1—C1—C2	152.98 (9)	O1—C5—C6—O5	-81.24 (14)
C2—C1—O1—C5	-63.63 (14)	C4—C5—C6—O5	38.67 (18)

S1—C1—O1—C5	175.85 (8)	C16—N11—C12—C13	-0.8 (2)
O1—C1—C2—O2	176.40 (11)	C16—N11—C12—S1	179.18 (12)
S1—C1—C2—O2	-65.14 (13)	C1—S1—C12—N11	-2.08 (14)
O1—C1—C2—C3	57.13 (14)	C1—S1—C12—C13	177.90 (11)
S1—C1—C2—C3	175.59 (8)	N11—C12—C13—C14	1.4 (2)
O2—C2—C3—O3	66.09 (13)	S1—C12—C13—C14	-178.62 (11)
C1—C2—C3—O3	-172.97 (10)	N11—C12—C13—C17	-178.71 (16)
O2—C2—C3—C4	-174.93 (11)	S1—C12—C13—C17	1.31 (19)
C1—C2—C3—C4	-54.00 (14)	C12—C13—C14—C15	-0.6 (2)
O3—C3—C4—O4	-62.97 (14)	C17—C13—C14—C15	179.51 (16)
C2—C3—C4—O4	176.42 (10)	C12—C13—C14—C18	-179.89 (15)
O3—C3—C4—C5	176.33 (12)	C17—C13—C14—C18	0.2 (2)
C2—C3—C4—C5	55.72 (15)	C13—C14—C15—C16	-0.7 (3)
C1—O1—C5—C6	-173.06 (10)	C18—C14—C15—C16	178.65 (17)
C1—O1—C5—C4	64.49 (13)	C12—N11—C16—C15	-0.5 (3)
O4—C4—C5—O1	179.92 (10)	C12—N11—C16—C19	178.95 (18)
C3—C4—C5—O1	-59.70 (14)	C14—C15—C16—N11	1.3 (3)
O4—C4—C5—C6	60.23 (16)	C14—C15—C16—C19	-178.17 (19)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H02 \cdots O3 ⁱ	0.80 (2)	2.01 (2)	2.7909 (16)	165 (3)
O3—H03 \cdots O5 ⁱⁱ	0.81 (2)	1.93 (2)	2.7394 (14)	174 (2)
O4—H04 \cdots O1 ⁱⁱⁱ	0.80 (2)	2.06 (2)	2.7490 (15)	144 (2)
O5—H05 \cdots O4 ^{iv}	0.79 (2)	1.96 (2)	2.7324 (18)	165 (3)
C2—H2 \cdots O5 ^{iv}	1.00	2.47	3.3326 (19)	144
C5—H5 \cdots N12 ^v	1.00	2.64	3.638 (2)	179

Symmetry codes: (i) $-x+1/2, y+1/2, -z+1/2$; (ii) $x-1, y, z$; (iii) $-x+3/2, y-1/2, -z+1/2$; (iv) $-x+3/2, y+1/2, -z+1/2$; (v) $x, y-1, z$.