



Crystal structures of two isomeric 2-aryl-3-phenyl-1,3-thiazepan-4-ones

Hemant P. Yennawar,^a Samuel D. Peterson^b and Lee J. Silverberg^{b*}

^aPennsylvania State University, Department of Biochemistry and Molecular Biology, 108 Althouse Laboratory, University Park, PA, 16801, USA, and ^bPennsylvania State University Schuylkill Campus, 200 University Drive, Schuylkill Haven, PA, 17972, USA. *Correspondence e-mail: ljs43@psu.edu

Received 8 July 2019
Accepted 22 July 2019

Edited by W. T. A. Harrison, University of Aberdeen, Scotland

Keywords: crystal structure; thiazepan; seven membered ring; chair pucker; isomers.


CCDC references: 1942358; 1942357

Supporting information: this article has supporting information at journals.iucr.org/e

The crystal of 6-(3-nitrophenyl)-7-phenyl-5-thia-7-azaspiro[2.6]nonan-8-one (**1**), C₁₉H₁₈N₂O₃S, has monoclinic (*P*2₁/*n*) symmetry while that of its isomer 6-(4-nitrophenyl)-7-phenyl-5-thia-7-azaspiro[2.6]nonan-8-one (**2**), has orthorhombic (*Pca*2₁) symmetry: compound **1** has two molecules, *A* and *B*, in the asymmetric unit while **2** has one. In all three molecules, the seven-membered thiazepan ring exhibits a chair conformation with *Q*2 and *Q*3 values (Å) of 0.521 (3), 0.735 (3) and 0.485 (3), 0.749 (3) in **1** and 0.517 (5), 0.699 (5) in **2**. In each structure, the phenyl rings attached to adjacent atoms of the thiazepan ring have interplanar angles ranging between 41 and 47°. Except for the nitro groups, the three molecules have similar conformations when overlaid in pairs. Both crystal structures are consolidated by C—H···O hydrogen bonds.

1. Chemical context


The seven-membered 1,3-thiazepan-4-one ring system, like the similar six-membered 1,3-thiazin-4-one and five-membered 1,3-thiazolidin-4-one systems, is biologically active and of potential medicinal use. For example, the Bristol-Myers Squibb ACE/NEP inhibitor omapatrilat advanced to Phase II clinical trials (Graul *et al.*, 1999; Robl *et al.* 1997; Tabrizchi, 2001; Cozier *et al.* 2018). In fact, nearly all of the known compounds with this ring system are related in structure to omapatrilat.




Previously we reported the synthesis and crystal structure of 6,7-diphenyl-5-thia-7-azaspiro[2.6]nonan-8-one (Yennawar & Silverberg, 2013). Herein we report the T3P-promoted synthesis and crystal structures of two new analogs: 6-(4-nitrophenyl)-7-phenyl-5-thia-7-azaspiro[2.6]nonan-8-one (**1**) and 6-(3-nitrophenyl)-7-phenyl-5-thia-7-azaspiro[2.6]nonan-8-one (**2**), in which a nitro group substitutes at the *para* and *meta* positions, respectively, of the C-2 aromatic ring.

2. Structural commentary


Compound **1** crystallizes with two molecules, *A* (containing S1) and *B* (containing S2), in the asymmetric unit (Fig. 1) and




Figure 1

The molecular structure of **1** with displacement ellipsoids drawn at the 50% probability level. C—H...O interactions are shown as dashed lines.

2 crystallizes with one molecule (Fig. 2). The configurations of the stereogenic centers in the arbitrarily chosen asymmetric molecules are (*S*) at C1 and (*R*) at C20 for **1** and (*S*) at C1 for **2**; in both structures, crystal symmetry generates a racemic mixture. These molecules adopt similar conformations and


Figure 2

The molecular structure of **2** with displacement ellipsoids drawn at the 50% probability level.

Table 1

Hydrogen-bond geometry (Å, °) for **1**.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C1—H1...O6	0.98	2.41	3.390 (3)	175
C3—H3B...O6	0.97	2.50	3.469 (3)	174
C10—H10...O6 ⁱ	0.93	2.60	3.408 (3)	146
C17—H17...O2 ⁱⁱ	0.93	2.46	3.216 (4)	138
C20—H20...O3 ⁱⁱⁱ	0.98	2.41	3.369 (3)	165
C28—H28...O3 ⁱⁱⁱ	0.93	2.56	3.384 (4)	147

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, y - 1, z$; (iii) $x, y + 1, z$.

Table 2

Hydrogen-bond geometry (Å, °) for **2**.


<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C1—H1...O1 ⁱ	0.98	2.38	3.352 (6)	173
C3—H3B...O1 ⁱ	0.97	2.48	3.444 (5)	171

Symmetry code: (i) $-x + \frac{1}{2}, y, z + \frac{1}{2}$.

overlay closely (Fig. 3) apart from the nitro groups. The seven-membered thiazepan rings in both structures adopt chair conformations. The puckering parameters [*Q*2 and *Q*3 (Å)] as calculated by *PLATON* (Spek, 2009) for molecules *A* and *B* in **1** are 0.521 (3), 0.735 (3) and 0.485 (3), 0.749 (3), respectively, with equivalent values of 0.517 (5), 0.699 (5) for **2**. The dihedral angles between the aromatic rings attached to the 2 and 3 positions of the thiazepan rings are 46.93 (15) (molecule **1A**), 42.50 (15) (**1B**) and 42.0 (3)° (**2**).

3. Supramolecular features

The extended structure of **1** has more extensive hydrogen bonding compared to that of **2** (Tables 1 and 2). In **1**, the


Figure 3

Overlay of the molecule of **2** on the similarly handed molecule of **1**.

molecules are arranged into layers propagating in the *ab* plane, with C—H...O hydrogen bonds in both the *a*- and *b*-axis directions, but not in the *c*-axis direction (Fig. 4). In **2**, the molecules link up *via* C—H...O 'head-to-tail' hydrogen bonds in the *c*-axis direction (Fig. 5) and hydrophobic interactions between adjacent chains consolidate the packing in the *a*- and *b*-axis directions.

4. Database survey

A 1,3-thiazepan-4-one with a 5,6-fused benzene and a 2,3-fused triazole has been reported, but only an *ORTEP* representation was given, without any other data (Bakavoli *et al.*, 2002). The structures of omapatrilat bound to proteins have been published recently (Cozier, *et al.* 2018). The 2,3-diphenyl structure that we previously reported showed a chair-type conformation for the thiazepan ring [CSD (Groom *et al.*, 2016) refcode MIHVOQ; Yennawar & Silverberg, 2013] like those reported here.

5. Synthesis and crystallization

A two-necked 25 ml round-bottom flask was oven-dried, cooled under N₂, and charged with a stir bar. 3- or 4-Nitrobenzaldehyde (0.907 g, 6 mmol), aniline (0.571 g, 6 mmol), and [1-(sulfanylmethyl)cyclopropyl] acetic acid (0.877 g, 6 mmol) were added. Pyridine (1.95 ml, 24 mmol) was added. Finally, 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphorinane-2,4,6-trioxide (T3P) in 2-methyltetrahydrofuran (50 weight %; 7.3 ml, 12 mmol) was added. The reaction was stirred at room temperature and followed by TLC. The mixture was poured

into a separatory funnel with dichloromethane and distilled water. The layers were separated and the aqueous layer was then extracted twice with dichloromethane. The organic fractions were combined and washed with saturated sodium bicarbonate and saturated sodium chloride and then dried over sodium sulfate and concentrated under vacuum. Further purification was carried out as indicated below for each compound.

6-(3-Nitrophenyl)-7-phenyl-5-thia-7-azaspiro[2.6]nonan-8-one (**1**): Chromatography on 30 g flash silica gel with mixtures of ethyl acetate and hexanes gave a solid. Recrystallization from 2-propanol solution gave crystals (0.5192 g, 26%), m.p. 457–458 K. X-ray-quality crystals were grown by slow evaporation from a 2-propanol solution.

6-(4-Nitrophenyl)-7-phenyl-5-thia-7-azaspiro[2.6]nonan-8-one (**2**): Chromatography on 30 g flash silica gel with mixtures of ethyl acetate and hexanes gave a solid. Recrystallization from ethyl acetate solution gave colorless crystals (0.1804 g, 9%), m.p. 480–482 K (decomposition). X-ray-quality crystals were grown by slow evaporation from an ethyl acetate solution.




Figure 4
Packing diagram for **1** showing C—H...O hydrogen bonds between molecules arranged in the *ab* planes.




Figure 5
Packing diagram for **2** viewed down the *b*-axis direction.

Table 3
Experimental details.

	1	2
Crystal data		
Chemical formula	C ₁₉ H ₁₈ N ₂ O ₃ S	C ₁₉ H ₁₈ N ₂ O ₃ S
<i>M_r</i>	354.41	354.41
Crystal system, space group	Monoclinic, <i>P2₁/n</i>	Orthorhombic, <i>Pca2₁</i>
Temperature (K)	298	298
<i>a</i> , <i>b</i> , <i>c</i> (Å)	16.993 (4), 9.955 (2), 21.243 (5)	17.478 (3), 10.4125 (19), 9.7129 (17)
α , β , γ (°)	90, 99.531 (4), 90	90, 90, 90
<i>V</i> (Å ³)	3543.9 (15)	1767.6 (5)
<i>Z</i>	8	4
Radiation type	Mo <i>K</i> α	Mo <i>K</i> α
μ (mm ⁻¹)	0.20	0.20
Crystal size (mm)	0.24 × 0.12 × 0.09	0.27 × 0.1 × 0.04
Data collection		
Diffractometer	Bruker SMART CCD area detector	Bruker SMART CCD area detector
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2001)	Multi-scan (<i>SADABS</i> ; Bruker, 2001)
<i>T_{min}</i> , <i>T_{max}</i>	0.780, 0.9	0.769, 0.9
No. of measured, independent and observed [<i>I</i> > 2σ(<i>I</i>)] reflections	30602, 8495, 3560	15240, 4261, 2403
<i>R_{int}</i>	0.083	0.056
(sin θ /λ) _{max} (Å ⁻¹)	0.669	0.668
Refinement		
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.057, 0.155, 0.94	0.081, 0.249, 1.01
No. of reflections	8495	4261
No. of parameters	451	226
No. of restraints	0	1
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	0.25, -0.26	0.55, -0.30
Absolute structure	–	Flack (1983)
Absolute structure parameter	–	0.47 (19)

Computer programs: *SMART* and *SAINT* (Bruker, 2001), *SHELXS* (Sheldrick, 2008), *SHELXL* (Sheldrick, 2015) and *OLEX2* (Dolomanov *et al.*, 2009).

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. The hydrogen atoms were placed geometrically (C–H = 0.93–0.98 Å) and refined as riding on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Acknowledgements

We thank Euticals (AMRI) for the gift of T3P in 2-methyltetrahydrofuran, Oakwood Chemicals for [1-(sulfanylmethyl)cyclopropyl] acetic acid, and Penn State Schuylkill for financial support.

References

Bakavoli, M., Davoodnia, A., Rahimizadeh, M., Heravi, M. M. & Ghasemzadeh, M. (2002). *J. Chem. Res.* pp. 178–179.
 Bruker (2001). *SADABS*, *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.

Cozier, G. E., Arendse, L. B., Schwager, S. L., Sturrock, E. D. & Acharya, K. R. (2018). *J. Med. Chem.* **61**, 10141–10154.
 Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Graul, A., Leeson, P. & Castañer, J. (1999). *Drugs Fut.* **24**, 269–277.
 Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
 Robl, J. A., Sun, C.-Q., Stevenson, J., Ryono, D. E., Simpkins, L. M., Cimarusti, M. P., Dejneka, T., Slusarchyk, W. A., Chao, S., Stratton, L., Misra, R. N., Bednarz, M. S., Asaad, M. M., Cheung, H. S., Abboa-Offei, B. E., Smith, P. L., Mathers, P. D., Fox, M., Schaeffer, T. R., Seymour, A. A. & Trippodo, N. C. (1997). *J. Med. Chem.* **40**, 1570–1577.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Tabrizchi, R. (2001). *Curr. Opin. Investig. Drugs*, **2**, 1414–1422.
 Yennawar, H. P. & Silverberg, L. J. (2013). *Acta Cryst.* **E69**, o1659.

supporting information

Acta Cryst. (2019). E75, 1270-1273 [https://doi.org/10.1107/S2056989019010429]

Crystal structures of two isomeric 2-aryl-3-phenyl-1,3-thiazepan-4-ones

Hemant P. Yennawar, Samuel D. Peterson and Lee J. Silverberg

Computing details

For both structures, data collection: *SMART* (Bruker, 2001); cell refinement: *SAINTE* (Bruker, 2001); data reduction: *SAINTE* (Bruker, 2001); program(s) used to solve structure: *SHELXS* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL* (Sheldrick, 2015); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2* (Dolomanov *et al.*, 2009).

6-(3-Nitrophenyl)-7-phenyl-5-thia-7-azaspiro[2.6]nonan-8-one (1)

Crystal data

$C_{19}H_{18}N_2O_3S$

$M_r = 354.41$

Monoclinic, $P2_1/n$

$a = 16.993$ (4) Å

$b = 9.955$ (2) Å

$c = 21.243$ (5) Å

$\beta = 99.531$ (4)°

$V = 3543.9$ (15) Å³

$Z = 8$

$F(000) = 1488$

$D_x = 1.329$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2681 reflections

$\theta = 2.4$ – 28.3 °

$\mu = 0.20$ mm⁻¹

$T = 298$ K

Block, colorless

$0.24 \times 0.12 \times 0.09$ mm

Data collection

Bruker SMART CCD area detector
diffractometer

Radiation source: fine-focus sealed tube

Parallel-graphite monochromator

Detector resolution: 8.34 pixels mm⁻¹

phi and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2001)

$T_{\min} = 0.780$, $T_{\max} = 0.9$

30602 measured reflections

8495 independent reflections

3560 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.083$

$\theta_{\max} = 28.4$ °, $\theta_{\min} = 1.4$ °

$h = -21$ → 22

$k = -12$ → 13

$l = -26$ → 28

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.057$

$wR(F^2) = 0.155$

$S = 0.94$

8495 reflections

451 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0603P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.25$ e Å⁻³

$\Delta\rho_{\min} = -0.26$ e Å⁻³

Special details

Experimental. The data collection nominally covered a full sphere of reciprocal space by a combination of 4 sets of ω scans each set at different φ and/or 2θ angles and each scan (10 s exposure) covering -0.300° degrees in ω . The crystal to detector distance was 5.86 cm.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.38777 (15)	0.3850 (3)	0.24330 (13)	0.0472 (7)
H1	0.4111	0.4733	0.2374	0.057*
C2	0.52215 (16)	0.2834 (3)	0.24439 (13)	0.0458 (7)
C3	0.53440 (16)	0.3814 (3)	0.19259 (13)	0.0497 (7)
H3A	0.5906	0.3831	0.1890	0.060*
H3B	0.5197	0.4707	0.2048	0.060*
C4	0.48636 (17)	0.3471 (3)	0.12818 (13)	0.0529 (7)
C5	0.40156 (18)	0.3958 (3)	0.11569 (14)	0.0673 (9)
H5A	0.3793	0.3748	0.0718	0.081*
H5B	0.4021	0.4928	0.1198	0.081*
C6	0.5063 (2)	0.2202 (3)	0.09693 (16)	0.0785 (10)
H6A	0.4628	0.1711	0.0717	0.094*
H6B	0.5485	0.1646	0.1197	0.094*
C7	0.5293 (2)	0.3518 (3)	0.07165 (15)	0.0748 (10)
H7A	0.4996	0.3826	0.0313	0.090*
H7B	0.5853	0.3761	0.0793	0.090*
C8	0.32870 (15)	0.4017 (3)	0.28901 (13)	0.0461 (7)
C9	0.26482 (16)	0.3140 (3)	0.28908 (14)	0.0551 (8)
H9	0.2578	0.2431	0.2602	0.066*
C10	0.21165 (16)	0.3302 (3)	0.33116 (15)	0.0598 (8)
H10	0.1691	0.2710	0.3301	0.072*
C11	0.22164 (17)	0.4337 (3)	0.37465 (15)	0.0582 (8)
H11	0.1861	0.4458	0.4031	0.070*
C12	0.28542 (16)	0.5189 (3)	0.37502 (13)	0.0496 (7)
C13	0.33890 (16)	0.5053 (3)	0.33310 (13)	0.0484 (7)
H13	0.3813	0.5649	0.3345	0.058*
C14	0.43844 (15)	0.1901 (3)	0.31425 (14)	0.0456 (7)
C15	0.45269 (18)	0.2164 (3)	0.37871 (15)	0.0593 (8)
H15	0.4730	0.2995	0.3935	0.071*
C16	0.4370 (2)	0.1200 (4)	0.42134 (16)	0.0750 (10)
H16	0.4467	0.1380	0.4649	0.090*
C17	0.4070 (2)	-0.0028 (3)	0.39960 (18)	0.0771 (10)
H17	0.3961	-0.0679	0.4283	0.093*

C18	0.39337 (19)	-0.0286 (3)	0.33550 (18)	0.0715 (9)
H18	0.3733	-0.1118	0.3209	0.086*
C19	0.40877 (17)	0.0665 (3)	0.29254 (15)	0.0582 (8)
H19	0.3993	0.0479	0.2490	0.070*
N1	0.45322 (12)	0.2936 (2)	0.27006 (10)	0.0443 (6)
N2	0.29752 (18)	0.6265 (3)	0.42288 (12)	0.0603 (7)
O1	0.24710 (15)	0.6440 (2)	0.45664 (11)	0.0846 (7)
O2	0.35847 (16)	0.6932 (2)	0.42706 (11)	0.0858 (7)
O3	0.57139 (11)	0.19619 (19)	0.26254 (10)	0.0632 (6)
S1	0.33553 (5)	0.32922 (9)	0.16613 (4)	0.0704 (3)
C20	0.64468 (15)	0.8845 (3)	0.29410 (13)	0.0484 (7)
H20	0.6223	0.9700	0.2766	0.058*
C21	0.52037 (16)	0.7803 (3)	0.23659 (13)	0.0435 (7)
C22	0.52413 (16)	0.8762 (3)	0.18223 (13)	0.0538 (8)
H22A	0.5306	0.9668	0.1992	0.065*
H22B	0.4738	0.8727	0.1531	0.065*
C23	0.59097 (18)	0.8473 (3)	0.14525 (14)	0.0594 (8)
C24	0.67190 (18)	0.9037 (4)	0.17153 (14)	0.0725 (10)
H24A	0.6666	0.9999	0.1766	0.087*
H24B	0.7066	0.8894	0.1402	0.087*
C25	0.5874 (2)	0.7178 (4)	0.10892 (16)	0.0812 (10)
H25A	0.6374	0.6731	0.1062	0.097*
H25B	0.5429	0.6581	0.1114	0.097*
C26	0.5701 (2)	0.8472 (4)	0.07334 (16)	0.0862 (11)
H26A	0.5152	0.8656	0.0545	0.103*
H26B	0.6097	0.8807	0.0492	0.103*
C27	0.68160 (15)	0.9101 (3)	0.36255 (13)	0.0473 (7)
C28	0.66026 (17)	1.0246 (3)	0.39258 (15)	0.0611 (8)
H28	0.6247	1.0854	0.3700	0.073*
C29	0.6914 (2)	1.0492 (4)	0.45580 (16)	0.0747 (10)
H29	0.6770	1.1271	0.4752	0.090*
C30	0.74311 (19)	0.9604 (4)	0.49030 (15)	0.0686 (9)
H30	0.7634	0.9761	0.5331	0.082*
C31	0.76402 (16)	0.8479 (3)	0.45990 (14)	0.0525 (7)
C32	0.73499 (16)	0.8210 (3)	0.39678 (14)	0.0514 (7)
H32	0.7510	0.7442	0.3774	0.062*
C33	0.58190 (15)	0.6820 (3)	0.33613 (13)	0.0439 (7)
C34	0.55008 (18)	0.7007 (3)	0.39071 (15)	0.0611 (8)
H34	0.5234	0.7801	0.3966	0.073*
C35	0.5576 (2)	0.6016 (4)	0.43721 (16)	0.0797 (10)
H35	0.5369	0.6149	0.4746	0.096*
C36	0.5957 (2)	0.4837 (4)	0.42757 (18)	0.0775 (11)
H36	0.6007	0.4169	0.4586	0.093*
C37	0.62626 (18)	0.4640 (3)	0.37303 (19)	0.0719 (10)
H37	0.6514	0.3832	0.3667	0.086*
C38	0.62024 (16)	0.5631 (3)	0.32684 (14)	0.0549 (8)
H38	0.6418	0.5497	0.2898	0.066*
N3	0.57806 (12)	0.7883 (2)	0.28935 (10)	0.0425 (5)

N4	0.81967 (15)	0.7509 (3)	0.49575 (14)	0.0637 (7)
O4	0.85870 (14)	0.7871 (2)	0.54671 (11)	0.0852 (7)
O5	0.82594 (15)	0.6398 (3)	0.47322 (12)	0.0904 (8)
O6	0.46794 (11)	0.69436 (18)	0.23280 (9)	0.0551 (5)
S2	0.72025 (4)	0.83452 (9)	0.24715 (4)	0.0681 (3)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0411 (15)	0.0448 (17)	0.0557 (18)	0.0023 (13)	0.0080 (13)	0.0011 (14)
C2	0.0440 (16)	0.0494 (18)	0.0444 (17)	0.0019 (14)	0.0080 (13)	-0.0006 (14)
C3	0.0465 (16)	0.0500 (17)	0.0548 (19)	0.0047 (13)	0.0146 (14)	0.0025 (15)
C4	0.0653 (19)	0.0506 (18)	0.0436 (18)	0.0059 (15)	0.0115 (15)	0.0017 (15)
C5	0.070 (2)	0.077 (2)	0.053 (2)	0.0036 (18)	0.0039 (16)	0.0057 (17)
C6	0.110 (3)	0.066 (2)	0.060 (2)	0.008 (2)	0.016 (2)	-0.0039 (19)
C7	0.093 (2)	0.079 (3)	0.058 (2)	0.010 (2)	0.0283 (19)	0.0086 (19)
C8	0.0420 (15)	0.0431 (17)	0.0531 (18)	0.0012 (13)	0.0072 (14)	-0.0008 (14)
C9	0.0480 (16)	0.0530 (19)	0.064 (2)	-0.0036 (14)	0.0083 (15)	-0.0085 (15)
C10	0.0428 (16)	0.065 (2)	0.073 (2)	-0.0086 (15)	0.0132 (16)	-0.0031 (18)
C11	0.0503 (18)	0.067 (2)	0.059 (2)	0.0045 (16)	0.0153 (15)	0.0025 (18)
C12	0.0555 (18)	0.0419 (17)	0.0504 (19)	0.0056 (14)	0.0064 (15)	0.0006 (15)
C13	0.0439 (15)	0.0446 (17)	0.0564 (19)	-0.0010 (13)	0.0072 (14)	0.0023 (15)
C14	0.0462 (15)	0.0408 (17)	0.0502 (19)	-0.0002 (13)	0.0090 (13)	0.0037 (15)
C15	0.074 (2)	0.0500 (19)	0.053 (2)	-0.0070 (16)	0.0077 (16)	-0.0028 (16)
C16	0.112 (3)	0.064 (2)	0.052 (2)	-0.004 (2)	0.020 (2)	0.0035 (19)
C17	0.102 (3)	0.059 (2)	0.077 (3)	-0.001 (2)	0.035 (2)	0.013 (2)
C18	0.086 (2)	0.046 (2)	0.084 (3)	-0.0154 (17)	0.020 (2)	0.001 (2)
C19	0.067 (2)	0.0486 (19)	0.059 (2)	-0.0024 (16)	0.0108 (16)	-0.0057 (17)
N1	0.0422 (13)	0.0429 (13)	0.0483 (14)	0.0020 (10)	0.0087 (11)	0.0027 (11)
N2	0.0754 (19)	0.0507 (17)	0.0552 (18)	0.0073 (15)	0.0121 (15)	0.0002 (14)
O1	0.1002 (18)	0.0907 (18)	0.0683 (16)	0.0102 (15)	0.0300 (14)	-0.0169 (14)
O2	0.1053 (19)	0.0678 (16)	0.0867 (18)	-0.0234 (14)	0.0228 (15)	-0.0174 (13)
O3	0.0565 (12)	0.0635 (14)	0.0708 (14)	0.0210 (11)	0.0139 (11)	0.0136 (11)
S1	0.0524 (5)	0.0998 (7)	0.0557 (5)	-0.0098 (4)	-0.0008 (4)	-0.0024 (5)
C20	0.0453 (15)	0.0448 (17)	0.0525 (18)	-0.0066 (13)	0.0007 (14)	0.0001 (14)
C21	0.0384 (15)	0.0459 (17)	0.0455 (17)	0.0038 (13)	0.0051 (13)	-0.0022 (14)
C22	0.0504 (17)	0.0551 (19)	0.0525 (19)	-0.0061 (14)	-0.0017 (14)	0.0043 (15)
C23	0.0622 (19)	0.070 (2)	0.0460 (19)	-0.0133 (17)	0.0093 (15)	0.0019 (16)
C24	0.069 (2)	0.099 (3)	0.051 (2)	-0.0195 (19)	0.0147 (17)	0.0054 (19)
C25	0.086 (3)	0.085 (3)	0.076 (2)	-0.008 (2)	0.024 (2)	-0.015 (2)
C26	0.092 (3)	0.113 (3)	0.054 (2)	-0.027 (2)	0.0099 (19)	0.003 (2)
C27	0.0440 (15)	0.0468 (17)	0.0491 (18)	-0.0076 (14)	0.0019 (14)	-0.0021 (15)
C28	0.0618 (19)	0.056 (2)	0.063 (2)	0.0039 (16)	0.0020 (16)	-0.0065 (17)
C29	0.082 (2)	0.071 (2)	0.069 (2)	0.0066 (19)	0.006 (2)	-0.023 (2)
C30	0.066 (2)	0.084 (3)	0.053 (2)	-0.0101 (19)	0.0008 (17)	-0.013 (2)
C31	0.0452 (16)	0.059 (2)	0.0519 (19)	-0.0063 (15)	0.0033 (14)	0.0022 (17)
C32	0.0483 (16)	0.0521 (18)	0.0513 (19)	-0.0059 (14)	0.0013 (14)	-0.0077 (15)
C33	0.0386 (14)	0.0460 (17)	0.0453 (17)	-0.0038 (13)	0.0017 (13)	0.0016 (14)

C34	0.066 (2)	0.062 (2)	0.059 (2)	-0.0022 (16)	0.0184 (17)	0.0022 (18)
C35	0.092 (3)	0.093 (3)	0.056 (2)	-0.019 (2)	0.017 (2)	0.013 (2)
C36	0.075 (2)	0.080 (3)	0.070 (3)	-0.020 (2)	-0.012 (2)	0.031 (2)
C37	0.058 (2)	0.056 (2)	0.096 (3)	0.0022 (16)	-0.007 (2)	0.016 (2)
C38	0.0510 (17)	0.0523 (19)	0.060 (2)	0.0010 (15)	0.0058 (15)	0.0003 (17)
N3	0.0404 (12)	0.0415 (13)	0.0435 (14)	-0.0048 (10)	0.0007 (11)	0.0036 (11)
N4	0.0548 (16)	0.080 (2)	0.0546 (19)	-0.0085 (16)	0.0048 (14)	0.0122 (17)
O4	0.0749 (15)	0.117 (2)	0.0558 (15)	-0.0125 (14)	-0.0133 (13)	0.0113 (14)
O5	0.1009 (19)	0.0781 (18)	0.0858 (19)	0.0195 (15)	-0.0034 (15)	0.0073 (15)
O6	0.0462 (11)	0.0529 (12)	0.0628 (13)	-0.0086 (10)	-0.0012 (10)	0.0042 (10)
S2	0.0479 (4)	0.0946 (7)	0.0624 (6)	-0.0072 (4)	0.0112 (4)	-0.0025 (5)

Geometric parameters (Å, °)

C1—H1	0.9800	C20—H20	0.9800
C1—C8	1.517 (4)	C20—C27	1.507 (4)
C1—N1	1.476 (3)	C20—N3	1.474 (3)
C1—S1	1.817 (3)	C20—S2	1.821 (3)
C2—C3	1.510 (4)	C21—C22	1.508 (4)
C2—N1	1.375 (3)	C21—N3	1.364 (3)
C2—O3	1.223 (3)	C21—O6	1.228 (3)
C3—H3A	0.9700	C22—H22A	0.9700
C3—H3B	0.9700	C22—H22B	0.9700
C3—C4	1.512 (4)	C22—C23	1.512 (4)
C4—C5	1.502 (4)	C23—C24	1.505 (4)
C4—C6	1.491 (4)	C23—C25	1.499 (4)
C4—C7	1.505 (4)	C23—C26	1.510 (4)
C5—H5A	0.9700	C24—H24A	0.9700
C5—H5B	0.9700	C24—H24B	0.9700
C5—S1	1.801 (3)	C24—S2	1.814 (3)
C6—H6A	0.9700	C25—H25A	0.9700
C6—H6B	0.9700	C25—H25B	0.9700
C6—C7	1.492 (4)	C25—C26	1.498 (5)
C7—H7A	0.9700	C26—H26A	0.9700
C7—H7B	0.9700	C26—H26B	0.9700
C8—C9	1.393 (4)	C27—C28	1.384 (4)
C8—C13	1.384 (4)	C27—C32	1.385 (4)
C9—H9	0.9300	C28—H28	0.9300
C9—C10	1.382 (4)	C28—C29	1.381 (4)
C10—H10	0.9300	C29—H29	0.9300
C10—C11	1.376 (4)	C29—C30	1.370 (4)
C11—H11	0.9300	C30—H30	0.9300
C11—C12	1.375 (4)	C30—C31	1.369 (4)
C12—C13	1.380 (4)	C31—C32	1.376 (4)
C12—N2	1.468 (4)	C31—N4	1.472 (4)
C13—H13	0.9300	C32—H32	0.9300
C14—C15	1.376 (4)	C33—C34	1.371 (4)
C14—C19	1.379 (4)	C33—C38	1.381 (4)

C14—N1	1.444 (3)	C33—N3	1.446 (3)
C15—H15	0.9300	C34—H34	0.9300
C15—C16	1.375 (4)	C34—C35	1.387 (4)
C16—H16	0.9300	C35—H35	0.9300
C16—C17	1.374 (4)	C35—C36	1.372 (5)
C17—H17	0.9300	C36—H36	0.9300
C17—C18	1.367 (4)	C36—C37	1.360 (5)
C18—H18	0.9300	C37—H37	0.9300
C18—C19	1.370 (4)	C37—C38	1.383 (4)
C19—H19	0.9300	C38—H38	0.9300
N2—O1	1.217 (3)	N4—O4	1.226 (3)
N2—O2	1.221 (3)	N4—O5	1.216 (3)
C8—C1—H1	107.9	C27—C20—H20	107.1
C8—C1—S1	109.56 (18)	C27—C20—S2	110.86 (18)
N1—C1—H1	107.9	N3—C20—H20	107.1
N1—C1—C8	110.9 (2)	N3—C20—C27	111.6 (2)
N1—C1—S1	112.55 (18)	N3—C20—S2	112.84 (18)
S1—C1—H1	107.9	S2—C20—H20	107.1
N1—C2—C3	118.1 (2)	N3—C21—C22	118.5 (2)
O3—C2—C3	121.2 (3)	O6—C21—C22	120.9 (2)
O3—C2—N1	120.7 (3)	O6—C21—N3	120.6 (2)
C2—C3—H3A	108.9	C21—C22—H22A	108.7
C2—C3—H3B	108.9	C21—C22—H22B	108.7
C2—C3—C4	113.2 (2)	C21—C22—C23	114.1 (2)
H3A—C3—H3B	107.7	H22A—C22—H22B	107.6
C4—C3—H3A	108.9	C23—C22—H22A	108.7
C4—C3—H3B	108.9	C23—C22—H22B	108.7
C5—C4—C3	116.4 (2)	C22—C23—C26	117.1 (3)
C6—C4—C3	117.8 (3)	C24—C23—C22	117.1 (3)
C6—C4—C5	118.5 (3)	C24—C23—C26	115.0 (3)
C6—C4—C7	59.7 (2)	C25—C23—C22	117.5 (3)
C7—C4—C3	117.0 (3)	C25—C23—C24	117.9 (3)
C7—C4—C5	115.2 (3)	C25—C23—C26	59.7 (2)
C4—C5—H5A	108.1	C23—C24—H24A	108.3
C4—C5—H5B	108.1	C23—C24—H24B	108.3
C4—C5—S1	116.8 (2)	C23—C24—S2	116.0 (2)
H5A—C5—H5B	107.3	H24A—C24—H24B	107.4
S1—C5—H5A	108.1	S2—C24—H24A	108.3
S1—C5—H5B	108.1	S2—C24—H24B	108.3
C4—C6—H6A	117.7	C23—C25—H25A	117.7
C4—C6—H6B	117.7	C23—C25—H25B	117.7
C4—C6—C7	60.6 (2)	H25A—C25—H25B	114.8
H6A—C6—H6B	114.8	C26—C25—C23	60.5 (2)
C7—C6—H6A	117.7	C26—C25—H25A	117.7
C7—C6—H6B	117.7	C26—C25—H25B	117.7
C4—C7—H7A	117.8	C23—C26—H26A	117.8
C4—C7—H7B	117.8	C23—C26—H26B	117.8

C6—C7—C4	59.7 (2)	C25—C26—C23	59.8 (2)
C6—C7—H7A	117.8	C25—C26—H26A	117.8
C6—C7—H7B	117.8	C25—C26—H26B	117.8
H7A—C7—H7B	114.9	H26A—C26—H26B	114.9
C9—C8—C1	122.0 (3)	C28—C27—C20	119.0 (3)
C13—C8—C1	119.5 (2)	C28—C27—C32	119.0 (3)
C13—C8—C9	118.4 (3)	C32—C27—C20	122.0 (3)
C8—C9—H9	119.3	C27—C28—H28	119.8
C10—C9—C8	121.4 (3)	C29—C28—C27	120.4 (3)
C10—C9—H9	119.3	C29—C28—H28	119.8
C9—C10—H10	120.0	C28—C29—H29	119.5
C11—C10—C9	120.0 (3)	C30—C29—C28	121.0 (3)
C11—C10—H10	120.0	C30—C29—H29	119.5
C10—C11—H11	120.8	C29—C30—H30	121.0
C10—C11—C12	118.4 (3)	C29—C30—C31	118.0 (3)
C12—C11—H11	120.8	C31—C30—H30	121.0
C11—C12—C13	122.6 (3)	C30—C31—C32	122.7 (3)
C11—C12—N2	118.3 (3)	C30—C31—N4	119.2 (3)
C13—C12—N2	119.1 (3)	C32—C31—N4	118.2 (3)
C8—C13—H13	120.4	C27—C32—H32	120.5
C12—C13—C8	119.2 (3)	C31—C32—C27	119.0 (3)
C12—C13—H13	120.4	C31—C32—H32	120.5
C15—C14—C19	119.8 (3)	C34—C33—C38	119.9 (3)
C15—C14—N1	119.3 (2)	C34—C33—N3	120.2 (3)
C19—C14—N1	120.8 (3)	C38—C33—N3	119.9 (3)
C14—C15—H15	120.0	C33—C34—H34	119.9
C16—C15—C14	120.1 (3)	C33—C34—C35	120.2 (3)
C16—C15—H15	120.0	C35—C34—H34	119.9
C15—C16—H16	120.0	C34—C35—H35	120.3
C17—C16—C15	120.0 (3)	C36—C35—C34	119.5 (3)
C17—C16—H16	120.0	C36—C35—H35	120.3
C16—C17—H17	120.2	C35—C36—H36	119.8
C18—C17—C16	119.6 (3)	C37—C36—C35	120.5 (3)
C18—C17—H17	120.2	C37—C36—H36	119.8
C17—C18—H18	119.5	C36—C37—H37	119.7
C17—C18—C19	120.9 (3)	C36—C37—C38	120.5 (3)
C19—C18—H18	119.5	C38—C37—H37	119.7
C14—C19—H19	120.2	C33—C38—C37	119.4 (3)
C18—C19—C14	119.5 (3)	C33—C38—H38	120.3
C18—C19—H19	120.2	C37—C38—H38	120.3
C2—N1—C1	122.0 (2)	C21—N3—C20	122.3 (2)
C2—N1—C14	117.2 (2)	C21—N3—C33	117.8 (2)
C14—N1—C1	119.3 (2)	C33—N3—C20	118.8 (2)
O1—N2—C12	118.7 (3)	O4—N4—C31	118.1 (3)
O1—N2—O2	123.4 (3)	O5—N4—C31	118.9 (3)
O2—N2—C12	117.8 (3)	O5—N4—O4	123.0 (3)
C5—S1—C1	99.49 (14)	C24—S2—C20	97.13 (14)

C1—C8—C9—C10	179.8 (3)	C20—C27—C28—C29	178.3 (3)
C1—C8—C13—C12	-179.3 (2)	C20—C27—C32—C31	-177.4 (2)
C2—C3—C4—C5	84.2 (3)	C21—C22—C23—C24	-84.2 (3)
C2—C3—C4—C6	-65.5 (3)	C21—C22—C23—C25	65.0 (3)
C2—C3—C4—C7	-133.7 (3)	C21—C22—C23—C26	133.1 (3)
C3—C2—N1—C1	7.6 (4)	C22—C21—N3—C20	-1.7 (4)
C3—C2—N1—C14	173.2 (2)	C22—C21—N3—C33	-169.5 (2)
C3—C4—C5—S1	-63.3 (3)	C22—C23—C24—S2	66.5 (3)
C3—C4—C6—C7	-106.7 (3)	C22—C23—C25—C26	106.9 (3)
C3—C4—C7—C6	107.9 (3)	C22—C23—C26—C25	-107.6 (3)
C4—C5—S1—C1	59.6 (3)	C23—C24—S2—C20	-60.9 (3)
C5—C4—C6—C7	104.2 (3)	C24—C23—C25—C26	-104.1 (3)
C5—C4—C7—C6	-109.6 (3)	C24—C23—C26—C25	109.0 (3)
C6—C4—C5—S1	86.2 (3)	C25—C23—C24—S2	-82.6 (3)
C7—C4—C5—S1	154.0 (2)	C26—C23—C24—S2	-150.1 (3)
C8—C1—N1—C2	-166.4 (2)	C27—C20—N3—C21	159.1 (2)
C8—C1—N1—C14	28.2 (3)	C27—C20—N3—C33	-33.2 (3)
C8—C1—S1—C5	153.1 (2)	C27—C20—S2—C24	-149.8 (2)
C8—C9—C10—C11	-0.7 (4)	C27—C28—C29—C30	-0.8 (5)
C9—C8—C13—C12	-0.6 (4)	C28—C27—C32—C31	1.2 (4)
C9—C10—C11—C12	-0.3 (4)	C28—C29—C30—C31	1.1 (5)
C10—C11—C12—C13	0.8 (4)	C29—C30—C31—C32	-0.2 (5)
C10—C11—C12—N2	-178.0 (2)	C29—C30—C31—N4	179.9 (3)
C11—C12—C13—C8	-0.4 (4)	C30—C31—C32—C27	-0.9 (4)
C11—C12—N2—O1	-6.4 (4)	C30—C31—N4—O4	-15.7 (4)
C11—C12—N2—O2	172.9 (3)	C30—C31—N4—O5	165.6 (3)
C13—C8—C9—C10	1.1 (4)	C32—C27—C28—C29	-0.4 (4)
C13—C12—N2—O1	174.8 (3)	C32—C31—N4—O4	164.4 (3)
C13—C12—N2—O2	-5.9 (4)	C32—C31—N4—O5	-14.4 (4)
C14—C15—C16—C17	0.0 (5)	C33—C34—C35—C36	-1.3 (5)
C15—C14—C19—C18	0.4 (4)	C34—C33—C38—C37	-0.2 (4)
C15—C14—N1—C1	-91.8 (3)	C34—C33—N3—C20	91.3 (3)
C15—C14—N1—C2	102.2 (3)	C34—C33—N3—C21	-100.4 (3)
C15—C16—C17—C18	0.3 (5)	C34—C35—C36—C37	0.2 (5)
C16—C17—C18—C19	-0.3 (5)	C35—C36—C37—C38	0.8 (5)
C17—C18—C19—C14	0.0 (5)	C36—C37—C38—C33	-0.9 (4)
C19—C14—C15—C16	-0.3 (4)	C38—C33—C34—C35	1.2 (4)
C19—C14—N1—C1	86.9 (3)	C38—C33—N3—C20	-86.4 (3)
C19—C14—N1—C2	-79.1 (3)	C38—C33—N3—C21	81.9 (3)
N1—C1—C8—C9	-87.2 (3)	N3—C20—C27—C28	-99.3 (3)
N1—C1—C8—C13	91.4 (3)	N3—C20—C27—C32	79.3 (3)
N1—C1—S1—C5	-83.0 (2)	N3—C20—S2—C24	84.2 (2)
N1—C2—C3—C4	-77.3 (3)	N3—C21—C22—C23	71.4 (3)
N1—C14—C15—C16	178.3 (3)	N3—C33—C34—C35	-176.5 (3)
N1—C14—C19—C18	-178.3 (3)	N3—C33—C38—C37	177.5 (2)
N2—C12—C13—C8	178.4 (2)	N4—C31—C32—C27	179.0 (2)
O3—C2—C3—C4	101.7 (3)	O6—C21—C22—C23	-106.5 (3)
O3—C2—N1—C1	-171.4 (2)	O6—C21—N3—C20	176.2 (2)

O3—C2—N1—C14	−5.8 (4)	O6—C21—N3—C33	8.4 (4)
S1—C1—C8—C9	37.6 (3)	S2—C20—C27—C28	134.0 (2)
S1—C1—C8—C13	−143.7 (2)	S2—C20—C27—C32	−47.4 (3)
S1—C1—N1—C2	70.4 (3)	S2—C20—N3—C21	−75.3 (3)
S1—C1—N1—C14	−94.9 (2)	S2—C20—N3—C33	92.4 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C1—H1...O6	0.98	2.41	3.390 (3)	175
C3—H3B...O6	0.97	2.50	3.469 (3)	174
C10—H10...O6 ⁱ	0.93	2.60	3.408 (3)	146
C17—H17...O2 ⁱⁱ	0.93	2.46	3.216 (4)	138
C20—H20...O3 ⁱⁱⁱ	0.98	2.41	3.369 (3)	165
C28—H28...O3 ⁱⁱⁱ	0.93	2.56	3.384 (4)	147

Symmetry codes: (i) $-x+1/2, y-1/2, -z+1/2$; (ii) $x, y-1, z$; (iii) $x, y+1, z$.

6-(4-Nitrophenyl)-7-phenyl-5-thia-7-azaspiro[2.6]nonan-8-one (2)*Crystal data*

$C_{19}H_{18}N_2O_3S$

$M_r = 354.41$

Orthorhombic, *Pca*2₁

$a = 17.478$ (3) Å

$b = 10.4125$ (19) Å

$c = 9.7129$ (17) Å

$V = 1767.6$ (5) Å³

$Z = 4$

$F(000) = 744$

$D_x = 1.332$ Mg m^{−3}

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3246 reflections

$\theta = 2.3$ – 28.2°

$\mu = 0.20$ mm^{−1}

$T = 298$ K

Needle, colorless

$0.27 \times 0.1 \times 0.04$ mm

Data collection

Bruker SMART CCD area detector
diffractometer

Radiation source: fine-focus sealed tube

Parallel graphite monochromator

Detector resolution: 8.34 pixels mm^{−1}

phi and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2001)

$T_{\min} = 0.769, T_{\max} = 0.9$

15240 measured reflections

4261 independent reflections

2403 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.056$

$\theta_{\max} = 28.3^\circ, \theta_{\min} = 2.0^\circ$

$h = -23 \rightarrow 20$

$k = -13 \rightarrow 13$

$l = -12 \rightarrow 12$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.081$

$wR(F^2) = 0.249$

$S = 1.01$

4261 reflections

226 parameters

1 restraint

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.1501P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.55$ e Å^{−3}

$\Delta\rho_{\min} = -0.30$ e Å^{−3}

Absolute structure: Flack (1983)

Absolute structure parameter: 0.47 (19)

Special details

Experimental. The data collection nominally covered a full sphere of reciprocal space by a combination of 4 sets of ω scans each set at different φ and/or 2θ angles and each scan (20 s exposure) covering -0.300° degrees in ω . The crystal to detector distance was 5.82 cm.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6266 (3)	0.5630 (4)	0.3841 (4)	0.0635 (11)
H1	0.6481	0.5871	0.4736	0.076*
C2	0.7512 (2)	0.6299 (3)	0.2850 (5)	0.0559 (9)
C3	0.7521 (3)	0.7400 (4)	0.3845 (4)	0.0593 (9)
H3A	0.8027	0.7785	0.3838	0.071*
H3B	0.7429	0.7071	0.4764	0.071*
C4	0.6941 (3)	0.8423 (4)	0.3537 (5)	0.0684 (12)
C5	0.6136 (3)	0.8248 (5)	0.4002 (8)	0.0895 (16)
H5A	0.5854	0.9020	0.3769	0.107*
H5B	0.6138	0.8180	0.4998	0.107*
C6	0.5840 (3)	0.4389 (4)	0.4049 (5)	0.0677 (11)
C7	0.6041 (3)	0.3639 (5)	0.5202 (6)	0.0840 (15)
H7	0.6424	0.3920	0.5794	0.101*
C8	0.5672 (4)	0.2488 (6)	0.5459 (8)	0.0912 (16)
H8	0.5794	0.1990	0.6223	0.109*
C9	0.5123 (3)	0.2113 (5)	0.4549 (7)	0.0836 (15)
C10	0.4898 (3)	0.2793 (6)	0.3448 (8)	0.0952 (18)
H10	0.4516	0.2491	0.2864	0.114*
C11	0.5263 (3)	0.3988 (5)	0.3204 (7)	0.0794 (13)
H11	0.5109	0.4499	0.2469	0.095*
C12	0.6820 (2)	0.4565 (4)	0.1787 (4)	0.0550 (9)
C13	0.7041 (3)	0.3314 (4)	0.1968 (6)	0.0697 (11)
H13	0.7288	0.3067	0.2773	0.084*
C14	0.6893 (3)	0.2408 (5)	0.0934 (6)	0.0780 (14)
H14	0.7037	0.1555	0.1050	0.094*
C15	0.6533 (3)	0.2795 (5)	-0.0250 (6)	0.0763 (13)
H15	0.6425	0.2196	-0.0933	0.092*
C16	0.6336 (3)	0.4032 (5)	-0.0435 (5)	0.0734 (12)
H16	0.6105	0.4283	-0.1254	0.088*
C17	0.6473 (3)	0.4935 (5)	0.0581 (5)	0.0646 (10)
H17	0.6332	0.5787	0.0448	0.078*

C18	0.7205 (4)	0.9800 (4)	0.3554 (6)	0.0846 (15)
H18A	0.7735	0.9967	0.3789	0.102*
H18B	0.6845	1.0446	0.3871	0.102*
C19	0.7041 (4)	0.9149 (4)	0.2180 (5)	0.0822 (15)
H19A	0.7474	0.8924	0.1603	0.099*
H19B	0.6584	0.9404	0.1685	0.099*
N1	0.69179 (18)	0.5453 (3)	0.2889 (3)	0.0568 (8)
N2	0.4715 (4)	0.0877 (6)	0.4819 (10)	0.120 (2)
O1	0.80230 (16)	0.6180 (3)	0.1992 (3)	0.0659 (7)
O2	0.4223 (4)	0.0545 (6)	0.3992 (11)	0.169 (3)
O3	0.4905 (4)	0.0269 (6)	0.5872 (9)	0.151 (3)
S1	0.56211 (7)	0.69074 (13)	0.3337 (2)	0.0942 (5)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.058 (2)	0.080 (3)	0.053 (2)	−0.0090 (19)	0.0124 (18)	−0.0078 (19)
C2	0.052 (2)	0.061 (2)	0.055 (2)	−0.0039 (17)	−0.0023 (19)	−0.0011 (18)
C3	0.064 (2)	0.063 (2)	0.050 (2)	−0.0029 (18)	−0.0001 (19)	−0.0036 (18)
C4	0.084 (3)	0.055 (2)	0.067 (3)	0.0001 (18)	−0.003 (2)	−0.005 (2)
C5	0.075 (3)	0.077 (3)	0.116 (4)	0.012 (2)	0.023 (3)	−0.008 (3)
C6	0.052 (2)	0.075 (3)	0.076 (3)	−0.0025 (19)	0.011 (2)	−0.002 (2)
C7	0.077 (3)	0.094 (4)	0.081 (4)	0.002 (3)	0.015 (3)	0.005 (3)
C8	0.086 (4)	0.091 (4)	0.096 (4)	0.007 (3)	0.017 (3)	0.010 (3)
C9	0.076 (3)	0.077 (3)	0.098 (4)	0.000 (2)	0.033 (3)	−0.005 (3)
C10	0.054 (3)	0.110 (4)	0.122 (5)	−0.014 (2)	0.015 (3)	−0.017 (4)
C11	0.059 (3)	0.088 (3)	0.091 (3)	−0.012 (2)	0.005 (3)	0.000 (3)
C12	0.053 (2)	0.064 (2)	0.049 (2)	−0.0016 (15)	−0.0009 (17)	−0.0045 (18)
C13	0.076 (3)	0.066 (2)	0.068 (3)	0.004 (2)	−0.007 (2)	0.004 (2)
C14	0.100 (4)	0.060 (3)	0.074 (3)	0.000 (2)	−0.001 (3)	−0.008 (2)
C15	0.079 (3)	0.077 (3)	0.073 (3)	−0.014 (2)	0.004 (3)	−0.015 (2)
C16	0.071 (3)	0.093 (3)	0.055 (2)	−0.015 (2)	−0.006 (2)	−0.006 (2)
C17	0.068 (3)	0.070 (3)	0.057 (2)	−0.006 (2)	−0.009 (2)	0.000 (2)
C18	0.099 (4)	0.061 (3)	0.095 (4)	−0.003 (2)	−0.014 (3)	−0.013 (3)
C19	0.112 (4)	0.074 (3)	0.060 (3)	−0.003 (3)	−0.020 (3)	0.005 (2)
N1	0.0504 (18)	0.072 (2)	0.0480 (18)	−0.0073 (13)	0.0019 (15)	−0.0070 (15)
N2	0.075 (3)	0.086 (3)	0.200 (7)	0.002 (3)	0.055 (4)	0.010 (5)
O1	0.0552 (16)	0.0804 (17)	0.0621 (18)	−0.0054 (13)	0.0086 (15)	−0.0031 (15)
O2	0.107 (4)	0.113 (4)	0.288 (10)	−0.043 (3)	0.024 (5)	−0.013 (5)
O3	0.148 (5)	0.105 (4)	0.201 (7)	−0.004 (3)	0.043 (5)	0.037 (4)
S1	0.0546 (6)	0.0848 (8)	0.1432 (14)	0.0082 (5)	−0.0026 (8)	−0.0113 (9)

Geometric parameters (Å, °)

C1—H1	0.9800	C9—N2	1.494 (8)
C1—C6	1.505 (6)	C10—H10	0.9300
C1—N1	1.479 (5)	C10—C11	1.418 (8)
C1—S1	1.811 (5)	C11—H11	0.9300

C2—C3	1.500 (5)	C12—C13	1.370 (6)
C2—N1	1.361 (5)	C12—C17	1.373 (6)
C2—O1	1.228 (5)	C12—N1	1.425 (5)
C3—H3A	0.9700	C13—H13	0.9300
C3—H3B	0.9700	C13—C14	1.402 (7)
C3—C4	1.501 (6)	C14—H14	0.9300
C4—C5	1.488 (7)	C14—C15	1.371 (8)
C4—C18	1.506 (7)	C15—H15	0.9300
C4—C19	1.530 (7)	C15—C16	1.345 (8)
C5—H5A	0.9700	C16—H16	0.9300
C5—H5B	0.9700	C16—C17	1.384 (7)
C5—S1	1.782 (6)	C17—H17	0.9300
C6—C7	1.409 (8)	C18—H18A	0.9700
C6—C11	1.366 (7)	C18—H18B	0.9700
C7—H7	0.9300	C18—C19	1.524 (7)
C7—C8	1.384 (8)	C19—H19A	0.9700
C8—H8	0.9300	C19—H19B	0.9700
C8—C9	1.362 (9)	N2—O2	1.226 (11)
C9—C10	1.342 (9)	N2—O3	1.248 (10)
C6—C1—H1	106.9	C11—C10—H10	121.1
C6—C1—S1	111.0 (3)	C6—C11—C10	120.0 (6)
N1—C1—H1	106.9	C6—C11—H11	120.0
N1—C1—C6	111.0 (3)	C10—C11—H11	120.0
N1—C1—S1	113.7 (3)	C13—C12—C17	120.0 (4)
S1—C1—H1	106.9	C13—C12—N1	119.1 (4)
N1—C2—C3	119.0 (4)	C17—C12—N1	120.8 (4)
O1—C2—C3	120.4 (4)	C12—C13—H13	120.1
O1—C2—N1	120.6 (3)	C12—C13—C14	119.8 (5)
C2—C3—H3A	108.8	C14—C13—H13	120.1
C2—C3—H3B	108.8	C13—C14—H14	120.4
C2—C3—C4	114.0 (4)	C15—C14—C13	119.1 (5)
H3A—C3—H3B	107.6	C15—C14—H14	120.4
C4—C3—H3A	108.8	C14—C15—H15	119.6
C4—C3—H3B	108.8	C16—C15—C14	120.7 (5)
C3—C4—C18	117.8 (4)	C16—C15—H15	119.6
C3—C4—C19	116.4 (4)	C15—C16—H16	119.6
C5—C4—C3	119.4 (4)	C15—C16—C17	120.7 (5)
C5—C4—C18	113.8 (4)	C17—C16—H16	119.6
C5—C4—C19	115.5 (5)	C12—C17—C16	119.6 (4)
C18—C4—C19	60.2 (3)	C12—C17—H17	120.2
C4—C5—H5A	107.9	C16—C17—H17	120.2
C4—C5—H5B	107.9	C4—C18—H18A	117.7
C4—C5—S1	117.6 (4)	C4—C18—H18B	117.7
H5A—C5—H5B	107.2	C4—C18—C19	60.6 (3)
S1—C5—H5A	107.9	H18A—C18—H18B	114.8
S1—C5—H5B	107.9	C19—C18—H18A	117.7
C7—C6—C1	117.3 (5)	C19—C18—H18B	117.7

C11—C6—C1	123.2 (5)	C4—C19—H19A	117.9
C11—C6—C7	119.5 (5)	C4—C19—H19B	117.9
C6—C7—H7	119.8	C18—C19—C4	59.1 (3)
C8—C7—C6	120.5 (6)	C18—C19—H19A	117.9
C8—C7—H7	119.8	C18—C19—H19B	117.9
C7—C8—H8	121.3	H19A—C19—H19B	115.0
C9—C8—C7	117.4 (6)	C2—N1—C1	121.6 (3)
C9—C8—H8	121.3	C2—N1—C12	119.4 (3)
C8—C9—N2	118.0 (7)	C12—N1—C1	117.3 (3)
C10—C9—C8	124.9 (6)	O2—N2—C9	117.5 (8)
C10—C9—N2	117.1 (7)	O2—N2—O3	125.6 (7)
C9—C10—H10	121.1	O3—N2—C9	116.9 (8)
C9—C10—C11	117.7 (6)	C5—S1—C1	99.4 (2)
C1—C6—C7—C8	179.8 (5)	C12—C13—C14—C15	0.6 (8)
C1—C6—C11—C10	178.7 (5)	C13—C12—C17—C16	1.3 (7)
C2—C3—C4—C5	82.2 (6)	C13—C12—N1—C1	-92.4 (5)
C2—C3—C4—C18	-132.8 (4)	C13—C12—N1—C2	102.5 (5)
C2—C3—C4—C19	-64.1 (5)	C13—C14—C15—C16	1.2 (9)
C3—C2—N1—C1	3.0 (6)	C14—C15—C16—C17	-1.7 (8)
C3—C2—N1—C12	167.4 (4)	C15—C16—C17—C12	0.5 (7)
C3—C4—C5—S1	-61.4 (7)	C17—C12—C13—C14	-1.8 (7)
C3—C4—C18—C19	106.1 (5)	C17—C12—N1—C1	84.3 (5)
C3—C4—C19—C18	-108.4 (5)	C17—C12—N1—C2	-80.7 (5)
C4—C5—S1—C1	55.8 (5)	C18—C4—C5—S1	152.2 (4)
C5—C4—C18—C19	-106.9 (5)	C19—C4—C5—S1	85.2 (5)
C5—C4—C19—C18	104.0 (5)	N1—C1—C6—C7	95.0 (5)
C6—C1—N1—C2	-160.3 (4)	N1—C1—C6—C11	-87.0 (6)
C6—C1—N1—C12	35.0 (5)	N1—C1—S1—C5	-81.7 (4)
C6—C1—S1—C5	152.3 (4)	N1—C2—C3—C4	-71.9 (5)
C6—C7—C8—C9	0.9 (8)	N1—C12—C13—C14	175.0 (4)
C7—C6—C11—C10	-3.3 (8)	N1—C12—C17—C16	-175.4 (4)
C7—C8—C9—C10	-2.1 (9)	N2—C9—C10—C11	178.0 (5)
C7—C8—C9—N2	-179.5 (5)	O1—C2—C3—C4	106.4 (5)
C8—C9—C10—C11	0.5 (8)	O1—C2—N1—C1	-175.3 (4)
C8—C9—N2—O2	-179.8 (6)	O1—C2—N1—C12	-10.9 (6)
C8—C9—N2—O3	0.2 (8)	S1—C1—C6—C7	-137.6 (4)
C9—C10—C11—C6	2.3 (8)	S1—C1—C6—C11	40.5 (6)
C10—C9—N2—O2	2.5 (8)	S1—C1—N1—C2	73.7 (4)
C10—C9—N2—O3	-177.4 (6)	S1—C1—N1—C12	-91.0 (4)
C11—C6—C7—C8	1.7 (8)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C1—H1 \cdots O1 ⁱ	0.98	2.38	3.352 (6)	173

C3—H3B···O1 ⁱ	0.97	2.48	3.444 (5)	171
--------------------------	------	------	-----------	-----

Symmetry code: (i) $-x+3/2, y, z+1/2$.