

# (Z)-2-(4-Methoxyanilino)-1,4-diphenylbut-2-ene-1,4-dione

A. J. Ravi,<sup>a</sup> M. Vinduvahini,<sup>b</sup> A. C. Vinayaka,<sup>c</sup> M. P. Sadashiva<sup>c</sup> and H. C. Devarajewda<sup>a\*</sup>

<sup>a</sup>Department of Physics, Yuvaraja's College (Constituent College), University of Mysore, Mysore 570 005, Karnataka, India, <sup>b</sup>Department of Physics, Sri D. Devaraja Urs Govt. First Grade College, Hunsur 571 105, Mysore District, Karnataka, India, and <sup>c</sup>Department of Studies in Chemistry, Manasagangotri, University of Mysore, Mysore 570 006, India. \*Correspondence e-mail: devarajewda@yahoo.com

Received 18 December 2016

Accepted 10 January 2017


Edited by K. Fejfarova, Institute of Biotechnology CAS, Czech Republic

Keywords: crystal structure; enamines; hydrogen bonding.

CCDC reference: 1526406

Structural data: full structural data are available from [iucrdata.iucr.org](http://iucrdata.iucr.org)

In the molecule of the title compound, C<sub>23</sub>H<sub>19</sub>NO<sub>3</sub>, the mean plane of the methoxyphenyl ring makes dihedral angles of 51.63 (8) and 50.86 (8)° with the terminal phenyl rings. An intramolecular N—H···O hydrogen bond occurs. The crystal structure features C—H···O hydrogen bonds.



## Structure description

$\beta$ -Enaminones containing a conjugated  $\text{—N—C=C—C=O}$  system can be used to synthesise many heterocycles: for instance, biologically important heterocycles, such as isoxazoles (Lin *et al.*, 1980), pyrroles (Yan *et al.*, 2010), indoles (Würtz *et al.*, 2008) and pyrazoles (Neumann *et al.*, 2010), have been synthesized conveniently from suitably substituted  $\beta$ -enaminones. As part of our studies in this area, the title compound was synthesized and its single-crystal structure has been reported here.

In the molecular structure of the title compound (Fig. 1), the mean plane of the methoxyphenyl ring (C6–C11) makes dihedral angles of 51.63 (8) and 50.86 (8)°, respectively, with the C14–C19 and C22–C27 phenyl rings. An intramolecular N—H···O hydrogen bond occurs. The crystal structure features C—H···O hydrogen bonds (Table 1, Fig. 2).

## Synthesis and crystallization


(Z)-2-Methylsulfanyl-1,4-diphenylbut-2-ene-1,4-dione (2.0 mmol, 1 equivalent), 4-methoxyaniline (2.6 mmol, 1.6 equivalents) and 5 volume of acidic silica with respect to starting substrate was thoroughly ground using a pestle and mortar. The solid reaction

**Table 1**  
Hydrogen-bond geometry (Å, °).


<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
N4–H4···O1	0.86	1.96	2.6346 (17)	135
C7–H7···O1 <sup>i</sup>	0.93	2.51	3.334 (2)	148
C26–H26···O2 <sup>ii</sup>	0.93	2.58	3.261 (2)	130

Symmetry codes: (i)  $-x + 1, -y + 1, -z$ ; (ii)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ .

mixture was transferred to an oven-dried 20 ml screw-cap reaction vial with magnetic stir-bar followed by the addition of anhydrous AlCl<sub>3</sub> (0.03 equivalents). The reaction mixture was stirred vigorously for 6 h. After completion of the reaction (monitored by TLC), the crude reaction mixture was purified through silica-gel column chromatography. Crystals suitable



**Figure 1**  
The molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level. H atoms are shown as spheres of arbitrary radius.



**Figure 2**  
Packing diagram of the molecule viewed parallel to the *b* axis.

**Table 2**  
Experimental details.

Crystal data	
Chemical formula	C <sub>23</sub> H <sub>19</sub> NO <sub>3</sub>
<i>M</i> <sub>r</sub>	357.39
Crystal system, space group	Monoclinic, <i>P</i> 2 <sub>1</sub> / <i>c</i>
Temperature (K)	296
<i>a</i> , <i>b</i> , <i>c</i> (Å)	10.1567 (5), 8.5396 (4), 21.8019 (10)
$\beta$ (°)	99.329 (2)
<i>V</i> (Å <sup>3</sup> )	1865.96 (15)
<i>Z</i>	4
Radiation type	Cu <i>K</i> $\alpha$
$\mu$ (mm <sup>-1</sup> )	0.68
Crystal size (mm)	0.24 × 0.20 × 0.12
Data collection	
Diffractometer	Bruker SMART CCD area-detector
Absorption correction	multi-scan ( <i>SADABS</i> ; Bruker, 2009)
<i>T</i> <sub>min</sub> , <i>T</i> <sub>max</sub>	0.770, 1.000
No. of measured, independent and observed [ <i>I</i> > 2σ( <i>I</i> )] reflections	18393, 3060, 2801
<i>R</i> <sub>int</sub>	0.048
(sin θ/λ) <sub>max</sub> (Å <sup>-1</sup> )	0.584
Refinement	
<i>R</i> [ <i>F</i> <sup>2</sup> > 2σ( <i>F</i> <sup>2</sup> )], <i>wR</i> ( <i>F</i> <sup>2</sup> ), <i>S</i>	0.050, 0.151, 1.06
No. of reflections	3060
No. of parameters	245
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}$ , $\Delta\rho_{\text{min}}$ (e Å <sup>-3</sup> )	0.23, -0.22

Computer programs: *SMART* and *SAINT* (Bruker, 2009), *SHELXS97* (Sheldrick, 2008), *SHELXL2014* (Sheldrick, 2015) and *ORTEP-3 for Windows* (Farrugia, 2012).

for analysis were obtained from an ethyl acetate–hexane (2:8 *v/v*) solution by slow evaporation at room temperature.

## Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2.

## Acknowledgements

The authors thank to Universities Sophisticated Instrumental Centre, University of Mysore, Mysuru, for CCD X-ray facilities,

## References

- Bruker (2009). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Lin, Y.-I. & Lang, S. A. (1980). *J. Org. Chem.* **45**, 4857–4860.
- Neumann, J. J., Suri, M. & Glorius, F. (2010). *Angew. Chem. Int. Ed.* **49**, 7790–7794.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Würtz, S., Rakshit, S., Neumann, J. J., Dröge, T. & Glorius, F. (2008). *Angew. Chem.* **120**, 7340–7343.
- Yan, R. L., Luo, J., Wang, C. X., Ma, C. W., Huang, G. S. & Liang, Y. M. (2010). *J. Org. Chem.* **75**, 5395–5397.

## full crystallographic data

*IUCrData* (2017). 2, x170043 [https://doi.org/10.1107/S2414314617000438]

**(Z)-2-(4-Methoxyanilino)-1,4-diphenylbut-2-ene-1,4-dione**

A. J. Ravi, M. Vinduvahini, A. C. Vinayaka, M. P. Sadashiva and H. C. Devarajegowda

**(Z)-2-(4-Methoxyanilino)-1,4-diphenylbut-2-ene-1,4-dione***Crystal data*

$C_{23}H_{19}NO_3$	$D_x = 1.272 \text{ Mg m}^{-3}$
$M_r = 357.39$	Melting point: 375 K
Monoclinic, $P2_1/c$	Cu $K\alpha$ radiation, $\lambda = 1.54178 \text{ \AA}$
$a = 10.1567 (5) \text{ \AA}$	Cell parameters from 3060 reflections
$b = 8.5396 (4) \text{ \AA}$	$\theta = 4.1\text{--}64.3^\circ$
$c = 21.8019 (10) \text{ \AA}$	$\mu = 0.68 \text{ mm}^{-1}$
$\beta = 99.329 (2)^\circ$	$T = 296 \text{ K}$
$V = 1865.96 (15) \text{ \AA}^3$	Prism, colourless
$Z = 4$	$0.24 \times 0.20 \times 0.12 \text{ mm}$
$F(000) = 752$	

*Data collection*

Bruker SMART CCD area-detector diffractometer	18393 measured reflections
Radiation source: fine-focus sealed tube	3060 independent reflections
Graphite monochromator	2801 reflections with $I > 2\sigma(I)$
$\omega$ and $\varphi$ scans	$R_{\text{int}} = 0.048$
Absorption correction: multi-scan (SADABS; Bruker, 2009)	$\theta_{\text{max}} = 64.3^\circ$ , $\theta_{\text{min}} = 4.1^\circ$
$T_{\text{min}} = 0.770$ , $T_{\text{max}} = 1.000$	$h = -11 \rightarrow 11$
	$k = -9 \rightarrow 8$
	$l = -25 \rightarrow 25$

*Refinement*

Refinement on $F^2$	H-atom parameters constrained
Least-squares matrix: full	$w = 1/[\sigma^2(F_o^2) + (0.1038P)^2 + 0.2139P]$
$R[F^2 > 2\sigma(F^2)] = 0.050$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.151$	$(\Delta/\sigma)_{\text{max}} < 0.001$
$S = 1.06$	$\Delta\rho_{\text{max}} = 0.23 \text{ e \AA}^{-3}$
3060 reflections	$\Delta\rho_{\text{min}} = -0.22 \text{ e \AA}^{-3}$
245 parameters	Extinction correction: SHELXL2014 (Sheldrick 2015), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
0 restraints	Extinction coefficient: 0.0053 (10)
Hydrogen site location: inferred from neighbouring sites	

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.64804 (12)	0.77341 (13)	0.62059 (5)	0.0584 (3)
O2	0.87120 (11)	0.58731 (12)	0.44792 (5)	0.0543 (3)
O3	0.37864 (16)	0.77684 (17)	0.26306 (6)	0.0781 (4)
N4	0.63382 (13)	0.67640 (14)	0.50499 (6)	0.0478 (3)
H4	0.5974	0.7271	0.5319	0.057*
C5	0.3437 (3)	0.6556 (3)	0.21998 (10)	0.1011 (8)
H5A	0.3013	0.6990	0.1812	0.152*
H5B	0.4226	0.5999	0.2138	0.152*
H5C	0.2834	0.5849	0.2355	0.152*
C6	0.44064 (16)	0.7394 (2)	0.32159 (7)	0.0530 (4)
C7	0.46065 (16)	0.86315 (19)	0.36271 (7)	0.0542 (4)
H7	0.4300	0.9624	0.3500	0.065*
C8	0.52575 (15)	0.84044 (17)	0.42243 (7)	0.0489 (4)
H8	0.5399	0.9248	0.4497	0.059*
C9	0.57053 (14)	0.69229 (17)	0.44240 (7)	0.0427 (4)
C10	0.54586 (14)	0.56800 (17)	0.40179 (7)	0.0488 (4)
H10	0.5718	0.4677	0.4152	0.059*
C11	0.48277 (16)	0.59083 (18)	0.34112 (7)	0.0536 (4)
H11	0.4689	0.5068	0.3137	0.064*
C12	0.74285 (14)	0.59305 (15)	0.52800 (6)	0.0414 (4)
C13	0.81574 (13)	0.50796 (16)	0.48246 (6)	0.0412 (4)
C14	0.81966 (13)	0.33465 (16)	0.48181 (6)	0.0423 (4)
C15	0.87236 (17)	0.2625 (2)	0.43389 (8)	0.0554 (4)
H15	0.9038	0.3232	0.4039	0.066*
C16	0.87819 (19)	0.1019 (2)	0.43071 (9)	0.0671 (5)
H16	0.9121	0.0544	0.3982	0.080*
C17	0.83407 (18)	0.0114 (2)	0.47545 (10)	0.0689 (5)
H17	0.8398	-0.0971	0.4736	0.083*
C18	0.78127 (18)	0.0815 (2)	0.52307 (9)	0.0642 (5)
H18	0.7515	0.0199	0.5532	0.077*
C19	0.77249 (15)	0.24275 (18)	0.52614 (7)	0.0503 (4)
H19	0.7351	0.2896	0.5578	0.060*
C20	0.79907 (15)	0.59469 (16)	0.58952 (6)	0.0455 (4)
H20	0.8718	0.5298	0.6026	0.055*
C21	0.75096 (16)	0.69174 (16)	0.63469 (7)	0.0461 (4)
C22	0.82350 (16)	0.69738 (17)	0.70020 (7)	0.0477 (4)
C23	0.94938 (19)	0.6344 (2)	0.71859 (8)	0.0632 (5)
H23	0.9919	0.5833	0.6896	0.076*
C24	1.0125 (2)	0.6467 (3)	0.77957 (9)	0.0729 (5)
H24	1.0977	0.6056	0.7913	0.088*
C25	0.9490 (2)	0.7201 (2)	0.82313 (8)	0.0651 (5)
H25	0.9910	0.7279	0.8642	0.078*
C26	0.8240 (2)	0.7811 (2)	0.80563 (8)	0.0709 (5)
H26	0.7808	0.8296	0.8350	0.085*
C27	0.7618 (2)	0.7709 (2)	0.74466 (8)	0.0632 (5)

H27                    0.6773                    0.8140                    0.7332                    0.076\*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0743 (8)	0.0535 (6)	0.0464 (6)	0.0173 (5)	0.0065 (5)	-0.0035 (5)
O2	0.0662 (7)	0.0557 (7)	0.0433 (6)	-0.0025 (5)	0.0162 (5)	0.0092 (5)
O3	0.0962 (10)	0.0822 (9)	0.0477 (7)	0.0060 (7)	-0.0130 (6)	0.0084 (6)
N4	0.0566 (7)	0.0490 (7)	0.0376 (7)	0.0071 (5)	0.0072 (5)	-0.0003 (5)
C5	0.1192 (19)	0.115 (2)	0.0568 (12)	0.0020 (15)	-0.0224 (12)	-0.0074 (12)
C6	0.0529 (8)	0.0607 (10)	0.0437 (8)	0.0016 (7)	0.0023 (7)	0.0077 (7)
C7	0.0605 (9)	0.0458 (8)	0.0551 (9)	0.0084 (7)	0.0059 (7)	0.0102 (7)
C8	0.0550 (8)	0.0419 (8)	0.0490 (9)	0.0034 (6)	0.0065 (6)	-0.0023 (6)
C9	0.0452 (7)	0.0441 (8)	0.0388 (7)	0.0002 (6)	0.0066 (6)	0.0030 (6)
C10	0.0539 (8)	0.0401 (8)	0.0500 (9)	0.0001 (6)	0.0010 (6)	0.0014 (6)
C11	0.0570 (9)	0.0516 (9)	0.0495 (9)	-0.0013 (7)	0.0005 (7)	-0.0077 (7)
C12	0.0502 (8)	0.0363 (7)	0.0379 (7)	-0.0023 (5)	0.0074 (6)	0.0026 (5)
C13	0.0451 (7)	0.0460 (8)	0.0313 (7)	-0.0020 (6)	0.0021 (5)	0.0017 (5)
C14	0.0411 (7)	0.0457 (8)	0.0380 (7)	0.0002 (5)	0.0006 (6)	-0.0029 (6)
C15	0.0570 (9)	0.0612 (10)	0.0483 (9)	0.0011 (7)	0.0098 (7)	-0.0084 (7)
C16	0.0661 (10)	0.0616 (11)	0.0727 (12)	0.0084 (8)	0.0087 (9)	-0.0224 (9)
C17	0.0610 (10)	0.0450 (9)	0.0956 (14)	0.0039 (7)	-0.0023 (9)	-0.0125 (9)
C18	0.0670 (10)	0.0454 (9)	0.0779 (12)	-0.0060 (7)	0.0053 (9)	0.0073 (8)
C19	0.0527 (8)	0.0468 (8)	0.0516 (9)	-0.0031 (6)	0.0088 (7)	0.0001 (6)
C20	0.0556 (8)	0.0439 (8)	0.0368 (7)	0.0047 (6)	0.0067 (6)	0.0005 (6)
C21	0.0606 (9)	0.0379 (7)	0.0402 (8)	0.0014 (6)	0.0091 (6)	0.0021 (6)
C22	0.0640 (9)	0.0410 (8)	0.0385 (8)	-0.0022 (6)	0.0094 (7)	0.0011 (6)
C23	0.0714 (11)	0.0756 (11)	0.0427 (9)	0.0093 (9)	0.0094 (7)	-0.0030 (8)
C24	0.0747 (11)	0.0884 (14)	0.0515 (10)	0.0096 (10)	-0.0023 (8)	0.0019 (9)
C25	0.0878 (13)	0.0673 (11)	0.0378 (8)	-0.0092 (9)	0.0031 (8)	-0.0025 (7)
C26	0.0919 (14)	0.0795 (12)	0.0417 (9)	0.0064 (10)	0.0122 (9)	-0.0135 (8)
C27	0.0756 (11)	0.0681 (11)	0.0458 (9)	0.0108 (8)	0.0095 (8)	-0.0095 (8)

*Geometric parameters (Å, °)*

O1—C21	1.253 (2)	C14—C15	1.393 (2)
O2—C13	1.2170 (16)	C15—C16	1.374 (3)
O3—C6	1.366 (2)	C15—H15	0.9300
O3—C5	1.405 (3)	C16—C17	1.375 (3)
N4—C12	1.3433 (19)	C16—H16	0.9300
N4—C9	1.4173 (19)	C17—C18	1.380 (3)
N4—H4	0.8600	C17—H17	0.9300
C5—H5A	0.9600	C18—C19	1.383 (2)
C5—H5B	0.9600	C18—H18	0.9300
C5—H5C	0.9600	C19—H19	0.9300
C6—C7	1.380 (2)	C20—C21	1.432 (2)
C6—C11	1.384 (2)	C20—H20	0.9300
C7—C8	1.375 (2)	C21—C22	1.498 (2)

C7—H7	0.9300	C22—C23	1.385 (2)
C8—C9	1.390 (2)	C22—C27	1.387 (2)
C8—H8	0.9300	C23—C24	1.383 (2)
C9—C10	1.379 (2)	C23—H23	0.9300
C10—C11	1.387 (2)	C24—C25	1.382 (3)
C10—H10	0.9300	C24—H24	0.9300
C11—H11	0.9300	C25—C26	1.368 (3)
C12—C20	1.3701 (19)	C25—H25	0.9300
C12—C13	1.5166 (19)	C26—C27	1.379 (3)
C13—C14	1.481 (2)	C26—H26	0.9300
C14—C19	1.389 (2)	C27—H27	0.9300
C6—O3—C5	118.70 (16)	C16—C15—H15	119.9
C12—N4—C9	128.91 (12)	C14—C15—H15	119.9
C12—N4—H4	115.5	C15—C16—C17	120.21 (17)
C9—N4—H4	115.5	C15—C16—H16	119.9
O3—C5—H5A	109.5	C17—C16—H16	119.9
O3—C5—H5B	109.5	C16—C17—C18	120.08 (16)
H5A—C5—H5B	109.5	C16—C17—H17	120.0
O3—C5—H5C	109.5	C18—C17—H17	120.0
H5A—C5—H5C	109.5	C17—C18—C19	120.28 (16)
H5B—C5—H5C	109.5	C17—C18—H18	119.9
O3—C6—C7	115.13 (14)	C19—C18—H18	119.9
O3—C6—C11	125.08 (16)	C18—C19—C14	119.80 (15)
C7—C6—C11	119.79 (14)	C18—C19—H19	120.1
C8—C7—C6	120.32 (14)	C14—C19—H19	120.1
C8—C7—H7	119.8	C12—C20—C21	122.99 (13)
C6—C7—H7	119.8	C12—C20—H20	118.5
C7—C8—C9	120.51 (14)	C21—C20—H20	118.5
C7—C8—H8	119.7	O1—C21—C20	121.16 (14)
C9—C8—H8	119.7	O1—C21—C22	118.56 (13)
C10—C9—C8	118.89 (14)	C20—C21—C22	120.28 (13)
C10—C9—N4	123.46 (13)	C23—C22—C27	118.28 (15)
C8—C9—N4	117.58 (13)	C23—C22—C21	123.49 (14)
C9—C10—C11	120.80 (14)	C27—C22—C21	118.22 (15)
C9—C10—H10	119.6	C24—C23—C22	120.73 (16)
C11—C10—H10	119.6	C24—C23—H23	119.6
C6—C11—C10	119.63 (14)	C22—C23—H23	119.6
C6—C11—H11	120.2	C25—C24—C23	119.99 (18)
C10—C11—H11	120.2	C25—C24—H24	120.0
N4—C12—C20	123.21 (13)	C23—C24—H24	120.0
N4—C12—C13	118.02 (12)	C26—C25—C24	119.77 (16)
C20—C12—C13	118.40 (12)	C26—C25—H25	120.1
O2—C13—C14	122.32 (12)	C24—C25—H25	120.1
O2—C13—C12	117.54 (12)	C25—C26—C27	120.29 (17)
C14—C13—C12	120.13 (11)	C25—C26—H26	119.9
C19—C14—C15	119.33 (14)	C27—C26—H26	119.9
C19—C14—C13	123.03 (13)	C26—C27—C22	120.94 (17)

C15—C14—C13	117.64 (13)	C26—C27—H27	119.5
C16—C15—C14	120.28 (16)	C22—C27—H27	119.5
C5—O3—C6—C7	-174.3 (2)	C19—C14—C15—C16	-0.3 (2)
C5—O3—C6—C11	5.7 (3)	C13—C14—C15—C16	-179.77 (14)
O3—C6—C7—C8	-178.02 (15)	C14—C15—C16—C17	-1.1 (3)
C11—C6—C7—C8	1.9 (2)	C15—C16—C17—C18	1.3 (3)
C6—C7—C8—C9	-0.9 (2)	C16—C17—C18—C19	-0.1 (3)
C7—C8—C9—C10	-1.5 (2)	C17—C18—C19—C14	-1.4 (2)
C7—C8—C9—N4	-178.42 (13)	C15—C14—C19—C18	1.5 (2)
C12—N4—C9—C10	44.7 (2)	C13—C14—C19—C18	-179.03 (14)
C12—N4—C9—C8	-138.48 (15)	N4—C12—C20—C21	-3.3 (2)
C8—C9—C10—C11	2.9 (2)	C13—C12—C20—C21	169.52 (13)
N4—C9—C10—C11	179.62 (13)	C12—C20—C21—O1	4.8 (2)
O3—C6—C11—C10	179.39 (15)	C12—C20—C21—C22	-175.16 (13)
C7—C6—C11—C10	-0.6 (2)	O1—C21—C22—C23	-168.49 (17)
C9—C10—C11—C6	-1.9 (2)	C20—C21—C22—C23	11.5 (2)
C9—N4—C12—C20	176.00 (14)	O1—C21—C22—C27	10.9 (2)
C9—N4—C12—C13	3.1 (2)	C20—C21—C22—C27	-169.14 (15)
N4—C12—C13—O2	66.90 (17)	C27—C22—C23—C24	-0.9 (3)
C20—C12—C13—O2	-106.31 (15)	C21—C22—C23—C24	178.44 (17)
N4—C12—C13—C14	-114.13 (14)	C22—C23—C24—C25	1.1 (3)
C20—C12—C13—C14	72.67 (17)	C23—C24—C25—C26	-0.4 (3)
O2—C13—C14—C19	171.17 (13)	C24—C25—C26—C27	-0.6 (3)
C12—C13—C14—C19	-7.8 (2)	C25—C26—C27—C22	0.8 (3)
O2—C13—C14—C15	-9.4 (2)	C23—C22—C27—C26	-0.1 (3)
C12—C13—C14—C15	171.68 (12)	C21—C22—C27—C26	-179.45 (16)

## Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N4—H4...O1	0.86	1.96	2.6346 (17)	135
C7—H7...O1 <sup>i</sup>	0.93	2.51	3.334 (2)	148
C26—H26...O2 <sup>ii</sup>	0.93	2.58	3.261 (2)	130

Symmetry codes: (i)  $-x+1, -y+1, -z$ ; (ii)  $x, -y+1/2, z-1/2$ .