

Dicarbonyl-1 κ^2 C- μ -chlorido-2:3 κ^2 Cl:Cl-pentachlorido-2 κ^2 Cl,3 κ^3 Cl-[1(η^6)-toluene]-digallium(III)ruthenium(I)(Ru—Ga)

Danielle Smith and George N. Harakas*

PO Box 6949, Radford University, Radford, Virginia 24142, USA. *Correspondence e-mail: gharakas@radford.edu


Received 2 July 2024

Accepted 3 July 2024

Edited by M. Bolte, Goethe-Universität Frankfurt, Germany

Keywords: crystal structure; ruthenium; gallium; metal–organic.**CCDC reference:** 2367755**Structural data:** full structural data are available from iucrdata.iucr.org

The title compound, $[\text{RuGa}_2\text{Cl}_6(\text{C}_7\text{H}_8)(\text{CO})_2]$ or $[(\text{CO})_2(\text{GaCl}_2)(\eta^6\text{-toluene})\text{Ru}]^+[\text{GaCl}_4]^-$, was isolated from the reaction of Ga_2Cl_4 with diphenylsilanediol in toluene, followed by the addition of $\text{Ru}_3(\text{CO})_{12}$. The compound contains a ruthenium–gallium metal–metal bond with a length of 2.4575 (2) Å.



Structure description

The reaction of Ga_2Cl_4 with $\text{Ru}_3(\text{CO})_{12}$ in toluene was demonstrated to produce two ruthenium–gallium metal clusters (Harakas & Whittlesey, 1997). The title compound (Fig. 1) was isolated during an attempt to synthesize the ruthenium–gallium diphenylsiloxane analogue of the previously reported iron–gallium dimethylsiloxane cluster (Demmin *et al.*, 2024).

The title compound can be described as $[(\text{CO})_2(\text{GaCl}_2)(\eta^6\text{-toluene})\text{Ru}]^+[\text{GaCl}_4]^-$. A single positive charge on the ruthenium complex provides a total of 18 electrons for the metal center with $[\text{GaCl}_4]^-$ for charge balance. This bonding model is supported by the $\text{Ga1}-\text{Cl3}$ bond length of 2.4619 (5) Å, which is significantly longer than the other $\text{Ga}-\text{Cl}$ bond lengths (Table 1) observed in the title compound. In contrast, $[(\text{CpFe}(\text{CO})_2)\text{Ga}(\text{Cl}^*\text{GaCl}_3)(\mu\text{-Cl})_2]$ is described as a Lewis acid–base complex *i.e.* $[(\text{CpFe}(\text{CO})_2)\text{GaCl}_2]_2\cdot 2\text{GaCl}_3$ (Borovik *et al.*, 1999). The bond angles for the terminal GaCl_3 are 112.87, 114.74, and 114.09°, which are all significantly greater than the 109.5° of tetrahedral geometry. For the title compound, the corresponding angles around Ga2 (Table 1) are much closer to the ideal tetrahedral geometry, which is consistent with $[\text{GaCl}_4]^-$. In tetraethyl ammonium tetrachloridogallium (Bolte *et al.*, 2023), the $\text{Cl}-\text{Ga}-\text{Cl}$ bond angles range from 108.1 to 110.1°. An analogous Lewis acid–base bonding model for the title complex would require a 19 electron ruthenium metal center or that Ga1 carries one formal negative charge, both of which are unlikely.




Figure 1
The molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level.

The Ru1–Ga1 bond length of 2.4575 (2) Å for the title compound is very similar to the value of 2.453 (1) Å observed for Ru₂{GaCl₂(THF)}₂(CO)₈ (Harakas & Whittlesey, 1997). The packing is shown in Fig. 2

During the work-up of the reaction, the title compound was isolated directly from the toluene solution. It is unknown at this time the role of diphenylsilanediol, if any, in the formation of the title compound. A solid that was insoluble in toluene in the reaction flask was extracted with THF forming an orange solution. This reaction product, which may contain the desired diphenylsiloxane metal cluster, has not yet been fully characterized.

Synthesis and crystallization

All manipulations were carried out under argon using standard Schlenk line techniques. Our previous work (Demmin *et al.*, 2024) demonstrated that silicone-based vacuum grease can contaminate gallium halide reactions. Therefore, PTFE




Figure 2
Crystal packing diagram viewed along the *b* axis. Hydrogen atoms have been omitted for clarity.

Table 1
Selected geometric parameters (Å, °).

Ru1–Ga1	2.4575 (2)	Ga2–Cl6	2.1413 (6)
Ga1–Cl2	2.1665 (5)	Ga2–Cl4	2.1456 (6)
Ga1–Cl1	2.1888 (5)	Ga2–Cl5	2.1521 (6)
Ga1–Cl3	2.4619 (5)	Ga2–Cl3	2.2583 (5)
Cl6–Ga2–Cl4	114.95 (3)	Cl6–Ga2–Cl3	110.96 (2)
Cl6–Ga2–Cl5	110.36 (3)	Cl4–Ga2–Cl3	106.43 (2)
Cl4–Ga2–Cl5	109.67 (3)	Cl5–Ga2–Cl3	103.82 (2)

Table 2
Experimental details.

Crystal data	[RuGa ₂ Cl ₆ (C ₇ H ₈)(CO) ₂]
Chemical formula	601.36
<i>M_r</i>	601.36
Crystal system, space group	Monoclinic, <i>P</i> ₂ / <i>c</i>
Temperature (K)	248
<i>a</i> , <i>b</i> , <i>c</i> (Å)	13.1598 (6), 9.7142 (4), 15.3369 (7)
β (°)	115.257 (1)
<i>V</i> (Å ³)	1773.19 (14)
<i>Z</i>	4
Radiation type	Mo <i>K</i> α
μ (mm ^{−1})	4.76
Crystal size (mm)	0.36 × 0.34 × 0.31
Data collection	
Diffractometer	Bruker D8 Quest Eco, Photon II 7
Absorption correction	Multi-scan (<i>SADABS</i> ; Krause <i>et al.</i> , 2015)
<i>T_{min}</i> , <i>T_{max}</i>	0.21, 0.32
No. of measured, independent and observed [<i>I</i> > 2σ(<i>I</i>)] reflections	66270, 4424, 4133
<i>R_{int}</i>	0.031
(sin θ/λ) _{max} (Å ^{−1})	0.668
Refinement	
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.016, 0.041, 1.09
No. of reflections	4424
No. of parameters	182
H-atom treatment	H-atom parameters constrained
Δρ _{max} , Δρ _{min} (e Å ^{−3})	0.65, −0.43

Computer programs: *APEX3* and *SAINT* (Bruker, 2019), *SHELXT2018/2* (Sheldrick, 2015a), *SHELXL2018/3* (Sheldrick, 2015b) and *ShelXle* (Hübschle *et al.*, 2011).

sleeves and non-silicone based vacuum grease were used on all glassware in this experiment. In a 250 ml Schlenk flask, gallium (5.60 g, 80.3 mmole) and GaCl₃ 5.00 g (28.4, mmole) were combined followed by toluene (175 ml). The mixture was heated to reflux for 24 h to produce a solution containing gallium(II) chloride (Ga₂Cl₄), excess gallium was present.

Diphenylsilanediol (0.306 g, 1.41 mmol) was added to a 150 ml Schlenk flask followed by toluene (50 ml). To this flask, 10 ml of the Ga₂Cl₄ stock solution was added *via* cannula. The dark-gray mixture was refluxed under argon for 72 h resulting in a light-gray mixture. The reaction flask was cooled to 25°C and Ru₃(CO)₁₂ (0.225 g, 0.352 mmol) was then added. The mixture was heated to reflux for an additional 72 h. This resulted in a mixture with a suspended gray solid/gel and colorless solution. The colorless solution was decanted into a 150 ml Schlenk flask *via* cannula. After standing at 25°C for 10 days, colorless crystals were observed.

A single crystal was coated with NVH oil and mounted on a MiTeGen loop under a stream of argon gas then cooled to −25°C for data collection.

Refinement

Crystal data, data collection, and structure refinement details are summarized in Table 2.

References

- Bolte, M., Georg, I. & Wagner, M. (2023). *CSD Communication* (CCDC 2250783, Refcode VIDYAM). CCDC, Cambridge, England.
- Borovik, A. S., Bott, S. G. & Barron, A. R. (1999). *Organometallics*, **18**, 2668–2676.
- Bruker (2019). *APEX3* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Demmin, M. E., Bauer, C., Ruf, M. & Harakas, G. N. (2024). *IUCr-Data*, **9**, x240620.
- Harakas, G. N. & Whittlesey, B. R. (1997). *Inorg. Chem.* **36**, 2704–2707.
- Hübschle, C. B., Sheldrick, G. M. & Dittrich, B. (2011). *J. Appl. Cryst.* **44**, 1281–1284.
- Krause, L., Herbst-Irmer, R., Sheldrick, G. M. & Stalke, D. (2015). *J. Appl. Cryst.* **48**, 3–10.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.

full crystallographic data

IUCrData (2024). **9**, x240657 [https://doi.org/10.1107/S2414314624006576]

Dicarbonyl-1 κ^2 C- μ -chlorido-2:3 κ^2 Cl:Cl-pentachlorido-2 κ^2 Cl,3 κ^3 Cl-[1(η^6)-toluene]digallium(III)ruthenium(I)(Ru—Ga)

Danielle Smith and George N. Harakas

Dicarbonyl-1 κ^2 C- μ -chlorido-2:3 κ^2 Cl:Cl-pentachlorido-2 κ^2 Cl,3 κ^3 Cl-[1(η^6)-toluene]digallium(III)ruthenium(I)(Ru—Ga)

Crystal data

[RuGa₂Cl₆(C₇H₈)(CO)₂]

$M_r = 601.36$

Monoclinic, $P2_1/c$

$a = 13.1598$ (6) Å

$b = 9.7142$ (4) Å

$c = 15.3369$ (7) Å

$\beta = 115.257$ (1)°

$V = 1773.19$ (14) Å³

$Z = 4$

$F(000) = 1144$

$D_x = 2.253$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9595 reflections

$\theta = 2.7$ – 28.3 °

$\mu = 4.76$ mm⁻¹

$T = 248$ K

Block, clear colourless

$0.36 \times 0.34 \times 0.31$ mm

Data collection

Bruker D8 Quest Eco, Photon II 7
diffractometer

Detector resolution: 7.3910 pixels mm⁻¹

phi and ω scans

Absorption correction: multi-scan
(SADABS; Krause *et al.*, 2015)

$T_{\min} = 0.21$, $T_{\max} = 0.32$

66270 measured reflections

4424 independent reflections

4133 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\max} = 28.3$ °, $\theta_{\min} = 2.7$ °

$h = -17$ → 17

$k = -12$ → 12

$l = -20$ → 20

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.016$

$wR(F^2) = 0.041$

$S = 1.09$

4424 reflections

182 parameters

0 restraints

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0179P)^2 + 1.0566P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.002$

$\Delta\rho_{\max} = 0.65$ e Å⁻³

$\Delta\rho_{\min} = -0.43$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ru1	0.24301 (2)	0.66776 (2)	0.55005 (2)	0.02484 (4)
Ga1	0.37587 (2)	0.52947 (2)	0.68455 (2)	0.02795 (5)
Ga2	0.21861 (2)	0.26035 (2)	0.77462 (2)	0.03228 (5)
Cl1	0.53434 (4)	0.47105 (6)	0.68030 (4)	0.04053 (11)
Cl2	0.41255 (5)	0.58077 (6)	0.83247 (3)	0.04424 (12)
Cl3	0.31993 (4)	0.28855 (5)	0.68851 (4)	0.03697 (10)
Cl4	0.09559 (5)	0.42180 (6)	0.73078 (5)	0.05227 (13)
Cl5	0.13673 (5)	0.06499 (6)	0.72620 (5)	0.05478 (14)
Cl6	0.32572 (4)	0.25357 (7)	0.92593 (4)	0.05277 (15)
O1	0.12790 (15)	0.73771 (17)	0.67684 (13)	0.0519 (4)
O2	0.10336 (14)	0.40984 (18)	0.48352 (13)	0.0573 (4)
C1	0.17226 (16)	0.7071 (2)	0.63130 (14)	0.0338 (4)
C2	0.15713 (16)	0.5047 (2)	0.51002 (14)	0.0367 (4)
C3	0.1370 (2)	1.0026 (3)	0.48306 (18)	0.0502 (5)
H3A	0.151190	1.085031	0.454310	0.075000*
H3B	0.148544	1.022284	0.548647	0.075000*
H3C	0.060080	0.972800	0.445748	0.075000*
C4	0.21607 (15)	0.8907 (2)	0.48377 (13)	0.0324 (4)
C5	0.18667 (17)	0.7958 (2)	0.40924 (13)	0.0389 (4)
H5	0.107892	0.790389	0.359964	0.047000*
C6	0.2656 (2)	0.6935 (3)	0.41157 (15)	0.0478 (5)
H6	0.241232	0.618942	0.362932	0.057000*
C7	0.37233 (19)	0.6881 (2)	0.48817 (17)	0.0458 (5)
H7	0.423488	0.610611	0.493398	0.055000*
C8	0.40090 (17)	0.7834 (2)	0.56321 (16)	0.0432 (5)
H8	0.471964	0.772111	0.622005	0.052000*
C9	0.32405 (16)	0.8811 (2)	0.56150 (14)	0.0371 (4)
H9	0.340902	0.937264	0.619916	0.044000*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ru1	0.02522 (6)	0.02903 (7)	0.02071 (6)	−0.00104 (5)	0.01024 (5)	0.00055 (5)
Ga1	0.02861 (9)	0.03056 (10)	0.02508 (9)	0.00314 (7)	0.01183 (7)	0.00206 (7)
Ga2	0.03015 (10)	0.03326 (11)	0.03336 (10)	−0.00027 (8)	0.01347 (8)	0.00417 (8)
Cl1	0.0333 (2)	0.0472 (3)	0.0458 (3)	0.00639 (19)	0.0213 (2)	0.0039 (2)
Cl2	0.0559 (3)	0.0468 (3)	0.0281 (2)	0.0091 (2)	0.0161 (2)	−0.00377 (19)
Cl3	0.0476 (2)	0.0293 (2)	0.0424 (2)	−0.00163 (19)	0.0273 (2)	0.00116 (18)
Cl4	0.0467 (3)	0.0490 (3)	0.0610 (3)	0.0157 (2)	0.0228 (3)	0.0080 (3)
Cl5	0.0554 (3)	0.0405 (3)	0.0623 (3)	−0.0155 (2)	0.0192 (3)	0.0040 (3)
Cl6	0.0389 (2)	0.0864 (4)	0.0323 (2)	0.0041 (3)	0.0145 (2)	0.0060 (3)
O1	0.0721 (11)	0.0459 (9)	0.0607 (10)	0.0051 (8)	0.0503 (9)	0.0003 (8)
O2	0.0500 (9)	0.0449 (9)	0.0626 (10)	−0.0152 (8)	0.0103 (8)	−0.0092 (8)
C1	0.0408 (9)	0.0295 (9)	0.0349 (9)	0.0002 (7)	0.0198 (8)	0.0036 (7)
C2	0.0334 (9)	0.0387 (10)	0.0334 (9)	−0.0019 (8)	0.0099 (7)	−0.0010 (8)

C3	0.0545 (13)	0.0464 (12)	0.0567 (14)	0.0116 (10)	0.0304 (11)	0.0103 (10)
C4	0.0365 (9)	0.0339 (9)	0.0302 (8)	-0.0027 (7)	0.0175 (7)	0.0051 (7)
C5	0.0422 (10)	0.0474 (11)	0.0233 (8)	-0.0055 (9)	0.0101 (7)	0.0060 (8)
C6	0.0754 (15)	0.0475 (12)	0.0321 (10)	-0.0045 (11)	0.0342 (11)	-0.0027 (9)
C7	0.0519 (12)	0.0497 (12)	0.0532 (12)	0.0122 (10)	0.0392 (11)	0.0147 (10)
C8	0.0308 (9)	0.0520 (12)	0.0473 (11)	-0.0053 (9)	0.0172 (8)	0.0101 (10)
C9	0.0371 (9)	0.0378 (10)	0.0343 (9)	-0.0111 (8)	0.0133 (8)	0.0000 (8)

Geometric parameters (Å, °)

Ru1—C1	1.8856 (19)	Ga2—C14	2.1456 (6)
Ru1—C2	1.890 (2)	Ga2—C15	2.1521 (6)
Ru1—C6	2.2803 (19)	Ga2—C13	2.2583 (5)
Ru1—C7	2.2834 (19)	O1—C1	1.125 (2)
Ru1—C8	2.294 (2)	O2—C2	1.127 (3)
Ru1—C9	2.3027 (19)	C3—C4	1.502 (3)
Ru1—C5	2.3223 (19)	C4—C5	1.389 (3)
Ru1—C4	2.3536 (19)	C4—C9	1.415 (3)
Ru1—Ga1	2.4575 (2)	C5—C6	1.426 (3)
Ga1—Cl2	2.1665 (5)	C6—C7	1.396 (3)
Ga1—Cl1	2.1888 (5)	C7—C8	1.398 (3)
Ga1—Cl3	2.4619 (5)	C8—C9	1.379 (3)
Ga2—Cl6	2.1413 (6)		
C1—Ru1—C2	89.37 (9)	Cl2—Ga1—Cl1	107.54 (2)
C1—Ru1—C6	152.94 (9)	Cl2—Ga1—Ru1	120.934 (17)
C2—Ru1—C6	94.99 (9)	Cl1—Ga1—Ru1	117.546 (16)
C1—Ru1—C7	157.35 (9)	Cl2—Ga1—Cl3	97.36 (2)
C2—Ru1—C7	112.70 (9)	Cl1—Ga1—Cl3	93.035 (19)
C6—Ru1—C7	35.62 (9)	Ru1—Ga1—Cl3	115.292 (15)
C1—Ru1—C8	121.78 (9)	Cl6—Ga2—C14	114.95 (3)
C2—Ru1—C8	146.39 (9)	Cl6—Ga2—C15	110.36 (3)
C6—Ru1—C8	63.59 (9)	Cl4—Ga2—C15	109.67 (3)
C7—Ru1—C8	35.56 (9)	Cl6—Ga2—Cl3	110.96 (2)
C1—Ru1—C9	96.35 (8)	Cl4—Ga2—Cl3	106.43 (2)
C2—Ru1—C9	166.51 (8)	Cl5—Ga2—Cl3	103.82 (2)
C6—Ru1—C9	74.46 (8)	Ga2—Cl3—Ga1	112.77 (2)
C7—Ru1—C9	63.31 (8)	O1—C1—Ru1	175.86 (18)
C8—Ru1—C9	34.91 (8)	O2—C2—Ru1	177.7 (2)
C1—Ru1—C5	117.03 (8)	C5—C4—C9	118.47 (18)
C2—Ru1—C5	103.73 (8)	C5—C4—C3	121.72 (19)
C6—Ru1—C5	36.09 (8)	C9—C4—C3	119.81 (19)
C7—Ru1—C5	64.28 (8)	C5—C4—Ru1	71.49 (11)
C8—Ru1—C5	74.78 (7)	C9—C4—Ru1	70.35 (11)
C9—Ru1—C5	62.79 (7)	C3—C4—Ru1	130.19 (13)
C1—Ru1—C4	94.20 (7)	C4—C5—C6	119.83 (19)
C2—Ru1—C4	132.21 (8)	C4—C5—Ru1	73.96 (11)
C6—Ru1—C4	63.40 (8)	C6—C5—Ru1	70.35 (11)

C7—Ru1—C4	75.08 (7)	C7—C6—C5	120.5 (2)
C8—Ru1—C4	63.34 (7)	C7—C6—Ru1	72.31 (11)
C9—Ru1—C4	35.37 (7)	C5—C6—Ru1	73.56 (11)
C5—Ru1—C4	34.55 (7)	C6—C7—C8	119.2 (2)
C1—Ru1—Ga1	86.05 (6)	C6—C7—Ru1	72.07 (12)
C2—Ru1—Ga1	85.82 (6)	C8—C7—Ru1	72.63 (11)
C6—Ru1—Ga1	120.86 (7)	C9—C8—C7	120.2 (2)
C7—Ru1—Ga1	90.38 (6)	C9—C8—Ru1	72.89 (11)
C8—Ru1—Ga1	84.29 (6)	C7—C8—Ru1	71.81 (12)
C9—Ru1—Ga1	106.70 (5)	C8—C9—C4	121.73 (19)
C5—Ru1—Ga1	154.66 (5)	C8—C9—Ru1	72.20 (12)
C4—Ru1—Ga1	141.95 (5)	C4—C9—Ru1	74.28 (11)
